





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 879-4503

1.5
1.6
1.8
2.0
2.2
2.5
2.8
3.2
3.6
4.0
4.5

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

1.5
1.6
1.8
2.0
2.2
2.5
2.8
3.2
3.6
4.0
4.5

© 1981

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- | | |
|---|---|
| <input checked="" type="checkbox"/> Coloured covers/ Couverture de couleur | <input type="checkbox"/> Coloured pages/ Pages de couleur |
| <input type="checkbox"/> Covers damaged/ Couverture endommagée | <input type="checkbox"/> Pages damaged/ Pages endommagées |
| <input type="checkbox"/> Covers restored and/or laminated/ Couverture restaurée et/ou pelliculée | <input type="checkbox"/> Pages restored and/or laminated/ Pages restaurées et/ou pelliculées |
| <input type="checkbox"/> Cover title missing/ Le titre de couverture manque | <input type="checkbox"/> Pages discoloured, stained or foxed/ Pages décolorées, tachetées ou piquées |
| <input type="checkbox"/> Coloured maps/ Cartes géographiques en couleur | <input type="checkbox"/> Pages detached/ Pages détachées |
| <input type="checkbox"/> Coloured ink (i.e. other than blue or black)/ Encre de couleur (i.e. autre que bleue ou noire) | <input type="checkbox"/> Showthrough/ Transparence |
| <input type="checkbox"/> Coloured plates and/or illustrations/ Planches et/ou illustrations en couleur | <input type="checkbox"/> Quality of print varies/ Qualité inégale de l'impression |
| <input checked="" type="checkbox"/> Bound with other material/ Relié avec d'autres documents | <input type="checkbox"/> Includes supplementary material/ Comprend du matériel supplémentaire |
| <input type="checkbox"/> Tight binding may cause shadows or distortion along interior margin/ La reliure serrée peut causer de l'ombre ou de la distortion le long de la marge intérieure | <input type="checkbox"/> Only edition available/ Seule édition disponible |
| <input type="checkbox"/> Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/ Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées. | <input type="checkbox"/> Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image/ Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible. |
| <input type="checkbox"/> Additional comments:/ Commentaires supplémentaires: | |

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

| | | | | | | | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|-------------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 10X | 12X | 14X | 16X | 18X | 20X | 22X | 24X | 26X | 28X | 30X | 32X |
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input checked="" type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

The copy filmed here has been reproduced thanks to the generosity of:




Library of Congress
Photoduplication Service

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Library of Congress
Photoduplication Service

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".


Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

ils
u
ifier
ne
age

ata

elure,
à

32X



BULLETIN OF THE UNIVERSITY OF WISCONSIN

SCIENCE SERIES, VOL. 1, NO. 5, PP. 157-366. (CONCLUSION.)

ANALYTIC KEYS TO THE GENERA AND SPECIES OF
NORTH AMERICAN MOSSES

BY

CHARLES REID BARNES

Professor of Botany.

REVISED AND EXTENDED BY
FRED DEFOREST HEALD,


Fellow in Botany.

WITH THE COOPERATION OF THE AUTHOR.

PUBLISHED BY AUTHORITY OF LAW AND WITH THE APPROVAL OF
THE REGENTS OF THE UNIVERSITY

MADISON, WIS.
PUBLISHED BY THE UNIVERSITY
JANUARY, 1897.

PRICE \$1.00



Committee of Publication

CHARLES KENDALL ADAMS, PRESIDENT OF THE UNIVERSITY

EDITORS


WILLIAM H. HOBBS (Chairman), *Science*

NELSON O. WHITNEY, *Engineering*

FREDERICK J. TURNER, *Economics, Political Science, and History*

UNITY

History



BULLETIN OF THE UNIVERSITY OF WISCONSIN

SCIENCE SERIES, VOL. 1, NO. 5, PP. 187-388. (CONCLUSION.)

ANALYTIC KEYS TO THE GENERA AND SPECIES OF
NORTH AMERICAN MOSSES


BY

CHARLES REID BARNES

REVISED AND EXTENDED BY
FRED DUFOREST HEALD,
WITH THE COOPERATION OF THE AUTHOR.

PUBLISHED BY AUTHORITY OF LAW AND WITH THE APPROVAL OF
THE REGENTS OF THE UNIVERSITY

MADISON, WIS.
PUBLISHED BY THE UNIVERSITY
DECEMBER, 1896.



INTRODUCTION.

PREVIOUS PUBLICATIONS.

In the summer of 1886 I published for free distribution a key to the genera of mosses recognized in the Manual of Lesquereux and James. A large edition of that key was soon exhausted, and I have been frequently assured that it proved of real service to students of mosses. Although much misgiving was felt as to its accuracy very few changes have been found necessary, and it is now presented for the third time substantially as it was at the first, except for the changes involved in its extension to include the sub-genera of *Hypnum*, most of which, as given in the Manual referred to, are certainly of generic rank.

Encouraged by the approval of the key to genera, in May, 1890, I published keys to the species of North American mosses recognized by Lesquereux and James. These keys were printed in the eighth volume of the Transactions of the Wisconsin Academy of Sciences, Arts and Letters, and a considerable edition of separates was also struck off. In those keys I included the comparatively small number of new species which had been described since the issue of the Manual. The work made no pretensions to a critical study of North American mosses, but was designed only as a convenience for students. The new edition here presented is meant to serve the same purpose more fully.

THE NEW EDITION.

Three reasons have induced me to prepare a new edition.

- (1) The separate edition printed in 1890 was exhausted sev-

eral years ago, and requests for copies of it were being received continually. (2) It will be several years before the new Manual of North American mosses can be completed, and much information is needed to make critical study for that work possible. (3) A very large number of new species have been described since 1890, about which it is especially desirable to have more accurate information.

I was fortunate in being able to associate with me in this work Mr. Fred D. Heald, sometime fellow in botany in the University of Wisconsin, who has prepared the first draft of the revision. Of some particular parts of the work for which credit is due him I shall speak later.

COLLECTED DESCRIPTIONS.

Comparatively few who would be glad to collect and study mosses have access to the various publications in which the species new to our flora have been described. We have thought it best, therefore, at the risk of adverse criticism, to append to the keys collected descriptions of all species not found in Lesquereux and James' Manual. For the specialist this may be looked upon as wasted energy; but these keys are prepared chiefly for amateurs, who will doubtless be glad to have these descriptions brought to hand.

The source of the descriptions reprinted in the Appendix is always cited. The descriptions are not always the original ones, though generally they are if the species is one recently named. We have not thought it necessary in other cases even to cite the original description which can be found readily from the clue given.

KINDBERG'S NEW SPECIES.

A great number of new species have been described by Dr. N. C. Kindberg, of Linköping, Sweden, and by Dr. C. Müller, of Halle, in collaboration with Kindberg, from the collections brought together by John Macoun, botanist of

the Dominion of Canada. There is good reason to believe that a majority of these are not well founded.

It is, of course, true that species as such do not exist in nature, and, therefore, that there will be diverse judgments as to what individuals should be grouped to form a species. But there is a general agreement that changes induced by the immediate action of the environment upon the individual can not furnish a proper basis for specific distinctions. Many of Kindberg's species are established upon such slight differences in size, shape or habit as may readily be induced by the immediate action of deficient or excessive moisture, light or shade, etc.

Other species are based upon wholly insufficient material. When immature or barren specimens of *Barbula* and *Bryum* are described as new species, one who knows the remarkable variability of the vegetative parts of plants can not but doubt the value of distinctions based upon them. Only the repeated collection of barren specimens with some persistent peculiarities can justify the establishment of new species upon such material.

Moreover, critical examination of Kindberg's new species in several genera have been made by a number of bryologists, among whom may be named Mrs. Britton, Best, Grout, Barnes, True, Cheney, Renauld, and Cardot. Without exception these students have declared a considerable number of the plants described as new to be referable to those already described, of which they are either slight varieties or forms unworthy a separate name. In a revision of the genus *Dicranum* (ined.), for example, Barnes and True have been compelled to reject seventeen out of eighteen species described by Kindberg.

These species are in many cases not only ill founded. Few of the descriptions are either accurate or sufficient. Very many are so brief and so purely comparative that it is impossible to obtain a definite idea of the diagnostic characters.

SCOPE OF THE KEYS.

In the keys we have endeavored to include all the species reported or described as belonging to our flora, unless a later special study of the genus has shown the addition to be untenable. Such special studies are cited in the keys. The incompleteness of many descriptions, and the lack of diagnostic characters have seriously hampered us in rearranging the keys, and they will fail, therefore, in many cases to differentiate described species. We have been at considerable pains to include as many of the barren and insufficiently described species as possible, in order that these may be recognized, if they exist, or may be referred to their appropriate place. Material which will enable students either to diagnose them clearly or to overthrow them would be of great value.

The keys do not generally discriminate varieties, but it is very important that collectors and amateurs keep in mind the variability of the mosses and seek to bring to light as many facts as possible bearing on this problem, so that the range of variability may be ascertained. To facilitate this we have also collected descriptions of varieties not appearing in the Manual. In these varietal descriptions the comparisons, unless otherwise stated, are with the characters of the species.

NOMENCLATURE.

It seemed to us highly desirable to avoid the making of new names in a compilation like this. We have therefore used, as a basis for our work, Renauld and Cardot's *Musci Americae Septentrionalis*.¹ In basing our keys upon that list we do not thereby express our adherence to the nomenclature employed. Of many species therein neither the

¹Published in the *Revue Bryologique* 19: 65-69. 1892 and 20: 1-32. 1893. Afterward this was issued as a separate and, most unfortunately, re-paged and revised.

le all the species
r flora, unless a
the addition to
ted in the keys.
and the lack of
pered us in re-
efore, in many
e have been at
the barren and
e, in order that
ay be referred
ch will enable
or to overthrow

varieties, but it
ateurs keep in
eek to bring to
his problem, so
ined. To facil-
ons of varieties
arietal descrip-
tated, are with

t the making of
have therefore
t Cardot's *Musci*
s upon that list
to the nomencla-
in neither the

1892 and 20: 1-32.
most unfortunately,

present names nor the autonomy can be maintained. But it enables us to avoid constructing new names in this publication, so that it need not be cited in the future literature of taxonomy.

This list also saved us much labor of compilation, both of names and of geographical distribution. Only such changes were made in the species included as seemed warranted by the special studies before mentioned. To the species therein enumerated we have added those described since its publication up to January 1, 1896.

We have departed from this list and published monographs in only two genera. In *Dicranum* we have used the results of the study of Barnes and True, and in *Amblystegium* those of Cheney (both at present unpublished), so far as the omission of certain species of these genera is concerned. We have not included some new species yet in MSS., believing that this bulletin is not a suitable place for publishing them.

It is scarcely necessary to add that no attention has been paid to *nomina nuda*.

CITATIONS.

For the sake of brevity we have cited *Macoun's Catalogue of Canadian Plants, Part VI. Mosses*, by Macoun and Kindberg, simply as *Mac. Cat.*, to which its independent paging lends itself. Likewise *Rabenhorst's Kryptogamen-Flora von Deutschland, Oesterreich und der Schweiz, Part IV. Die Laubmoose*, by Limpricht, is cited as *Limpr. Laubm.* Other citations are given in full or with abbreviated accurate titles. In all cases we have cited the original publication of Renauld and Cardot's *Musci Americæ Septentrionalis* in the *Revue Bryologique*.

ACKNOWLEDGMENTS.

Upon Mr. Heald has fallen the bulk of the work of collating the list of species to be included and selecting the descriptions to be reprinted. Many of the latter are trans-

lated from Latin, French, or German, with most of which he is also to be credited. Perhaps his most difficult task was in preparing the new part of the key to the genera of Hypnaceæ, and the keys to the species of *Hypnum*. The incorporation of the many new species in the keys to the larger genera, such as *Sphagnum*, *Barbula*, *Grimmia* and *Bryum*, was also a difficult task, made doubly difficult by imperfect descriptions and want of figures. There is scarcely a key which he has not altered more or less to permit these insertions, and some are completely rearranged.

In this work advantage has been freely taken of the suggestions, and sometimes of the very characters, in the keys found in Limpricht's *Laubmoose*, Braithwaite's *British Moss Flora*, Husnot's *Muscologia Gallica*, and Warnstorf's *Charakteristik und Uebersicht der nord-, mittel-, und südamerikanischen Torfmoose*.

For the key of *Dicranum* thanks are due to Professor R. H. True and for that of *Amblystegium* to Professor L. S. Cheney, both of the University of Wisconsin.

The first draft of the manuscript prepared by Mr. Heald I have carefully edited, revised, and, in some parts, rearranged. In the absence of Mr. Heald from the country, the not inconsiderable labor of attending to the details of publication has fallen upon me.

For corrections to the previous edition acknowledgments are due to Mrs. E. G. Britton, Miss Clara E. Cummings, Mr. E. A. Rau, Dr. C. W. Swan, Mr. Marshall A. Howe, Mr. Edo Claassen, Professor L. S. Cheney and Professor R. H. True.

The unfailing kindness of Mrs. E. G. Britton in supplying specimens, information as to particular species, and copies of several inaccessible descriptions receives also our grateful recognition.

Professor Conway MacMillan has also put us under obligations by the loan of periodicals not accessible here.

We shall be grateful for notice of errors or omissions in the present edition.

USE OF THE KEYS.

Upon page 157 is a key to the four orders of Musci. The first three orders each have a single genus only. To the genera of the fourth order, BRYALES, the key upon the following twelve pages is devoted.

In this key there are always at least two choices presented, whose equivalence is indicated by similar numbers or signs preceding the line, or by equal indention of the lines, which usually begin with the same word or its opposite. When the second member does not follow on the same page a reference to the page upon which it does occur follows the first. For an example see page 158. When there are more than two choices offered, all except the first and last are followed by a number or sign to indicate that a succeeding one is to be sought. For an example see page 160.

When the accepted choice is followed by a dotted line leading to the name of a genus, the description of that genus will be found in Lesquereux and James' Manual of the Mosses of North America on the page indicated by Arabic figures of ordinary face (999). The key to the species of that genus will be found on the page of this work indicated by the bold face figures (999) following. If the genus be not described in the Manual it will be found described in the Appendix, on the page indicated by italic figures (999.)

In the keys to species the correlative choices are indicated in the same way. Arabic figures (999) following the name of a species show that the species will be found described in the Manual on the page named. If followed by italic figures (999), a description will be found in the Appendix on the page indicated.

APOLOGIA PRO LABORE SUO.

No one realizes more clearly than the writer that such work as that which is now presented to students is in one

sense "dead work." Yet in another it is vital. It is in itself of only temporary value. It perishes with the using. The user may soon get beyond the need of a crutch.

The only justification for such work is its usefulness to amateurs, who may by its help put bryologists in possession of facts regarding the distribution, variation, and relationships of our moss flora which could not be gained without their aid. These keys are published, therefore, purely as an encouragement to the study of our mosses, which with the Hepaticæ are more neglected than any other group of which we have accessible descriptions.

An earnest student equipped with patience, some skill in dissection, a compound microscope, and the Manual, ought to be able with the assistance of these keys to determine the names of most of the mosses which he can collect. Those which remain uncertain he can refer to those who possess the illustrations and exsiccati which are often indispensable for identification.

CHARLES R. BARNES.

University of Wisconsin,
December 15, 1896.

NSIN.

ital. It is in
with the using.
crutch.

usefulness to
sts in posses-
tion, and re-
not be gained
d, therefore,
our mosses,
ed than any
riptions.

some skill in
annual, ought
to determine
can collect.
to those who
re often in-

BARNES.

ANALYTIC KEY

TO THE

GENERA OF MOSSES.

ORDER I.—Sphagnales.

Capsule dehiscing by a deciduous operculum, peristome
none, leaves composed of large hyaline cells, with
intervening rows of small chlorophyllose ones.

Genus single *Sphagnum*, 12. 170.

ORDER II.—Andreæales.

Capsule dehiscing by four (rarely six) longitudinal slits.

Genus single *Andreæa*, 25. 175.

ORDER III.—Archidiales.

Capsule bursting irregularly, spores few and very large.

Genus single *Archidium*, 49. 176.

ORDER IV.—Bryales.

Capsule bursting irregularly (spores numerous) or gen-
erally dehiscing by a deciduous operculum, in the
latter case usually furnished with a peristome.
Leaves not sphagnoid.

Genera numerous as follows:

I. CLEISTOCARPI.—*Capsule without a deciduous operculum.*
(II on p. 158.)

A. *Green protonema persistent.* (B on p. 158.)

Leaves ecostate.

Capsule colorless *Micromitrium*, 37. 176.

Capsule colored *Ephemerum*, 37. 176.

Leaves costate *Ephemerum*, 37. 176.

B. *Green protonema not persistent.*

Margins of leaves flat or incurved.

Leaves lance-obovate to broad ovate or ovate lanceolate,
not abruptly pointed.Margins dentate or serrate **Physcomitrella**, 39.Margins entire **Voitlia**, 42, 43.

Leaves linear-lanceolate to subulate or abruptly pointed.

Calyptra mitrate **Bruchia**, 45. 177.Calyptra cucullate **Pleuridium**, 43. 177.[*Astomum* may be sought here.]

Margins of leaves more or less revolute.

Capsule spherical **Sphaerangium**, 40. 176.

Capsule short-pointed.

Calyptra mitrate **Microbryum**, 45, 37.Calyptra cucullate **Phaseum**, 41. 177.II. STEGOCARPI.—*Capsule with a deciduous operculum.*A. ACROCARPI.—*Capsule at the apex of the stem, either the main
shoot or a well developed branch.* (B on p. 166.)1. *Mouth of the capsule naked.* (2 on p. 159.)[*Weisia viridula* vars. may be sought here.]* *Leaf-cells isodiametric, at least above the middle of the leaf, often
obscure.* (** on p. 159.)Lid imperfectly formed, persistent **Astomum**, 51. 178.

Lid perfect, deciduous.

Capsule immersed.

Leaves lamellose **Pharomitrium**, 100.

Leaves not lamellose.

Apex hyaline or prolonged into a hyaline hair.

Perichaetial leaves ciliate **Hedwigia**, 152.Perichaetial leaves not ciliate **Grinnia**, 134. 197.Apex obtuse, plants robust, blackish **Scouleria**, 137. 197.

Capsule exserted, ribbed when dry.

Calyptra cucullate.

Leaves with long hyaline points **Braunia**, 152.

Leaves not hyaline pointed.

Capsule contracted below the orifice **Amphoridium**, 153. 201Capsule not contracted below the orifice **Zygodon**, 206. 201.

Calyptra campanulate-mitrate, plicate, usually

hairy **Macromitrium**, 178. 206.

- Calyptra long clavate-companulate, not plicate nor hairy.
 Costa with loose cells in center of section . . . **Encalypta**, 180. 206.
 Costa with central stereid strand . . . **Merceya**,¹ 242.
- Capsule exerted, not ribbed when dry.
 Calyptra persistent, plicate, twisted . . . **Calymperes**, 184. 207.
 Calyptra persistent, smooth, surpassing capsule.
 Costa with loose cells in center of section . . . **Encalypta**, 180. 206.
 Costa with central stereid strand . . . **Merceya**,¹ 242.
- Calyptra not persistent, often fugacious.
 Leaves ciliate at the base, obtuse, flat . . . **Edipodium**, 244, 245.
 Leaves not ciliate at the base, pointed, carinate,
 distichous **Eustichia**, 94.
 Leaves not ciliate at the base, pluriseriate.
 Linear-lanceolate, margins plane.
 Costa filling elongated point, plants minute, . . . **Anodus**, 96.
 Costa percurrent.
 Capsule on a short lateral branch . . . **Anoctangium**, 54. 179.
 Capsule terminating main axis . . . **Gymnostomum**, 52. 178.
 Ovate lanceolate, margins reflexed . . . **Barbula**, 115. 191.
 Broad, often hair pointed, plants minute . . . **Pottia**, 100. 188.

**** Leaf cells plainly elongated, distinct.**

- Lid small, convex or short-conic, capsule microstome.
 Leaves vertically inserted **Schistostega**, 188.
 Leaves subulate, dentate **Bartramia**, 203. 210.
 Leaves broad, entire, calyptra enclosing capsule **Pyramidula**, 196.
 Lid large (rarely small), capsule macrostome.
 Capsule splitting at the middle . . . **Aphanorhagma**, 196.
 Capsule dehiscing regularly above the middle, not
 covered by calyptra **Phycomitrium**, 196. 209.

2. Mouth of the capsule furnished with a peristome.

* *Peristome single.* (** on p. 164.)

+ *Teeth articulate.* (+ + on p. 163.)

++ *Teeth eight.* (++ ++ on p. 160.)

- Leaves thick, coriaceous **Octoblepharum**, 91.

[*Orthotrichum* and *Ptychomitrium* (§ *Notarisia*) may be sought here.]

¹ Genus only known in sterile state; wherefore its place in the key cannot be determined except as its leaves ally it to *Encalypta*.

++ ++ *Teeth sixteen, calyptra mitrate.* (++ ++ ++)

= *Calyptra plicate.*

- Teeth cribose, purple *Coscinodon*, 154. 201.
 Teeth filiform, trifid *Ptychomitrium*, 156. 201.
 Teeth approximate or connate in pairs.
 Lanceolate to subulate, papillose *Ptychomitrium*, 156. 201.
 Triangular-lanceolate, articles quadrate.
 Basal leaf-cells linear, chlorophyllose *Ulota*, 160. 202.
 Basal leaf-cells hexagono-rectangular, hyaline *Orthotrichum*, 164. 203.
 Teeth short, pale, fragile *Macromitrium*, 178. 206.

= = *Calyptra not plicate.*

Aquatic, floating.

- Leaves distichous *Fissidens* (§ *Octodiceras*), 89. 186.
 Leaves pluriseriate *Cinclidotus*, 134.

Terrestrial.

Very small, gregarious.

- Teeth broad, erose-truncate, hyaline *Brachyodus*, 98.
 Teeth linear-lanceolate, deeply bifid *Campylostellium*, 99.

Larger, above 1 cm. in height.

- Leaf-cells small, quadrate or punctate, obscure.
 Beak long-clavate *Encalypta*, 180. 206.
 Beak long or short, not clavate.

- Teeth lanceolate, flat, subentire or cribose or
 2-3-fid to the middle *Grimmia*, 134. 197.

- Teeth linear-lanceolate, 2-3-fid to below middle, or
 cleft to base into filiform segments *Rhacomitrium*, 147. 199.

Leaf-cells large, very distinct, pedicel with a
 prominent apophysis.

- Apophysis smaller than the capsule.
 Leaves entire, obtuse *Dissodon*, 189. 207.
 Leaves serrate, acute or acuminate *Tayloria*, 190. 208.
 Apophysis exceeding the capsule *Splachnum*, 193. 208.

++ ++ ++ *Teeth sixteen, calyptra cucullate.* (++ ++ ++ ++ on p. 163.)

= *Leaves distichous.* (= = on p. 161.)

- Leaves subulate *Distichium*, 93. 187.
 Leaves broader, with a prominent vertical wing *Fissidens*, 81. 185.

== *Leaves pluriseriate.*¶ *Capsule unsymmetric, cernuous-inclined or arcuate.*Teeth filiform-bifid from a membranous base . . . **Desmatodon, 110. 190.**

Teeth irregularly lacerate or bifid to the middle or below.

Leaf cells not enlarged at the basal angles, roundish or quadrate above.

Lid long-beaked, leaves serrulate, peristome equaling

half the capsule **Dichodontium, 61. 180.**

Lid long-beaked, leaves crenulate or denticulate,

peristome shorter **Cynodontium, 59. 179.**Lid short-beaked **Oreoweisia, 58.**

Leaf-cells not enlarged at the basal angles, oblong above,

rectangular at base **Dicranella, 64. 180.**

Leaf-cells enlarged-quadrate at the basal angles.

Linear at base, capsule not strumose, dioicous . . . **Dicranum,¹ 67. 181.**Rectangular at base, capsule strumose, monoicous **Cynodontium, 59. 179.**Leaf-cells of two kinds, in two or three layers . . . **Leucobryum, 90. 187.**

Teeth bifid to near the base.

Lid conic, leaves subulate **Trichodon, 92. 187.**Lid conic, leaves lanceolate **Ceratodon, 92. 187.**Lid aristate, neck very long **Trematodon, 62. 180.**Teeth not cleft, short, irregular **Catoscopium, 211.**Teeth not cleft, cohering by their tips . . . **Conostomum, 207.**

Teeth not cleft, perforate.

Neck long, exceeding the capsule **Trematodon, 62. 180.**Neck inconspicuous, plants small **Discellum, 188.**Neck inconspicuous, plants large **Oreoweisia, 58.**

Teeth not cleft nor perforate.

Lid with a short thick oblique beak **Oreoweisia, 58.**Lid with a short slender oblique beak . . . **Cynodontium, 59. 179.**[*Mittelehferia* and *Funaria* may be sought here.]¶ ¶ *Capsule symmetric, pendulous on a flexuous pedicel.*

(¶ ¶ ¶ on p. 162.)

Teeth bifid to the middle **Campylopus, 77. 184.**Teeth bifid to the base, free **Dicranodontium, 77. 184.**

Teeth bifid to the common membranous base.

Connivent and slightly twisted **Desmatodon, 110. 190.**Erect, not twisted **Trichostomum, 108. 190.**Teeth entire, short, plants minute **Seligeria, 96. 188.**¹ The genus *Monocranum* C. Mull. is not included in the key on account of its doubtful nature.

¶ ¶ ¶ *Capsule symmetric, erect.*

Teeth bifid to the common membranous base.

Leaves subulate to lance-subulate from a broader

base *Ditrichum*, 105. 189.

Leaves broader.

Lid short, conic or beaked *Desmatodon*, 110. 190.Lid elongated, conic *Trichostomum*, 108. 190.[*Barbula* may be sought here.]

Teeth deeply bifid or cleft to the base, free.

Leaf-cells small, not enlarged at the angles, oblong

above *Dicranella*, 64. 180.Leaf-cells small, not enlarged at the angles, roundish
or quadrate above.Lid oblique *Cynodontium*, 59. 179.Lid straight *Leptodontium*, 116, 117.Leaf-cells small, enlarged-quadrate at the angles *Dicranum*, 67. 181.Leaf-cells large, distinct *Aongstromia*, 63.

Teeth cribose, perforate or slightly cleft.

Leaf-cells enlarged-quadrate at the angles.

Capsule broad-pyriform *Blindia*, 98. 108.Capsule oval to sub-cylindric *Dicranoweisia*, 57. 179.

Leaf-cells not enlarged at the angles.

Teeth large, mostly cribose.

Pedicel little exceeding the often hair-pointed

leaves *Grimmia*, 134. 197.Pedicel long, leaves hair-pointed *Desmatodon*, 110. 190.

Pedicel long, leaves not hair-pointed.

Leaves serrate just above sheathing base *Eucladium*, 46, 47.Leaves entire or crenulate above *Didymodon*, 104. 189.

Teeth small, often truncate or rudimentary.

Leaf-margins involute above *Weisia*, 55. 179.

Leaf-margins revolute or plane.

Leaves densely papillose in the upper part *Didymodon*, 104. 189.

Leaves not papillose.

Capsule long exserted, lid completely deciduous *Pottia*, 100. 188.

Capsule subimmersed, lid adhering to colu-

mella *Sconleria*, 137. 197.

Teeth entire.

[*Cynodontium* and *Eucladium* may also be sought here.]Capsule with a long, thick apophysis *Tetraplodon*, 191. 208.

Capsule oval to subcylindric.

Not ribbed when dry.

Teeth short, leaves entire, narrow *Weisia*, 55. 179.

- Teeth short, leaves serrate, broad . . . **Syrrophodon**, 185. 207.
 Teeth linear-filiform, connate at base . . . **Didymodon**, 104. 180.
 Teeth narrowly lanceolate, free . . . **Dicranoweisia**, 57. 179.
 Ribbed when dry **Rhabdoweisia**, 58. 179.
 Capsule short-pyriform, turbinate when dry.
 Teeth blunt **Seligeria**, 96. 188.
 Teeth acute **Blindia**, 98. 108.
 Capsule pyriform, not turbinate when dry.
 Plants gregarious or subcespitose . . . **Entosthodon**, 199. 209.
 Plants in deep compact tufts . . . **Mielichhoferia**, 214. 211.
 Capsule ovate-globose, lid obliquely long-beaked **Drummondia**, 160. 209.
 Capsule globose, lid beakless, small . . . **Bartramia**, 203. 210.

+++ Teeth thirty-two.

- Teeth cancellate **Barbula**, 115. 191.
 Teeth filiform or linear, almost terete, arising from a long
 or short basilar membrane.
 Short, slightly, if at all, twisted.
 Leaves (upper) convolute-sheathing at base **Leptobarbula**, 123, 124
 Leaves not sheathing at base.
 Subulate or lance-subulate from a broader
 base **Ditrichum**, 105. 189.
 Broader, lid elongated-conic . . . **Trichostomum**, 108. 190.
 [*Barbula rigidula* will be sought here.]
 Broader, lid short-conic or short-beaked **Desmatodon**, 110. 190.
 Long, twisted to the left **Barbula**, 115. 191.
 Teeth flat, not from a distinct basilar membrane.
 Cells of capsule linear-oblong . . . **Dicranodontium**, 77. 184.
 Cells of capsule irregularly polygonal . . . **Didymodon**, 104. 189.

+- Teeth not articulate.

++ Teeth four, solid.

- Capsule linear-oblong, stems long, conspicuous . . . **Tetraphis**, 186. 207.
 Capsule ovate, stems very short . . . **Tetrodontium**, 187.

+++ Teeth thirty-two or sixty-four.

[*Leptobarbula* may be sought here.]

- Calyptra cucullate, capsule symmetric or nearly so.
 Leaves undulate-cripsed when dry, lamellæ few (2-8),
 straight **Atrichum**, 255. 221.

- Leaves sub-tubulose at apex, lamellae undulate or numerous **Oligotrichum**, 268. 222.
 Calyptra cucullate, capsule unsymmetric, arcuate incurved **Psilopilum**, 269.
 Calyptra mitrate, densely hairy.
 Capsule not angular, teeth 32 **Pogonatum**, 260. 222.
 Capsule 4-6 angled, teeth 64 **Polytrichum**, 263. 223.

** *Peristome double.*

+ *Capsule symmetric, erect (sometimes inclined in age).*

- Teeth almost 0, imperfect or rudimentary . . . **Macromitrium**, 178. 206.
 Teeth perfect, linear or filiform.
 Capsule smooth.
 Teeth revolute **Schlotheimia**, 179.
 Teeth not revolute.
 Leaves ecostate **Fontinalis**, 268. 224.
 Leaves costate.
 Leaf-cells oval hexagonal, capsule long exserted **Leptotheca**, 251.
 Leaf-cells hexagonal-rhombic, capsule immersed **Brachelyma**.¹
 Leaf-cells linear-attenuate, capsule immersed or emergent **Dichelyma**, 272. 225.
 Capsule ribbed and twisted.
 Costa with loose cells in center of section . . . **Encalypta**, 180. 206.
 Costa with central stereide band **Merceya**,² 242.
 Teeth broadly or narrowly triangular-lanceolate, pale, capsule ribbed, not twisted.
 Leaf-cells at base linear, chlorophyllose **Ulota**, 160. 202.
 Leaf-cells at base hexagono-rectangular, hyaline.
 Lid straight, conical or acuminate **Orthotrichum**, 164. 203.
 Lid obliquely rostrate **Zygodon**, 206. 201.

+ + *Capsule unsymmetric, inclined or oblique or pendulous.*

+ + *Inner peristome a plaited cone. (+ + on p. 165.)*

- Pedical thick, red, densely verrucose **Buxbaumia**, 267. 223.
 Pedicel very short, almost none **Diphyseium**, 266.

¹ Including one species, *B. subulatum* Sch. (*Dichelyma subulatum* Myrin; *Cryphaea mundata* Nees. L. and J. Mannal, pp. 274 and 413.)

² See footnote, p. 159.

++ ++ *Inner peristome a membrane, carinate or cut into sixteen segments; these sometimes separated by cilia.*

= *Cilia very short, rudimentary or none.*

Membrane entire, 16-carinate *Cinclidium*, 249. 221.

Membrane cleft into segments.

Segments entire or interruptedly cleft along the middle line.

Shorter than the teeth or rudimentary *Funaria*, 200. 210.

Equaling the teeth in length.

Leaves squarrose-recurved from the middle . . . *Paludella*, 213.

Leaves not squarrose.

Pedicele long.

Leaf-cells narrowly rhombic-hexagonal, tending

to linear, leaves narrow *Webera*, 215. 211.

Leaf-cells and leaves broader *Bryum*, 223. 214.

Pedicele short, neck long *Zieria*, 240. 219.

Far exceeding the teeth in length.

Pedicele long, leaf-cells large, pellucid *Amblyodon*, 211.

Pedicele long, leaf-cells small, rectangular, chloro-

phyllous *Meesia*, 212. 211.

Pedicele short, neck long *Zieria*, 240. 219.

Segments bifid, divisions divaricate.

Leaves lanceolate to subulate, large *Bartramia*, 203. 210.

Leaves lanceolate or broader, smaller *Philonotis*, 208. 211.

Segments filiform, united by fours at their tips . . . *Timmia*, 254. 221.

= = *Cilia present.*

Appendiculate.

Leaves lance-subulate, cells linear *Leptobryum*, 215.

Leaves broader, cells rhombic-hexagonal *Bryum*, 223. 214.

Inappendiculate.

Capsule not ribbed when dry.

Leaves lanceolate, glossy, cells narrowly rhombic-

hexagonal, inclining to linear *Webera*, 215. 211.

Leaves ample, soft, oblong, ovate to obovate or

broader, cells round-hexagonal *Mnium*, 241. 219.

Leaves narrowly lanceolate, rigid *Rhizogonium*, 250.

Capsule ribbed when dry.

Oblong or elongated pyriform *Aulacomnium*, 252. 221.

Sub-globose *Philonotis*, 208. 211.

B. PLEUROCARPI. *Capsule on a very short lateral branch.*
 [Fontinalis, Dichelyma and Anactangium may be sought here.]

1. *Peristome single (rarely none), teeth eight or sixteen.*

[Species belonging to genera under "2" *infra* may be sought here.]

Leaves distichous, with broad vertical wing . . . Fissidens, 81. 185.

Leaves pluriseriate.

Entire (except perhaps at the apex) and ecostate.

Perichaetial leaves erose-dentate at apex . . . Habrodon, 296.

Perichaetial leaves entire . . . Leucodon, 287. 227.

Entire and costate.

Short acuminate, teeth 16, yellowish . . . Clasmatodon, 297.

Obtuse, teeth 8, red . . . Cryphaea, 275. 226.

Serrate, capsule emergent . . . Leptodon, 278. 226.

Serrate to ciliate-dentate, capsule long pedicelled . . . Fabronia, 294. 228.

2. *Peristome double, the inner often imperfect.*

* Segments none or short, or obscured by adhering to teeth.
 (* * on p. 167.)

+ *Leaves papillose.*

Entire, ovate to ovate-lanceolate.

Teeth ciliate-papillose . . . Leskea, 301. 229.

Teeth not papillose . . . Anomodon, 304. 230.

Entire or cristate-serrate, obovate or spatulate Pterigynandrum, 288. 228.

Spinulose-dentate to fimbriate (rarely entire) deltoid or

round-ovate . . . Thelia, 298. 229.

Serrate, broadly ovate . . . Pterogonium, 289. 228.

+ + *Leaves not papillose.*

Capsule straight.

Segments bifid or adherent to the teeth.

Plants small (1-2 cm.), capsules about 2 mm. . . Pylaissea, 308. 230.

Plants large (4-6 cm.), capsules about 4 mm. Cylindrothecium, 310. 231.

Segments not bifid nor adherent.

Leaves ecostate or obscurely bicostate . . . Neckera, 281. 226.

Leaves costate . . . Antitrichia, 290. 228.

Capsule curved or arcuate . . . Homalothecium, 309. 230.

at lateral branch.
may be sought here.]

eight or sixteen.

may be sought here.]

. Flissidens, 81. 185.

Habrodon, 296.

Leucodon, 287. 227.

Clasmatodon, 297.

. Cryphaea, 275. 226.

Leptodon, 278. 226.

Fabronia, 294. 228.

imperfect.

adhering to teeth.

. Leskea, 301. 229.

Anomodon, 304. 230.

gynandrum, 288. 228.

oid or

. Thelia, 298. 229.

terogonium, 289. 228.

Pylaisaea, 308. 230.

androthecium, 310. 231.

. Neckera, 281. 226.

Antitrichia, 290. 228.

alothecium, 309. 230.

** Segments not distinctly keeled, narrow. (***)

+ Leaves costate.

Cells isodiametric to oval-rhombic.

Papillose.

Stem and branch-leaves similar Leskea, 301. 229.

Stem-leaves much smaller than branch-leaves Anomodon, 304. 230.

Not papillose.

Annulus large, compound, perichaetial leaves
costate Cryphaea, 275. 226.

Annulus simple, perichaetial leaves ecostate Lescuraea, 414, 416.

Annulus none.

Endostome with cilia Alsia, 279. 226.

Endostome without cilia.

Teeth erect or incurved when dry.

Leaf-cells minute, obscure, alar cells elongated Neckera, 281. 226.

Leaf-cells distinct, alar cells quadrate Myrinea, 410, 411.

Teeth reflexed when dry Anacamptodon, 296.

Cells linear or vermicular.

Annulus none Neckera, 281. 226.

Annulus present Antitrichia, 290. 228.

[Cylindrothecium, with leaves obscurely costate, may be sought here.]

+ + Leaves ecostate.¹

Annulus none Neckera, 281. 226.

Annulus present.

Leaf-cells quadrate at basal angles.

Plants small, capsules about 2 mm. long Platygryium,² 307.

Plants large, capsule about 4 mm. long Cylindrothecium, 310. 231.

Leaf-cells not quadrate at the basal angles Orthothecium, 315. 232.

*** Segments distinctly keeled, often broad.

+ Capsule symmetric, erect. (+ + on p. 168.)

[Species of *Hypnum* with erect or sub-erect capsules will be sought here.]

+ + Leaves papillose. (+ + + on p. 168.)

Plants large; branches erect, dendroid Alsia, 279. 226.

Plants long, pendent from trees, branches filiform Meteorium, 286. 227.

Plants small, branches erect, julaceous Myurella, 300. 229.

¹*Taxithellum planum* (= *Hypnum planum* Brid. L. & J. Man. 411) falls here, but on account of lack of data in regard to the annulus it can not be exactly located.

See also Appendix, no. 416.

++ ++ *Leaves not papillose.*

- Leaves costate or ecostate, complanate, pseudo-distichous.
 Middle leaf-cells linear, cilia none or very short . . . *Homalia*, 285. 227.
 Middle leaf-cells rhomboidal, cilia solitary, $\frac{2}{3}$ length
 of the segments *Stereophyllum*,¹ 411.
 Leaves costate, plants dendroid *Climacium*, 313. 231.
 Leaves ecostate, annulus large (none in *Cyl. Drummondii*).
 Cells quadrate at basal angles.
 Teeth hyaline margined *Platygyrium*, 307.
 Teeth not hyaline margined *Cylindrothecium*, 310. 231.
 Cells not quadrate at basal angles *Orthothecium*, 315. 232.
 Leaves ecostate, annulus small, narrow *Pylaissea*, 308. 230.

+ + *Capsule unsymmetric, often arcuate.*++ *Leaf-cells large, calyptra mitrate.*

- Leaves mucronate or acute or acuminate *Hookeria*, 292. 228.
 Leaves obtuse *Pterigophyllum*, 293.

++ ++ *Leaf-cells small, calyptra cucullate.*[*Climacium Ruthenicum* will be sought here.]¶ *Leaf-cells short (1 : 5 or less).* (¶ ¶ on page 169.)[*Isothecium Brewertianum* may be sought here.]

Leaves papillose.

Paraphyllia present.

- Costa short or none or double *Heterocladium*, 320. 232.
 [*Pseudoleskea malacoclada* will be sought here.]

Costa strong.

- Capsule oval or oblong, lid convex-conic *Pseudoleskea*, 319. 232.

Capsule cylindrical or if oval oblong then lid long

- rostrate *Thuidium*, 321. 233.

- Paraphyllia none *Claopodium*, 327. 234.

Leaves not papillose.

Shortly bicostate.

- Stem leaves filiform pointed *Heterocladium*, 320. 232.

- Stem leaves not filiform pointed *Tripterocladium*, 330. 234.

Unicostate or ecostate.

[*Tripterocladium rupestre* will be sought here.]

- Plants dendroid, leaves coarsely serrate *Thamnum*, 361. 240.

- Plants creeping, leaves entire or denticulate
 above *Amblystegium*, 371. 242.

¹ *Stereophyllum Wrightii* R. and C. Rev. Bryol. 20: 23. 1893. (*Hypnum Wrightii* Sull. L. and J. Man. 411.)

¶ ¶ *Leaf-cells long (1:5 or more).*¹

Leaves uncostate half way or more.

Seta rough.²

Leaves deeply plicate lengthwise . . . *Camptothecium*, 331. 234.

Leaves not deeply plicate.

Lid convex conic to long conic (rostellate in *Scler. caespitosum*).

Leaf-cells not abruptly enlarged at base, upper
usually distinct, elongated rhombic *Brachythecium*, 334. 235.

Leaf-cells abruptly enlarged at the base, indistinct,
linear-vermicular³ . . . *Scleropodium*, 346. 237.

Lid more or less long rostrate . . . *Eurhynchium*, 351. 238.

Seta smooth.⁴

Lid more or less long rostrate.

Leaves ovate or ovate lanceolate . . . *Rhynchostegium*, 358. 240.

Leaves deltoid . . . *Eurhynchium*, 351. 238.

Lid convex to conic or apiculate.

Branches fasciculate, stoloniferous, leaves generally

papillose, not plicate . . . *Isothecium*, 347. 236.

Branches irregularly pinnate, leaves not pap-
illose . . . *Brachythecium*, 334. 235.

Branches regularly or irregularly pinnate, leaves not
papillose, when plicate also falcate-secund . . . *Hypnum*,⁵ 244.

Leaves ecostate, or costa very short or double.

Leaves secund.

Paraphyllia none . . . *Hylocomium*,⁶ 409. 250.

Paraphyllia few or abundant . . . *Hypnum*, 244.

Leaves complanate spreading.

Lid rostrate.

Leaves serrulate to the base . . . *Rhynchostegium*, 358. 240.

Leaves entire or serrulate at the apex . . . *Plagiothecium*, 362. 241.

Lid convex or conic . . . *Plagiothecium*, 362. 241.

Leaves equally spreading (also *Hylocomium* 185-188).

Lid long subulate-rostrate . . . *Raphidostegium*, 355. 239.

Lid conic or short rostrate.

Capsule symmetric, erect or cernuous . . . *Plagiothecium*, 362. 241.

Capsule unsymmetric, arcuate . . . *Hypnum*,⁵ 244.

¹ *Amblystegium* species, especially *riparium* and *vacillans*, may be sought here.

² *Isothecium lentum*, *Rhynchostegium curvisetum*, and *Hypnum chloropterum* will be sought here.

³ *Brachythecium populicum* will be sought here.

⁴ *Camptothecium nitens* will be sought here.

⁵ Including the subgenera XVII to XXVI of L. & J. Man. 407.

⁶ Including *Pleurozium* of L. & J. Man. 407.

ANALYTIC KEY
TO THE
SPECIES OF MOSSES.

SPHAGNUM, p. 12.

I. *Cortical cells of stem and branch without spiral fibrils; branch leaves mostly blunt and toothed, rarely acute.*

Branches 7—13 in a fascicle § IV. *Polyclada*, p. 173.

Branches 3—6 in a fascicle, or fewer.

Hyaline cells of stem leaves fibrillose.

Cortical cells thin walled.

Border of stem leaves much broader at base, chloro-

phylllose cells exposed on the inner face § I. *Acutifolia*, p. 171.

Border of stem leaves not or slightly broader at the

base § VI. *Subsecunda*, p. 173.

Cortical cells thick walled § III. *Cuspidata*, p. 172.

Hyaline cells of stem leaves without fibrils.

Chlorophyllose cells of branch leaves exposed on

inner face § I. *Acutifolia*, p. 171.

[*S. subsecundum* and *S. contortum* may be sought here.]

Chlorophyllose cells exposed on both faces, only on outer face, or included.

Cell membranes of stem leaves resorbed.

Hyaline cells of branch leaves with few pores on

the outer surface (4-6) § II. *Squarrosa*, p. 172.

Hyaline cells of branch leaves with numerous

pores on outer surface (12-16+) § V. *Rigida*, p. 173.

Cell membranes of stem leaves not resorbed, cortex

thick walled § III. *Cuspidata*, p. 172.

- II. *Cortical cells of stem and pendent branches always with fibrils and pores; branch leaves not blunt and toothed but cucullate and hyaline bordered* §VII. *Cymbifolia*, p. 175.

I. *Acutifolia*.

- A. *Branch leaves toothed only at blunt apex, without resorption furrows on margined border.*

1. *Stem leaves lacerate-fringed at apex.*

- a. *Stem leaves with completely resorbed cell membranes in upper part, without fibrils.*

Stem leaves widening upward, spatulate, apex and upper margins lacerate-fringed *S. fimbriatum*, 14.

Stem leaves not widening upward, linguiform and lacerate fringed only at broad round apex *S. Girgensohnii*,¹ 13.

- b. *Stem leaves with resorbed membranes only at apex, upper $\frac{1}{2}$ to $\frac{2}{3}$ with fibrils* *S. Bolanderi*, 1.

2. *Stem leaves usually dentate at apex, nowhere with completely resorbed cell membranes.*

- a. *Stem leaves slightly or not at all narrowed upward, linguiform.*

Outer wall of cortical cells always porose *S. Russowii*, 2.

Outer wall of cortical cells without pores.

Branch leaves curved when dry, erect-spreading, stem leaves without fibrils.

Pores of branch leaves very small, near apex, *S. Warnstorffii*, 3.

Pores of branch leaves of medium size or large *S. Vancouveriense*, 4.

Branch leaves frequently secund, stem leaves frequently with fibrils, wood-cylinder never brown *S. tenellum*, 5.

Branch leaves densely imbricate, stem leaves without fibrils, wood-cylinder always brown *S. fuscum*, 6.

b. *Stem leaves distinctly narrowed above.*

Branch leaves when dry plainly 5 ranked *S. quinquefarium*, 7.

¹ *S. strictum* Lindb.

Branch leaves when dry not 5 ranked.

Branch leaves with isolated pores on inner face toward apex.

Hyaline cells of stem leaves mostly with fibrils, branch leaves not shining.

Stem leaves extended into a blunt point with involute margins *S. tenerum*, 8.

Stem leaves scarcely pointed, not involute *S. acutifolium*, 13.

Hyaline cells of stem leaves mostly with fibrils, branch leaves silky shining *S. subnitens*, 9.

Branch leaves with numerous pores upon entire inner face *S. microphyllum*, 10.

B. *Branch leaves toothed at apex and upper margins; border with resorption-furrows.*

Stem leaves small (1.14-1.28 mm.), narrow border plainly broadened near the base *S. Labradorense*, 11.

Stem leaves larger (2-2.10 mm.), narrow border scarcely broader at the base *S. molle*,¹ 18.

II. Squarrosa.

Dioicous, perigonal leaves scarcely different from the leaves of the sterile branches *S. teres*, 16.

Monoicous, perigonal leaves smaller, curved, erect spreading *S. squarrosus*, 16.

III. Cuspidata.

A. *Branch leaves without fibrils.*

Pits on the outer surface large (10-12 μ diam.), 5-16 in a single row in the middle of cell *S. macrophyllum*, 24.

Pits on the outer surface smaller (4-5 μ diam.), 40-60 in each hyaline cell *S. Floridanum*, 12.

B. *Branch leaves always with fibrils.*

i. *Branch leaves ovate-lanceolate or lanceolate to almost long linear.*

a. *Stem leaves lacerate-fimbriate in the upper part.*

Stem leaves widened upward, spatulate, fringed at apex *S. Lindbergii*, 15.

Stem leaves triangular linguiform, lacerate two-cleft at the apex *S. riparium*, 13.

¹ Including *S. Muelleri* Sch., L. & J. Man. 17.

b. *Stem leaves entire or with isolated gaps at the tip only.*

Chlorophyllose cells on the outer face of the leaf, triangular in cross section.

Cortex plainly distinct from wood-cylinder, branch leaves almost always without pores on the inner face

S. Mendocinum, 20.

Cortex indistinct, branch leaves with numerous rounded pores on the inner face

S. recurvum,¹ 15.

Chlorophyllose cells exposed on both faces, quadrate or trapezoidal.

Borders of the branch leaves involute, pores of the outer face very small, in the upper cell corners

S. cuspidatum, 14.

Borders of the branch leaves not involute, pores of the outer face large (6 μ), in 1 or 2 rows

S. Dusenii, 14.

Borders of the branch leaves not involute, pores generally absent

S. Fitzgeraldii, 23.

2. *Branch leaves small, oval or long ovate*

S. molluscum,² 20.

IV. *Polyclada.*

This section has only one species

S. Wulfianum, 16.

V. *Rigida.*

Chlorophyllose cells of the branch leaves spindle-shaped in cross section, hyaline cells with papillae on the wall bordering the chlorophyllose cells

S. Garberi, 18.

Chlorophyllose cells elliptical in cross section, smooth on the wall bordering the chlorophyllose cells

S. compactum,³ 17.

VI. *Subsecunda.*A. *Stem and branch leaves without pores*

S. Pylalei,⁴ 26.

B. *Branch leaves always porose.*1. *Branch leaves with few pores on both faces, never in uninterrupted rows.*

Stem leaves enlarged from base upwards, upper margins involute, slightly toothed

S. Mohrianum, 15.

¹ *S. intermedium* Hoffm., L. & J. Man. 15.

² *S. tenellum* Ehrh., L. & J. Man. 20.

³ *S. rigidum* Sch., L. & J. Man. 17.

⁴ Including *S. sedoides* Brid., L. & J. Man. 23.

Stem leaves not enlarged from the base upwards, apex frequently cucullate and fringed.

- Hyaline cells not divided *S. obesum*, 16.
 Hyaline cells with 1-2 cross walls *S. dasyphyllum*, 17.

2. *Branch leaves with few pores on the inside, outside with numerous pores in bead-like rows.* (3)

a. *Chlorophyllose cells of branch leaves isosceles-triangular, or trapezoidal in cross section; mostly exposed on the inner face* *S. microcarpum*, 18.

b. *Chlorophyllose cells of branch leaves quadrate or barrel-shaped, central and exposed on both faces.* (c)

Stem mostly simple, rarely a single divergent branch *S. cyclophyllum*, 22.
 Stem fasciculate-branched.

Cortex of one or (in part) two cell layers *S. subsecundum*, 19.
 Cortex of two to several cell layers.

Hyaline cells with abundant fibrils, stem leaves large (1.3-1.4 mm. long) *S. platyphyllum*, 19.

Hyaline cells without fibrils, or with fibrils only at apex, stem leaves smaller (.86 mm. long) *S. contortum*,¹ 19.

Chlorophyllose cells triangular or trapezoidal, exposed on the outer face.

Branch leaves ovate-lanceolate, plicate *S. plicatum*, 20.
 Branch leaves round ovate, not plicate *S. Orlandense*, 21.

3. *Branch leaves with abundant pores on both faces.*

Chlorophyllose cells triangular or trapezoidal in cross section, mostly exposed on outer face (sometimes on both).

Hyaline cells much divided, chlorophyllose cells triangular *S. Mobilense*, 22.

Hyaline cells once divided, chlorophyllose cells trapezoidal, free on both faces *S. simile*, 23.

Chlorophyllose cells quadrate or barrel-shaped, central or exposed on both faces *S. rufescens*, 24.

¹ *S. larictinum* Spruce, L. & J. Man. 19.

S. obesum, 16.
S. dasyphyllum, 17.

side with numer-

les-triangular, or
osed on the inner
S. microcarpum, 18.

or barrel-shaped,
(c)

S. cyclophyllum, 22.

S. subsecundum, 19.

ge
S. platyphyllum, 19.
apex,

S. contortum, 19.

, exposed on the

S. plicatum, 20.

S. Oriandense, 21.

both faces.

ction,

S. Mobilense, 22.

S. simile, 23.

S. rufescens, 24.

VII. *Cymbifolia*.

A. *Chlorophyllose cells of branch leaves exposed on the inner face.*

Branch leaves minutely fringed all around . . . *S. Portoricense*, 22.
Branch leaves not fringed.

Walls of the hyaline cells adjoining the chlorophyllose
cells furnished with a fringe of fibrils . . . *S. imbricatum*, 21.

Walls of the hyaline cells adjoining the chlorophyllose
cells thickly papillose . . . *S. papillosum*, 21.

Walls smooth *S. cymbifolium*, 21.

B. *Chlorophyllose cells of branch leaves exposed on both faces or included.*

[*S. cymbifolium* may be sought here.]

Cortical cells with fibrils.

Chlorophyllose cells free on both faces, wood-cylinder
dark-brown to black *S. Waghornii*, 23a.

Chlorophyllose cells included, wood cylinder red . . . *S. medium*, 24a.

Cortical cells not fibrillose *S. Ludovicianum*, 25.

ANDRÆA, p. 25.

I. *Leaves ecostate.*

Leaves papillose beneath.

Margin hyaline.

Leaves incurved, minute, rotund obtuse, bi-ventri-
cose *A. parvifolia*, 26.

Leaves spreading or secund, acuminate, not ventri-
cose *A. petrophila*, 25.

Leaves spreading, obtuse, short-pointed *A. papillosa*, 27.

Margin not hyaline *A. alpestris*, 28.

Leaves not papillose, upper leaf-cells rhomboid . . . *A. obovata*, 29.

II. *Leaves costate.*

Perichaetial and comal leaves different.

Costa vanishing below apex *A. rupestris*, 25.

Costa excurrent.

Subula papillose.

Costa filling all the subula *A. crassinervis*, 25.

Costa filling half the subula *A. Huntii*, 30.

Subula smooth *A. Blyttii*, 31.

¹*S. Austriacum* Sull., L. & J. Man. 21.

Perichaetial and comal leaves similar.

- Leaves faintly papillose, costa narrow, sometimes ex-
current *A. Macounii*, 35.
Leaves strongly papillose, costa 30-40 μ wide, percurrent *A. nivalls*, 33.

ARCHIDIUM, p. 49.

Autoicous.

- Costa reaching to point of leaf *A. Ohioense*, 50.
Costa often long excurrent *A. Hallii*, 51.¹

Paroicous.

- Leaves serrulate *A. tenerrimum*, 50.
Leaves quite entire.
Cells oval or rhombic *A. Ravenelii*, 50
Cells quadrangular or quadrate *A. longifolium*, 50.

MICROMITRIUM, p. 37.

- Spores 63 μ diameter, leaves serrate *M. megalosporum*, 37.
Spores 25 μ diameter, nearly smooth, leaves serrate above *M. Austini*, 37.
Spores a little smaller, papillose, leaves nearly entire *M. synoicum*, 37.

EPHEMERUM, p. 37.

- Leaves not costate *E. serratum*, 37.
Leaves costate.
Costa ending below or at apex *E. coherens*, 39.
Costa excurrent.
Seta short, capsule acutely beaked *E. stenophyllum*, 39.
Seta wanting, capsule blunt pointed.
Leaves gradually long-accumbent, slightly and
irregularly serrate at apex *E. crassinervium*, 38.
Leaves with a long hyaline spinulose arista *E. spinulosum*, 38.
Leaves papillose both sides *E. papillosum*, 38.
Leaves long-spinulose on both sides *E. hystrix*, 38.

SPHÆRANGIUM, p. 40.

- Leaves papillose on both faces *S. Schimperianum*, 41.
Leaves smooth, or papillose on back.
Margins reflexed, plants triquetrous *S. triquetrum*, 41.
Margins almost plane, plants round or tetragonal.
Lower leaves ecostate *S. rufescens*, 40.
Lower leaves costate *S. muticum*, 40.

¹See also Appendix, no. 34.

ex-
A. Macounii, 38.
current A. nivalis, 33.

A. Ohioense, 50.
A. Hallii, 51.¹

A. tenerrimum, 50.

A. Ravenellii, 50
A. longifolium, 50.

M. megalosporum, 37.
M. Austlini, 37.
M. synolcum, 37.

E. serratum, 37.

E. coherens, 39.

E. stenophyllum, 39.

and
E. crassinervium, 38.

E. spinulosum, 38.

E. papillosum, 38.

E. hystrix, 38.

Schimperlanum, 41.

S. triquetrum, 41.

S. rufescens, 40.

S. muticum, 40.

PHASCUM, p. 41.

- Capsule sub-globose, apiculate.
Leaf margins plane or incurved, denticulate . . . P. Carniolicum, 42.
Leaf margins reflexed, quite entire.
Capsule immersed P. cuspidatum, 42.
Capsule sub-exserted P. subexsertum, 35.
Capsule ovate- or oblong-lanceolate P. bryoides, 42.

PLEURIDIUM, p. 43.

- Inflorescence parvicous.
Costa reaching the obscurely serrate apex . . . P. subulatum, 43.
Costa excurrent into a smooth awl-shaped point . . . P. Ravenellii, 43.
Inflorescence autocious.
Upper leaves long subulate.
Entire or apex denticulate or serrulate . . . P. alternifolium, 44.¹
Serrulate from middle upward P. Bolanderi, 44.
Upper leaves abruptly short pointed P. Sullivantii, 44.

BRUCHIA, p. 45.²

- I. Collum none B. palustris, 45.

II. Collum present.

- A. Exceeding the sporangium B. longicollis, 38.

B. Shorter than or equaling sporangium.

i. Capsule exserted.

- Leaves short, not subulate.
Spores papillose B. Hallii, 47.
Spores pitted.
Costa percurrent B. brevifolia, 48.
Costa vanishing below apex B. fusca, 39.
Leaves subulate.
Smooth or sub-papillose.
Costa filling point B. curviseta, 47.
Costa not filling point, apex serrulate.
Spores spinose.
Capsule ovoid, neck short B. flexuosa,¹ 46.
Capsule elongated, neck long B. Sullivantii, 46.

¹ See also Appendix, no. 38.

² See revision by Eliz. G. Britton, Bull. Torr. Bot. Club 21: 243. 1894.

| | |
|--------------------------------|-------------------|
| Spores reticulate | B. Texana, 48. |
| Spores papillose | B. Bolanderi, 46. |
| Distinctly papillose | B. Donnellii, 48. |

2. Capsule immersed.

| | |
|------------------------------|--------------------|
| Spores pitted | B. Caroline, 40. |
| Spores reticulate. | |
| Calyptra papillose | B. Ravenelli, 49. |
| Calyptra smooth | B. Drummondii, 48. |

B. Beyrichiana Müll. is rejected as mixed and uncertain; *B. Hamp-
eana* Müll. is a Chilean species.

ASTOMUM, p. 51.

| | |
|---|-------------------------|
| Leaves not crispate when dry. | |
| Capsule chestnut-brown, ovoid | A. nitidulum, 52. |
| Capsule orange, oval | A. nitidulum, var., 52. |
| Leaves crispate when dry. | |
| Capsules often clustered (2-3), oblong oval | A. Ludoviclanum, 52. |
| Capsule solitary. | |
| Brown globose, capsule immersed | A. crispum, 51. |
| Brown globose, capsule exerted | A. Drummondii, 41. |
| Orange, sub-globose, immersed | A. Sullivantii, 4 52. |

GYMNOSTOMUM, p. 52.

| | |
|---|---------------------|
| Lid long remaining attached to columella, capsule thick- walled, with 6-8 rows of transversely elongated cells at the mouth | G. curvirostre, 53. |
| Lid falling early, capsule thin-walled, with 3-4 rows of transversely elon- gated cells at mouth. | |
| Plants 1-2 mm. high, lid conic | G. tenue, 54. |
| Plants 5-10 mm. high, lid subulate, costa 24-35 μ wide at base with 2 guides ¹ | G. calcareum, 5 53. |
| Plants 1-7 cm. high, costa 70 μ wide at base, with 4-6 guides ² | G. rupestre, 53. |

¹ Incl. *B. brevicollis* L. & J. Man. 47.

² *B. Ravenelli mollis* L. & J. Man. 49. Fuller description, Appendix no. 40.

³ *B. brevipes* L. & J. Man. 48. *B. brevipes* Hook. is African.

⁴ *A. Sullivantii* is probably a variety of *A. crispum*. Most American specimens seem to be referable to *A. Sullivantii*. (Ren. & Card.)

⁵ As far as can be determined *G. platyphyllum* Kindb. (Appendix no. 44) falls here.

⁶ See explanation under *Dicranum*, p. 181.

B. Texana, 48.
 B. Bolanderi, 46.
 B. Donnellii, 48.

B. Carolinae,³ 40.

B. Ravenelli, 49.
 B. Drummondii,³ 48.
 uncertain; B. Hamp-

A. nitidulum, 52.
 nitidulum, var., 52.

Ludoviclanum, 52.

A. crispum, 51.
 A. Drummondii, 41.
 A. Sullivanii,⁴ 52.

ck-
 ells
 G. curvirostre, 53.
 f transversely elon-

G. tenue, 54.
 de
 G. calcareum,⁵ 53.
 -6
 G. rupestre, 53.

ndix no. 40.

merican specimens seem

x no. 44) falls here.

ANCECTANGIUM, p. 54.

Leaves long-lanceolate, subulate pointed A. Peckii, 55.
 Leaves lanceolate, acute A. compactum, 45.

WEISIA, p. 55.

Inflorescence autoicous.
 Leaf margins involute, costa stout, excurrent W. viridula, 55.¹
 Leaf margins not involute, costa thin, vanishing in the
 acumen W. convoluta, 48.
 Inflorescence dioicous.
 Teeth large, lacunose and bifid, capsule 8-sulcate W. longiseta,² 56.
 Teeth truncate, capsule not sulcate W. Wolfii, 57.

DICRANOWEISIA,³ p. 57.

Leaf cells at base thick-walled, linear (1:6-10) D. crispula, 57
 Leaf cells at base thin-walled, rectangular (1:2-3) D. cirrhata, 57.

RHABDOWEISIA, p. 58.

Leaves minutely denticulate or entire; teeth filiform,
 smooth, fugacious R. fugax, 59.
 Leaves coarsely dentate; teeth linear, obliquely crossed-
 striate R. denticulata, 59.

CYNODONTIUM, p. 59.

Annulus very narrow and persistent or none.
 Teeth bifid or trifid, pale C. subalpestre, 51.
 Teeth bifid, red to purple.
 Collum indistinct or none C. gracilescens, 60.
 Collum short, inflated and strumose C. virens, 61.
 Teeth not bifid, purple C. Schisti, 59.
 Annulus distinct.
 Leaves long taper-pointed, apex serrate C. polycarpum, 60.
 Leaves more obtuse, apex crenulate C. strumulosum, 52.

¹ See also Appendix, nos. 49, 50.

² From the examination of specimens of this species from Florida and Louisiana, Renaud & Cardot conclude that it is only a variety of the most variable and polymorphous *W. viridula*.

³ *D. obliqua* Kindb. Mac. Cat. 256 is *nomen nudum*.

DICHODONTIUM, p. 61.

- Capsule strumose *D. Olympicum*, 54.
 Capsule not strumose.
 Costa vanishing below apex, seta yellow *D. pellucidum*,¹ 62.
 Costa percurrent, seta red *D. Canadense*, 62.

TREMATODON, p. 62.

- Leaves lance-subulate.
 Collum equaling or somewhat exceeding the oval oblong
 sporangium, teeth deeply bifid *T. ambiguum*, 63.
 Collum greatly exceeding the cylindrical sporangium,
 teeth perforate *T. longicollis*, 63.
 Leaves ovate, short pointed, teeth entire or perforate *T. brevicollis*, 55.

DICRANELLA, p. 64.

- I. Cells of the exothecium rectangular quadrate; seta red; costa usually narrow and well defined below.

A. Leaves not sheathing, erect-spreading.

- Costa percurrent or excurrent.
 Annulus none, peristome papillose.
 Leaves short acuminate, blunt, costa $\frac{1}{4}$ width of leaf
 at base *D. Langloisii*, 53.
 Leaves long acuminate, costa $\frac{1}{4}$ width of leaf at base *D. Howei*, 61.
 Leaves long acuminate, acute, costa $\frac{1}{4}$ width of leaf at base.
 Capsule cernuous *D. varia*, 65.
 Capsule erect, symmetric *D. rufescens*, 66.
 Annulus present, peristome not papillose.
 Teeth orange, pale at the apex, costa excurrent *D. parvula*, 57.
 Teeth purple, costa percurrent *D. leptotrichoides*, 59.
 Costa ceasing within the apex, annulus large, simple *D. debilis*, 66.

B. Leaves from a sheathing base, squarrose.

- Broad, obtuse *D. squarrosa*, 65.
 Abruptly subulate.
 Capsule striate, substrumose, leaf apex entire *D. Grevilleana*, 64.
 Capsule not striate nor strumose, leaf apex serrulate *D. Schreberi*, 64.²

¹ See also Appendix, no. 53.² As near as can be determined *D. laxiretis* R. & C. falls here. See Appendix, no. 62, 63.³ See Appendix, no. 56.

II. *Cells of the exothecium prosenchymatous; seta often yellow; costa usually broad and indistinct below.*

A. *Seta red.*

- Leaves from a sheathing base, squarrose *D. crispa*, 64.
 Leaves not sheathing nor squarrose.
 Mostly erect, capsule cernuous *D. subulata*, 66.
 Secund, capsule erect *D. curvata*, 67.

B. *Seta yellowish.*

- Capsule symmetric, erect *D. Fitzgeraldi*, 60.
 Capsule cernuous.
 Strumose *D. cerviculata*, 65.
 Not strumose *D. heteromalla*, 66.

DICRANUM, p. 67.¹

In this genus the structure of the costa is of diagnostic value. It is either composed of similar cells (*homogeneous*), or composed of large parenchyma cells and small sclerenchyma cells (*stereids*). The large parenchyma cells (*guides* = "Deuter" of Lorentz²) form a row (seldom double) in the middle of the costa touching each other tangentially. They are comparatively large, but little thickened and either empty or starch-bearing.³

I. *Monoicous, stems radiculose only at base, costa long excurrent, homogeneous.*

- Capsule erect, not strumose.
 Striate and furrowed when dry *D. hyperboreum*,⁴ 64.
 Neither striate nor furrowed when dry *D. fulvellum*, 68.
 Capsule cernuous, strumose.
 Leaf cells not papillose, capsule oblong-cylindric *D. Starkii*, 68.
 Leaf cells with papillae over partitions, capsule short-ovate.
 Leaves fringed-secund *P. falcatum*,⁵ 68.
 Leaves spreading *D. Blyttii*, 68.

¹ Arranged by Dr. RODNEY H. TRUE.

² Pringsheim's *Jahrb. f. wissenschaft. Bot.* 3: 574.

³ Limpricht: *Die Laubmoose* 23.

⁴ See Appendix, no. 65.

⁵ See Appendix, no. 68a.

D. Olympleum, 54.

D. pellucidum,¹ 62.

D. Canadense, 62.

oblong

T. ambiguum, 63.

gium,

T. longicollis, 63.

T. brevicollis, 55.

ate; seta red; costa
below.

reading.

f leaf

D. Langloisii, 58.

base *D. Howei*, 61.

leaf at base.

D. varia, 65.

D. rufescens, 66.

D. parvula, 57.

leptotrichoides, 59.

D. debilis, 66.

quarrose.

D. squarrosa, 65.

D. Grevilleana, 64.

D. Schreberi, 64.²

See Appendix, no. 62, 63.

- II. *Monoicous, eradiculoae, costa vanishing, guides present.* (III) **D. molle, 66.**
- III. *Dioicous, stems subradiculoae above, costa very broad, 2-4-stratose, superficial cells thin, without chlorophyll, capsule erect, regular.* (IV.)

Costa one-half leaf-width at base.

- Costa not furrowed at back, smooth **D. albicans, 71.**
- Costa furrowed and toothed at back **D. longifolium, 70.¹**
- Costa one-fifth to one-fourth leaf-width at base **D. Sameri, 68.²**

- IV. *Dioicous, stems radiculoae, often densely so, costa with median guides.*

A. *Capsule cernuous, more or less arcuate.*

i. *Leaf cells pitted.*

- a. *Costa not reaching apex, leaves mostly transverse, undulate.*

Leaf cells above elongated.

- Costa serrate at back, not lamelloae **D. Bonjeani, 74.³**

Costa with serrate lamellae.

- Capsules clustered, perichaetial leaves differentiated **D. undulatum, 76.**

- Capsules solitary (?), perichaetial leaves like others **D. dipteroneuron, 75.⁴**

- Capsules solitary, perichaetial leaves tubulose, abruptly narrowed **D. Bonjeani, 74.³**

Leaf cells above isodiametric.

- Margin smooth to near tip **D. brachycaulon, 88.**

Margin serrate or denticulate.

Upper costa and lamina rough at back.

- Capsules clustered, leaf margins sharply serrate **D. Drummondii, 76.**

- Capsules solitary, leaf margins finely denticulate above widened base **D. spurium, 75.**

- Costa and lamina smooth at back, capsules solitary, margins irregularly denticulate in upper half **D. Bergeri, 77.⁵**

¹ See Appendix, no. 67.

² See Appendix, no. 68.

³ *D. patustre* La Pyl., L. & J. Man. See Appendix, nos. 85-87.

⁴ Ex descr. probably = *D. Bonjeani* DeNot.

⁵ *D. Schraderi* W. & M., L. & J. Man. 75.

ides

D. molle, 66.very broad, 2-4-strat-
rophyll,*D. albicans*, 71.*D. longifolium*, 70.¹*D. Santeri*, 68.²

costa with median

uate.

verely undulate.

D. Bonjeani, 74.³*D. undulatum*, 76.pteronenron, 75.⁴

ptly

D. Bonjeani, 74.³*D. brachycaulon*, 88.*Drummondii*, 76.*D. spurium*, 75.

7,

D. Bergeri, 77.⁵b. *Costa percurrent or excurrent, leaves not undulate.*[*D. Bonjeani* may be sought here.]Guides in two rows *D. majus*, 74.

Guides in one row.

Margin serrulate to middle of leaf, back of costa lamellose.

Seta yellow, capsule long and narrow (1:5-6) *D. consobrinum*, 84.Seta red, capsule broader (1:4) *D. scoparium*, 73.¹

Margin entire to near apex, back of costa not lamellose.

Cells above narrow (1:5-6) leaves brittle *D. neglectum*, 73.Cells above isodiametric, leaves not brittle *D. brachycaulon*, 88.2. *Leaf cells not pitted or faintly so.*

Leaves quite entire or denticulate, subulate.

Points very brittle, mostly broken *D. fragillifolium*, 73.Points not broken *D. elongatum*, 71.

Leaves entire, upper obtuse.

Cells throughout elongated, thickened, pitted *D. Groenlandicum*, 69.Cells short, quadrate above, not pitted *D. Miquelonense*, 72.

Leaves serrulate.

Upper cells very irregular.

Capsules cylindrical, striped or striated.

Costa at point of greatest leaf-width one-sixth or
more *D. Muhlenbeckii*, 72.²

Costa at point of greatest leaf-width one-tenth or less.

Cells of upper third of lamina short (8-10 ×
10-17 μ) *D. pallidum*, 76.Cells of upper third of lamina narrower, larger,
(15 × 50 μ) *D. rhabdocarpum*, 73.Capsules ovate-oblong, not striped or but faintly *D. congestum*, 72.³Upper cells regular, capsule obovate-oblong, striped *D. fuscescens*, 72.B. *Capsule erect, symmetric.*Costa without stereids *D. strictum*, 69.¹ See Appendix, nos. 77-83.² See Appendix, no. 74.³ *D. fuscescens* L. & J. Man. 72 in part.⁴ *D. fuscescens longirostre* and *angustifolium* L. & J. Man. 72. See Appendix, nos. 70, 71.

Costa with two stereid bands.

Lamina above more or less bi-stratose.

Margin and costa serrulate *D. fulvum*, 70.

Margin entire, apex usually broken *D. viride*, 69.

Lamina throughout uni-stratose.

Upper cells rectangular and mamillose *D. montanum*, 69.

Upper cells less regular, not mamillose *D. flagellare*, 70.

DICRANODONTIUM, p. 77.

Cells at the basal angles enlarged.

Quadrate, teeth deeply bifid *D. Millspaughii*,¹ 90.

Rectangular, teeth bifid to base *D. longirostre*, 77.

Cells at the basal angles not enlarged *D. Virginicum*, 89.

CAMPYLOPUS, p. 77.

I. *Costa smooth at back.*

A. *Auricles none.*

[*C. gracilicaulis* may belong here.]

Cells of the costa uniform in transverse section *C. Leanus*, 78.

Cells of the costa unlike in transverse section, stereids forming 2-3 dorsal layers.

Hyaline cells superior, in one row *C. Schimperl*, 91.

Hyaline cells superior, in two rows *C. Henricl*, 92.

B. *Auricles present.*

No lamina except small colored auricles *C. Hallii*, 79.

Lamina distinct.

Perichaetial leaves concolorous.

Auricles brown, plane, decurrent *C. Talliensis*, 78.

Auricles whitish, large.

Leaves serrulate at apex *C. subleucogaster*, 79.

Leaves spinulose serrate at apex *C. Donnellii*, 79.

Auricles dirty red *C. angustiretis*, 80.

Perichaetial leaves with hyaline points (may include *C. Donnellii* and *C. angustiretis*) *C. gracilicaulis*, 80.

¹ *Campylopus flexuosus* L. & J. Man. 78, not of Bridel. *C. flexuosus* Brid. is not found in America.

II. *Costa scabrous or lamellose at back.*

- Leaves with pellucid hair points C. *introflexus*, 78.
 Leaves not hair pointed.
 Alar cells round, lamina wanting C. *frigidus*, 79.
 No auricles C. *virginicus*, 80.

FISSIDENS, p. 81 (incl. *Conomitrium*, p. 89.¹)I. (EUFISSIDENS.) *Plants terrestrial or submersed but not floating; leaves soft, one layer of cells.*A. *Fruit terminal.*1. *Monoicous, male flowers axillary.*[*F. falcatus* may be sought here.]

- Leaf-cells small, densely chlorophyllose, in distinct rows F. *limbatus*, 82.
 Leaf-cells large, not densely chlorophyllose, nor in distinct rows F. *bryoides*, 81.

2. *Dioicous or monoicous with the male flowers terminal on a rooting branch at the base of the female stem.*a. *Leaf-cells 1½-2 times as long as wide, large, distinct.*

- Plants less than 1 mm. high, leaves two or three pairs F. *closteri*, 81.
 Plants 2-4 mm. high, wholly hyaline, leaves 3-5 pairs F. *hyalinus*, 84.

b. *Leaf-cells almost or quite isodiametric, often obscure.*[*F. limbatus* may be sought here.]

Leaves with a narrow border, at least on vaginant lamina.

Marginal leaf-cells not papillose.

Leaves acute, costa percurrent.

Vaginant lamina not reaching middle of leaf, peristome arising below the mouth F. *pusillus*, 84.Vaginant lamina reaching middle of leaf, peristome not arising below the mouth F. *incurvus*,² 82.Leaves obtuse, costa vanishing below apex F. *obtusifolius*, var. 85.¹ See Barnes: Bot. Gaz. 12: 1. 1882.² See Appendix, no. 93.

- Marginal leaf-cells papillose.
 Costa percurrent F. Ravenelli, 85.
 Costa ceasing below apex F. Garberi, 86.
 Leaves without a border.
 Acute, cells densely chlorophyllose, obscurely papillose F. Donnellii, 85.
 Acute, almost hyaline, smooth F. pauperculus, 99.
 Obtuse, cells pellucid, operculum conic F. obtusifolius, 86.
 Apiculate, operculum with acicular beak F. osmundoides, 87.
 Leaves with a thick reddish border. Plants submersed,
 rigid F. rufulus, 84.¹

B. *Fruit lateral.*

1. *Leaves without a border.*

- Obtuse, entire, plants 2-5 cm. high, fruit sub-terminal F. polypodifolides, 88.
 Rounded at apex, irregularly serrate, 1-2 cm. high, fruit
 sub-basal F. sub-basilaris, 88.
 Mucronate, regularly serrulate, fruit basal or sub-basal F. taxifolius,² 87.
 2. *Leaves bordered by several rows of paler, often incrassate, cells.*

Leaf margin serrulate.

- Leaf cells obscure ($6-9 \times 6-12\mu$) F. cristatus, Wils.,³ 87.
 Leaf cells distinct ($12-15 \times 18-24\mu$) F. adiantoides,⁴ 88.
 Leaves entire or slightly denticulate at the apex.
 Inferior lamina bordered, ceasing abruptly at base F. Floridanus, 83.
 Inferior lamina not bordered, tapering below F. falcatus, 98.

II. (PACHYFISSIDENS.) *Leaves rigid, composed of more than one layer of cells, opaque.*

- Plants growing in water or very wet places F. gradifrons, 89.

III. (OCTODICERAS.)⁵ *Plants aquatic, filiform, floating.*

- Plants large, much branched, pedicel shorter than the
 capsule F. Julianus, 89.
 Plants small, little branched, pedicel longer than the
 capsule F. Hallianus, 90.

¹ *F. ventricosus* of L. & J. Man.

² See Appendix, no. 96.

³ *F. deceptens* De Not., L. & J. Man.

⁴ See Appendix, no. 97.

⁵ *Conomitrium* of Manual.

F. Ravenelli, 85.
F. Garberl, 86.
F. Donnelli, 85.
F. pauperulus, 99.
F. obtusifolius, 86.
F. osmundoides, 87.
 ed,
F. rufulus, 84.¹

polypodioides, 88.
 nit
sub-basilaris, 88.
F. taxifolius,² 87.

incrassate, cells.

status, Wils.,³ 87.
adiantoides,⁴ 88.

F. Floridanus, 83.
F. falcatus, 98.

l of more than

F. gradifrons, 89.

n, floating.

F. Julianus, 89.

F. Hallianus, 80.

In the Revision of N. A. species of Fissidens,¹ *FF. inconstans, exiguus* and *minutus* were reduced to *F. incurvus*, the latter two forming varieties. *Fr. bryoides caespitans, crassipes, Hallii* and *Texanus* are relegated to the list of doubtful species. *FF. Bambergerti* Schimp. I regard as a form of *F. incurvus*; *F. viridulus* is a form, possibly a subspecies, of the same. It may be known by its thin-walled capsule, with the peristome inserted below the mouth. Neither are worthy of a distinct place in the key. "*F. tamarindifolius* Don" (Mac. Cat. 36), seems to be *F. tamarindifolius* Brid. which is a variety of *F. incurvus*.

LEUCOBRYUM,² p. 90.

Capsule apparently lateral (by innovations), leaves ovate lanceolate.

Robust, tufts 4-6 cm. deep *L. glaucum* (L.) Sch.,³ 90.

Smaller tufts scarcely 2 cm. deep *L. glaucum albidum*,⁴ 91.

Capsule exactly terminal, leaves squarrose, very short and very broad *L. minus*,⁵ 91.

CERATODON, p. 92.

Costa percurrent or vanishing below apex.

Capsule distinctly strumose, articulations of teeth few *C. heterophyllus*, 101.

Capsule not distinctly strumose, articulations of teeth many *C. purpureus*, 92.

Costa long excurrent, teeth articulate to the middle.

Leaf margin entire *C. conicus*, 100.

Leaf margin serrulate toward apex *C. minor*, 92.

TRICHODON, p. 92.

Cells of the leaf base linear, above rectangular or indistinct

T. cylindricus, 93.

Cells of leaf base rectangular (1:2-4), above quadrate

T. flexifolius, 102.

DISTICHIMUM, p. 93.

Costa long excurrent.

Capsule erect, spores 17-20 μ *D. capillaceum*, 93.

Capsule cernuous, spores 30-44 μ *D. inclinatum*, 94.

Costa percurrent or vanishing below the apex *D. Macounii*, 106.

¹ See Barnes: Bot. Gaz. 13: 1. 1889.

² See E. G. Britton, Bull. Torr. Bot. Club 19: 189. 1892.

³ *L. vulgare* L. & J. Man. 90.

⁴ *L. albidum* (Brid.) Lindb. *L. minus* Sulliv. non Hampe.

⁵ *L. sediforme* L. & J. Man. 91.

SELIGERIA, p. 96.

- Seta straight when moist.
 Leaves sharp pointed, cells above rectangular, spores
 10-14 μ *S. pusilla*, 96.
 Leaves blunt-pointed, cells above quadratic, spores
 14-18 μ *S. calcarea*, 97.
 Leaves mostly blunt-pointed, cells rectangular, spores
 24-32 μ *S. tristicha*, 97.
 Seta arcuate when moist.
 Leaves long-subulate, costa excurrent *S. recurvata*, 97.
 Leaves sub-linear, obtuse, costa not excurrent *S. campylopoda*, 107.

POTTIA, p. 100.

I. *Peristome wanting or rudimentary.*

- Costa with 2-4 lamellae above *P. cavifolia*, 101.
 Costa not lamellate.
 Leaf margins more or less revolute.
 Lid conic obtuse, spores echinate *P. minutula*, 101.
 Lid rostellate, spores papillose.
 Calyptra smooth.
 Capsule obovate spherical (1:1) or sub-hemi-
 spherical *P. truncata*, 101.
 Capsule oblong-oval (1:2).
 Leaves minutely papillose toward apex, basal
 cells 1:4 *P. intermedia*, 110.
 Leaves smooth, basal cells 1:5-6 *P. littoralis*, 111.
 Calyptra scabrous *P. Wilsoni*, 101.
 Leaf margins plane or involute.
 Lid abruptly rostrate, leaves sharply serrate above *P. Helmii*,¹ 102.
 Lid conic, leaves distantly denticulate above *P. riparia*, 102.
 Lid conic-subulate, leaves slightly crenulate above *P. barbula*, 102.

II. *Peristome distinct.*

- Leaves oblong-lanceolate, margins revolute *P. Starkeana*, 103.
 Leaves rounded or round-spatulate, margins plane *P. latifolia*, 103.

¹*P. helmtoides* Kindb. (Appendix, no. 109), evidently falls here. The great resemblance of the two makes the specific value of *P. helmtoides* extremely doubtful.

DIDYMODON, p. 104.

- Leaf cells throughout quadratic *D. luridus*, 104.
- Leaf cells below rectangular.
- Inflorescence synoicous *D. rubellus*, 104.
- Inflorescence dioicous.
- Lid conic, leaves denticulate *D. Baden-Powellii*, 113.
- Lid rostrate.
- Basal cells thick-walled.
- Lamina and costa long papillose *D. rufus*, 116.
- Lamina and costa minutely papillose *D. Hendersoni*, 114.
- Basal cells thin walled.
- Perichætal leaves gradually narrowed *D. cylindricus*, 105.
- Perichætal leaves abruptly attenuate *D. Canadensis*, 112.

DITRICHUM, p. 105.¹

- Dioicous.
- Leaves slightly twisted.
- Stem leaves spreading, perichætal leaves hardly sheathing *D. tortile*, 105.
- Stem leaves imbricate, perichætal leaves long sheathing *D. vaginans*, 106.
- Leaves not twisted.
- Plants short (1-2 cm.), not radiculose.
- Teeth without a basilar membrane *D. homomallum*, 107.
- Teeth with a broad basilar membrane *D. ambiguum*, 104.
- Plants long (to 10 cm.), densely radiculose. *D. flexicaule*, 107.²
- Monocous.
- Plants short (5-10 mm.).
- Teeth cylindric, legs unequal, nodose-articulate, leaves spreading *D. pallidum*, 107.
- Teeth cylindric, legs equal, obscurely and distinctly articulate, leaves spreading *D. montanum*, 109.
- Teeth flattened, linear, trabeculate, perforate, leaves secund *D. Schimperii*, 108.
- Plants longer (2-3 cm.), glaucous *D. glaucescens*, 108.

¹ *Leptotrichum* Hampe, L. & J. Man, 105.
² *Leptotrichum brevifolium* Kindb., Mac. Cat. 46, is described as a subspecies of *L. flexicaule* Hampe. The characters given however are not sufficient to justify a place in this key. See Appendix, no. 108.

TRICHOSTOMUM, p. 108.

I. *Lamina composed of one layer of cells, papillose.*

Margin reflexed or undulate, entire.

Annulus wanting *T. tophaceum*, 109.Annulus large, compound *T. pyriforme*, 109.

Margin plane or incurved.

Costa reaching apex or excurrent; serrate above.

Base of leaf yellowish, with thick walled rectangular cells *T. crispulum*, 109.

Base of leaf hyaline.

Abruptly mucronate or obtuse, with long papillae *T. flavo-virens*, 109.Gradually acuminate, papillae low *T. nitidum*, 118.Costa ceasing far below apex; entire *T. Coloradense*, 413.II. *Lamina of two layers, upper surface mamilliose, lower smooth.*Costa excurrent, leaves denticulate above *T. Vancouveriense*, 119.

Costa percurrent or vanishing below apex, serrate above.

Peristome not twisted, seta arcuate or variously bent *T. flexipes*, 110.Peristome twisted, seta subflexuous *T. anomalum*, 110.

DESMATODON, p. 110.

I. *Capsule erect or nearly so.*A. *Leaves without a hyaline or thickened border.*1. *Costa excurrent into a hair.*a. *Papillose.*

Capsule oblong (1:2 or 1:3 excl. lid), 16 teeth divided nearly or quite to base.

Plants of mountainous regions; calyptra reaching base of capsule *D. latifolius*, 111.Plants of lowlands; calyptra reaching half way to base of capsule *D. Guepinii*, 114.

Capsule cylindrical (1:5-6); teeth divided half way or entire.

[*D. obliquus* may be sought here.]Dioicous *D. plinthobius*, 112.Monoicous *D. Neo-Mexicanus*, 113.

b. *Not papillose* **D. Systilius**, 111.

2. *Costa vanishing at apex or forming a short point.*

Leaves hyaline $\frac{3}{4}$ of their length **D. obtusifolius**, 114.

Leaves hyaline only at base.

Margins revolute.

Capsule long cylindric, leaves crenulate **D. arenaceus**, 111.

Capsule elliptic, leaves entire **D. nervosus**,¹ 113.

Margins inflexed above **D. Garberi**, 112.

B. *Leaves with a pellucid border* **D. Porteri**, 112.

II. *Capsule nodding, or pendent, or arcuate.*

Leaves with a thickened border below.

Seta straight, capsule nodding or horizontal **D. cernuus**,² 114.

Seta reflexed, capsule pendent **D. Laureri**, 115.

Seta flexuous, capsule arcuate **D. camptothecius**, 121.

Leaves without a border **D. obliquus**, 115.

BARBULA, p. 115.

I. *Leaves with jointed dichotomous filaments on the costa.*

Costa broad ($\frac{1}{2}$ leaf), flattened, leaves thick, rigid §I. **Aloidella**, p. 192.

Costa narrow, round, leaves thin, broad §II. **Chloronota**, p. 192.

II. *Leaves not filamentose.*

Teeth from a low membrane, scarcely projecting from the mouth [excl. *B. brevipes*].

Plants small.

Leaf cells distinct §III. **Cuneifolia**, p. 192.

Leaf cells small.

Perichaetial leaves little different from the foliage.

§IV. **Unguiculata**, p. 193.

Perichaetial leaves long sheathing or convolute §V. **Convoluta**, p. 195.

Plants robust [excl. *B. caespitosa*].

Leaves entire; stems radiculose §VI. **Tortuosa**, p. 195.

Leaves serrate, stems not radiculose §VII. **Squarrosa**, p. 195.

Teeth from a high tessellated membrane §VIII. **Syntrichia**, p. 195.

¹*D. subtorquescens* C. M. & Kindb. (Appendix, no. 120) appears to be identical with *D. nervosus* B. & S.

²See Appendix, no. 122.

§I. Alcidellæ.

Synolcous *B. brevirostris*, 115.
 Dioicous.

Annulus broad, revoluble, lid $\frac{1}{2}$ to $\frac{3}{5}$ length of capsule.

Peristome twice twisted.

Leaves round-ovate, obtuse *B. macrorhyncha*, 125.

Leaves oblong, obtuse or apiculate *B. rigida*, 116.

Peristome straight or slightly twisted *B. aloides*, 162.

Annulus small, persistent, lid $\frac{1}{3}$ length of capsule *B. insignis*, 116.

§II. Chloronotæ.

Leaves with hair points.

Tip of leaf hyaline *B. membranifolia*, 116.

Tip of leaf concolorous.

Hair smooth, leaves acute or somewhat obtuse *B. chloronotos*, 116.

Hair serrate, leaves rounded obtuse *B. Henrieli*, 126.

Leaves without hair points *B. Mannii*, 127.

§III. Cuneifoliæ.

Leaves bordered by 2-4 rows of thickened cells *B. marginata*, 118.

Leaves bordered by 1 row of round yellowish cells with prominent papillæ, aristate *B. v. na*, 117.

Leaves with a broad yellowish border, not pointed *B. ?*, 128.

Leaves without a border.

Costa excurrent into a hoary hair *B. muralis*, 119.

Costa forming a short point or ceasing below apex.

Leaf cells smooth.

Margins plane.

Upper leaves opaque at the margins *B. cuneifolia*, 117.

Upper leaves pellucid at the margins *B. subcarnifolia*, 120.

Margins recurved above and below the middle at one side *B. carnifolia*, 129.

Leaf cells papillose [incl. *B. amplexa*?]

Peristome membrane long *B. brevipes*, 119.

Peristome membrane short.

Inner perichaetial leaves short *B. Bolanderi*, 118.

Inner perichaetial leaves long-sheathing, abruptly

reflexed *B. amplexa*, 118.

§IV. Ungulatae.

[*B. caespitosa* may be sought here.]

- I.
- Peristome wanting*
-
- B. rubiginosa*
- , 126.

II. *Peristome present.*A. *Teeth straight or scarcely twisted.*

Basal leaf cells rectangular.

- Teeth cancellate
- B. cancellata*
- , 122.

Teeth nodose, separate.

- Leaves long lanceolate, costa dark
- B. rigidula*
- , 123.

- Leaves ovate lanceolate, costa pale
- B. sparsidens*
- , 133.

- Basal leaf cells elongate-oval
- B. spadicea*
- , 139.

B. *Teeth plainly twisted.*1. *Leaves blunt or mucronate by the excurrent costa.*

Leaves short, ovate, the very apex obtuse.

- Capsule cylindric, calyptra reaching middle
- B. brachyphylla*
- ,
- ¹
- 123.

- Capsule ovate, calyptra reaching base
- B. purpurea*
- , 123.

Leaves longer, narrower, sharp pointed.

Cells at base rectangular and pellucid.

Teeth twisted 2-3 times, purple or red.

- Capsule oblong-elliptic to subcylindrical, sub-in-
-
- curved
- B. unguiculata*
- , 120.

- Capsule oblong, small, erect
- B. Jooriana*
- , 120.

- Teeth once loosely twisted, pale
- B. tortollifolia*
- , 144.

- Cells at the base quadrate, chlorophyllose
- B. Cruegeri*
- , 122.

2. *Leaves gradually pointed.*a. *Leaves not papillose* [incl. *B. artocarpa*?]

Leaf border plane.

- Annulus none, teeth reddish
- B. gracilis*
- , 127.

- Annulus large, simple, persistent, teeth whitish
- B. artocarpa*
- , 126.

Leaf border reflexed all around or revolute below.

Leaves long-subulate, costa percurrent or excurrent,

- red
- B. subgracilis*
- ,
- ²
- 136.

¹ See Appendix, no. 131.² See Appendix, no. 136.

Leaves long lanceolate, costa filling the acumen,
brown *B. pseudo-rigidula*, 140.

b. *Leaves papillose.*

i. *Cells at base roundish, quadrate, or short-rectangular.*

Annulus none or indistinct.

Leaves twisted when dry.

Costa 70μ wide at base and tapering gradually *B. fallax*, 121.

Costa 50μ wide, of equal breadth to middle *B. recurvifolia*, 122.

Leaves not twisted when dry.

Costa pale, percurrent *B. subicmadophila*, 137.

Costa brown, short excurrent *B. melanocarpa*, 138.

Annulus pale, compound *B. elata*, 125.

ii. *Cells at base rectangular, often elongated.*

[*B. fallax* may be sought here.]

Leaves erect-incurved, imbricate when dry.

Cells (above) $5-7\mu$ diameter *B. vinealis*, 124.

Cells twice as large *B. virescens*, 124.

Leaves squarrose-spreading or reflexed, twisted when dry.

Perichaetial leaves open, sheathing only at base, revolute

on edges *B. subfallax*, 121.

Perichaetial leaves half sheathing.

Annulus simple, narrow, persistent.

Leaves erect, half clasping at base *B. semitorta*, 126.

Leaves recurved, deeply concave *B. circinnatula*, 141.

Annulus double or triple.

Cells $5-7\mu$ in diameter { *B. cylindrica*, 125.

Cells twice as large { *B. flexifolia*, 124.

Cells twice as large *B. virescens*, 124.

BB. vinealis, flexifolia, virescens, cylindrica, with possibly *semitorta* and *circinnatula*, are doubtless forms of one species, so that the key will probably break down here.

BB. decursivula, Dieckii, horridifolia, robustifolia, and subcylindrica, belonging to this section, are described from sterile specimens and therefore cannot be included in the key, but descriptions will be found in the Appendix nos. 132, 134, 142, 143, and 145.

¹In L. & J. Man. 126, in note under *B. semitorta*, read "Comparable to *B. vinealis*" instead of *B. brachyphylla*. See Pacif. R. R. Rept. 4: 136.

on,
pseudo-rigidula, 140.

tri-rectangular.

B. fallax, 121.
B. recurvifolia, 122.

subcymadophila, 137.
B. melanocarpa, 138.
B. elata, 125.

elongated.

B. vinealis, 124.
B. virescens, 124.

dry.
involute
B. subfallax, 121.

B. semitorta,¹ 123.
B. circinnatula, 141.

{ B. cylindrica, 125.
B. flexifolia, 124.
B. virescens, 124.

with possibly semi-sterile specimens and species, so that the

folia, and subcylindrica, and subcylindrica-sterile specimens and species will be found in

comparable to *B. vinealis*"

§V. Convolutae.

Leaves involute on margin.

Aristulate by excurrent costa B. agraria, 123.
Acute or submucronate B. Donnellii, 128.

Leaves plane on margin or recurved.¹

Capsule costate when dry B. Rauli, 126.
Capsule smooth.

Leaves acute, costa percurrent B. convoluta, 127.
Leaves with hyaline point B. Closteri, 127.

Leaves obtuse.

Perichetial leaves rounded or truncate at apex B. chrysopoda, 143.
Perichetial leaves obtuse or short apiculate B. convoluta var. 147.

§VI. Tortuosae.

Leaf margin plane or undulate when moist.

Leaves long linear, acute, abruptly mucronate B. caespitosa, 129.
Leaves very long acuminate, cuspidate.

Twisted crispate when dry, above of one layer of cells B. tortuosa, 129.
Not crispate, brittle, two layers of cells above B. fragilis, 129.
Leaf margin involute, cucullate above B. inclinatula, 149.

§VII. Squarrosae.

Includes but one species B. squarrosa, 130.

§VIII. Syntrichiae.

[*B. brevipes* may be sought here.]

I. Leaves with a border of thickened cells.

Leaves with a distinct border, sometimes absent at the apex, plane.

Marginal cells elongated B. subulata,² 130.
Marginal cells roundish B. laevipila, 132.

Leaves with a narrow border at the base only.

Basal cells faintly chlorophyllose B. lato-exclisa, 160.
Basal cells strongly chlorophyllose B. papillinervis, 156.

¹ *B. platyneura* C. M. & Kindb., described from barren specimens falls here. See Appendix, no. 146.

² See Appendix, no. 150.

SCOULERIA, p. 137.¹*B. mucronifolia*, 131.

Leaves bordered by slightly larger, thick walled cells, peristome present *S. aquatica*,² 137.
 Leaves bordered by larger denser cells, peristome absent *S. marginata*, 164.

B. inermis, 131.
B. tubulata mutica, 130.

alpine

B. laevipila, 132.*B. alpina*, 151.

ulose

B. Muelleri, 133.*B. latifolia*, 132.*B. papillosa*, 133.

hair.

th in length.

B. ruralis,¹ 132.*B. leptotricha*, 159.*B. ruraliformis*, 153.*B. intermedia*, 157.

pale

B. laeviuscula, 155.*B. emarginata*, 161.

the

B. aciphylla, 158.

re-

B. brachyanga, 164.*B. brachyanga*, *laevi-**tricha* and *lato-ex-*

trms of the extremely

locarpa Kindb. is;

ishing them.

GRIMMIA, p. 134.

Capsule longer than the seta.

Straight, capsule symmetric §I. *Schistidium*.Crooked, capsule ventricose §II. *Gasterogrimmia*.

Capsule shorter than the seta.

Seta arcuate §III. *Eugrimmia*.Seta straight §IV. *Guembelia*.§I. *Schistidium*.

Leaves with hyaline points.

Capsule oblong *G. ambigua*, 135.

Capsule ovate-globose.

Leaves erect or appressed when dry.

In small, dense cushions, soft lurid green *G. conferta*,³ 135.

In lax cushions, coarse, fuscous.

Peristome distinct, orifice dilated *G. spherica*, 197.Peristome rudimentary, orifice not dilated *G. apocarpa*, 136.Leaves spreading when dry *G. heterophylla*, 168.

Leaves mucous.

Margin plané.

Entire or slightly serrate at the apex.

Ovate-oblong, short *G. atricha*, 169.Linear lanceolate *G. maritima*, 137.Coarsely dentate at the apex *G. Agassizii*, 136.Margins recurved or revolute { *G. conferta* } vars.⁴
 { *G. apocarpa* }

¹ *Grimmia* § *Scouleria*, L. & J. Man. 137. This genus has lately been revised by Mrs. Elisabeth G. Britton, Bull. Torr. Bot. Club 22: 38. 1896. *S. Muelleri* Kindb. and *S. aquatica catulliformis* Müll. are reduced to *S. aquatica* Hook. *S. Nevii* is referred to *S. aquatica nigrescens* Kindb.

² *Grimmia Scouleri* Müll. See Appendix, no. 163.

³ *G. chloroblasta* Kindb. (Appendix, no. 167) falls here. See also no. 166.

⁴ See Appendix, nos. 165, 166. *G. platyphylla* Mitt. (L. & J. Man. 136) = *G. apocarpa latifolia* Zett. sec. Ren. & Card. Rev. Bryol. 10: 83. 1892. A barren species, *G. pachyneurula* C. M. & Kindb., Mac. Cat. 65 (see Appendix, no. 170), evidently belongs to this section.

§II. *Gasterogrimmia*.

- Peristome wanting, lamina bistratose near apex . . . *G. anodon*, 138.
 Peristome present, lamina unistratose throughout . . . *G. plagipoda*, 138.

§III. *Eugrimmia*.

- Capsule costate when dry.
 Leaves homomalous-falcate when dry . . . *G. hamulosa*, 139.
 Leaves spirally twisted on stem when dry . . . *G. torquata*, 140.
 Leaves incurved-cirrhate when dry . . . *G. contorta*, 139.
 Leaves imbricate or slightly twisted when dry, hair pointed.
 Lamina of one layer of cells . . . *G. pachyphylla*, 181.
 Lamina 2-4-stratose in the upper part or at the margin.
 Hair point rough . . . *G. Muhlenbeckii*, 140.
 Hair point smooth.
 Annulus simple, monoicous . . . *G. pulvinata*, 138.
 Annulus compound (3-4), dioicous.
 Costa narrow, capsule 8-striate or costate.
 Lid short . . . *G. funalis*, 179.
 Lid long, straight . . . *G. trichophylla*, 141.
 Costa strong, capsule 10-striate . . . *G. elatior*, 172.
 Capsule not costate (or obscurely) when dry.
 Leaves falcate-reflexed when moist . . . *G. Watsoni*, 141.
 Leaves not reflexed.
 Margin plane, capsule elliptic, collum none . . . *G. Olneyi*, 142.
 [*G. Muhlenbeckii* may be sought here.]
 Margin reflexed.
 Leaves gemmiferous, capsule oval-oblong.
 Teeth papillose, leaf cells at base scarcely
 sinuous . . . *G. Philbertiana*, 171.
 Teeth smooth, leaf cells at base strongly
 sinuous . . . *G. Hartmanii*, 174.
 Leaves not gemmiferous.
 Open, erect, lanceolate . . . *G. California*, 142.
 Arcuate, ovate-lanceolate . . . *G. depilata*, 176.

Only incomplete descriptions of the following species can be obtained and they are therefore not included in the key: *G. aruatifolia* Kindb., *Arizonae* Ren. & Card., *prolifera* C. M. & Kindb., *tortifolia* Kindb., and *Hendersonii* Ren. & Card. See descriptions in Appendix, nos. 173, 176, 177, 178, 180.

§IV. *Guembelia*.

Lamina above 2-4-stratose.

Calyptra cucullate.

Leaves hair pointed.

Basal cells quadrate *G. tenerrima*, 188.Basal cells rectangular *G. commutata*, 145.

Leaves not hair-pointed, blunt or hyaline apiculate.

Teeth entire, annulus simple, indistinct *G. sulcata*, 186.Teeth irregularly lacerate or bifid, annulus compound, triple *G. unicolor*, 146.

Calyptra mitrate.

Leaf margins plane.

Hair point very rough, lid straight beaked *G. leucophæa*, 144.Hair point faintly denticulate, lid obliquely beaked *G. sarcocalyx*, 190.

Leaf margins recurved.

Walls of basal cells sinuate *G. Pennsylvanica*, 144.Walls of basal cells smooth *G. ovata*, 143.

Lamina having only the margin 2-4-stratose.

Leaves mucicous or hyaline, apiculate.

Costa reaching the hyaline apex, dioicous *G. elongata*, 185.Costa vanishing far below apex, monoicous *G. Coloradensis*, 143.

Leaves hair-pointed.

Annulus wanting.

Calyptra mitrate, covering whole capsule *G. calyptrata*, 144.Calyptra cucullate *G. montana*, 145.

Annulus present.

Cells of leaf base elongated (1:4 to 1:8) *G. Donriana*, 142.Cells of leaf base short (1:2), borders plane *G. alpestris*, 146.Cells of leaf base short, borders recurved *G. microtricha*, 187.Lamina of a single layer of cells *G. mollis*, 189.

The descriptions of *GG. cinclidodonteæ* C. Müll., *crassinervis* C. Müll., *tenella* C. Müll., and *Mannixæ* C. Müll. are incomplete and as no authentic material is at hand they are not included in the key. See Appendix, nos. 182, 183, 194, 184a.

RHACOMITRIUM, p. 147.

I. *Branches fastigiate*.Leaves with a short hyaline point *R. Sudeticum*, 149.

Leaves mucicous.

Costa with 2-4 lamellæ at back *R. patens*, 147.

Costa not lamellose.

Leaves with quadrate cells at the base, decurrent and semi-auricled *R. depressum*, 148.

Leaves with linear cells at the base, neither decurrent nor auricled.

Obtuse.

Perichætal leaves costate, seta long *R. aciculare*, 148.

Perichætal leaves ecostate, seta short *R. Nevil*, 148.

Acute.

Capsule oblong, not striate, teeth orange *R. Macounii*,¹ 193.

Capsule oval, striate or plicate when dry, teeth purple *R. robustifolium*, 195.

II. *Branches fasciculate.*

A. *Leaves muticous.*

Cells elongated above *R. fasciculare*, 150.

Cells quadratic above.

Costa percurrent.

Leaf cells smooth *R. varium*,² 150.

Leaf cells rough, simple papillæ over the lumen *R. canescens*, 151.

Costa vanishing below apex, cells with geminate papillæ over the partitions *R. protensum*, 192.

B. *Leaves with a hyaline point.*

Cells linear at least above.

Hyaline point not papillose *R. microcarpum*,³ 150.

Hyaline point strongly erose-serrate and papillose *R. lanuginosum*,⁴ 151.

Cells quadratic above.

[*R. micropus* may be sought here.]

Strongly papillose on both sides *R. canescens*,⁵ 151.

Smooth or nearly so.

Annulus broad.

Alar cells of leaf linear *R. heterostichum*,⁶ 149.

¹ *R. alternatum* C. M. & Kindsb., described from barren specimens, seems to be referable to this species. See Appendix, no. 194.

² *R. Oregonum* Ren. & Card., Bot. Gaz. 13: 198. pl. 15. 1888, is this species (sde J. Cardot in litt.) which seems to be *R. canescens lutescens* L. & J.; sde Mrs. E. G. Britton in litt.

³ See Appendix, no. 200.

⁴ See Appendix, no. 202.

⁵ See Appendix, nos. 203, 204.

⁶ See Appendix, nos. 196-198.

nt and
 . *R. depressum*, 148.
 urrent

. *R. aciculare*, 148.
 . *R. Nevii*, 148.
R. Macounii,¹ 193.
 teeth
R. robustifolium, 195.

R. fasciculare, 150.

R. varium,² 150.
R. canescens, 151.
 e pa-
R. protensum, 192.

it.
R. microcarpum,² 150.
R. lanuginosum,² 151.

R. canescens,² 151.

R. heterostichum,² 149.

mens, seems to be refera-

this species (*vide* J. Car-
de Mrs. E. G. Britton *in*

Alar cells of leaf quadrate *R. micropus*, 199.
 Annulus none *R. speciosum*, 201.

COSCINODON, p. 154.

Costa not entering the hyaline point, which is less than
 the leaf in length.
 Dioicous, leaves oblong lanceolate *C. pulvinatus*, 154.
 Autoicous, leaves obovate *C. Rauli*,¹ 155.
 Costa forming a rough hyaline point twice as long as the
 leaf *C. Wrightii*, 155.

PTYCHOMITRIUM, p. 156.

Plants large (3 cm. +), leaves acuminate, sharply dentate *P. Gardneri*, 156.
 Plants small (1 cm. —), leaves not acuminate, nearly or
 quite entire.
 Collum none.
 Teeth subulate (1:10), entire *P. incurvum*, 157.
 Teeth lanceolate (1:4), bi- or trifid *P. Drummondii*, 157.
 Collum equalling one-third sporangium *P. pygmaeum*, 157.

AMPHORIDIUM, p. 158.

Leaf margins plane, entire *A. Lapponicum*, 158.
 Leaf margins recurved or revolute.
 Leaves remote, recurved-spreading, serrate *A. Sullivantii*, 159.
 Leaves close.
 Costa excurrent, seta arcuate *A. Californicum*, 159.
 Costa vanishing below apex.
 Entire *A. Mougeotii*, 159.
 Serrulate *A. caespitosum*, 160.

ZYGODON.²

Capsule without a peristome *Z. viridissimus*, 207.
 Capsule with double peristome *Z. conoideus*, 208.

¹ Including *C. Renauldi* Card., Bot. Gaz. 15: 41, 1890.

² See appendix, no. 206.

ULOTA,¹ p. 160.I. *Leaves rigid, not crispate when dry.*[*U. Drummondii* may be sought here.]

- Costa percurrent *U. Hutchinsii*,² 163.
 Costa ceasing below apex *U. Barclayi*, 164.

II. *Leaves crispate when dry.*

- Capsule not constricted below the mouth when dry.
 Costate only at the mouth *U. Ludwigii*, 161.
 Costate for its whole length.
 Stems creeping, leaves slightly crispate, cilia 0 *U. Drummondii*, 161.
 Stems not creeping, leaves strongly crispate, cilia present.
 Capsule short oval, neck short.
 Upper leaves tipped with gemmæ *U. phyllantha*, 163.
 Upper leaves not tipped with gemmæ *U. Bruchii*,³ 162.
 Capsule cylindric, neck long *U. maritima*,⁴ 217.
 Capsule constricted below the mouth when dry.
 Cells uniform throughout the leaf *U. megalospora*,⁵ 210.
 Cells different at the angles.
 Teeth lacunose at the apex *U. curvifolia*,⁶ 161.
 Teeth not lacunose.
 Neck very long, teeth confluent *U. crispa*,⁷ 162.
 Neck shorter, teeth separated at apex *U. crispa*, 163.

¹ It seems best, while retaining for convenience the generic name *Ulota* (without expressing any opinion as to its validity), to utilize Mrs. Britton's careful study of our American species. The nomenclature used is also derived in part from her paper in Bull. Torr. Bot. Club 21: 65-76, 1894.

² See Appendix, no. 212.

³ From the description given (Mac. Cat. 82) I am unable to separate *U. obtusiuscula* C. M. & Kindb.

⁴ Renaud and Cardot think that *U. maritima* and *U. phyllantha* resemble each other so closely that *U. maritima* should not stand as a distinct species.

⁵ Including *U. subulata* and *U. subulifolia* C. M. & Kindb., Mac. Cat. 82, sec. Elis. G. Britton.

⁶ Including *U. Americana* Mitt., Man. 162, and probably *U. scabrata* Kindb., Mac. Cat. 83. (E. G. B.)

⁷ Including *U. camplopoda* and *U. connectens* Kindb., Mac. Cat. 85. (E. G. B.)

ORTHOTRICHUM, p. 164.

1. *Stomata superficial.*1. *Peristome simple.*

Capsule entirely smooth.

Wholly exerted *O. lævigatum*, 165.Immersed *O. Schlotthaueri*, 213.

Capsule strongly costate.

Leaves densely papillose.

Capsule wholly exerted, abrupt at base, teeth 8 *O. Douglasii*, 167.

Capsule immersed or emergent, defluent into seta, teeth 16.

Leaf bistratose in the upper part *O. Sturmii*, 166.Leaf unistratose *O. rhabdophorum*, 227.Leaves almost smooth *O. bullatum*, 222.

Capsule ribbed only near the mouth.

Teeth striolate, capsule exerted *O. Roellii*, 217.Teeth papillose, capsule immersed *O. Shawii*, 213.2. *Peristome double.*[*O. lævigatum*, *Sturmii*, *Roellii* and *Schlotthaueri* may be sought here.]a. *Capsule entirely smooth.*Immersed, papillæ simple *O. striatum*,¹ 174.

Exserted, papillæ bifurcate.

Alar cells thick walled, quadrate to hexagonal *O. arcticum*, 223.

Alar cells not thickened, rectangular.

Leaf margin revolute, capsule cylindrical when

moist *O. Macounii*, 223a.Leaf margin reflexed, capsule obovate when moist *O. Kingianum*, 170.b. *Capsule strongly costate.* (c)Leaves beset with clavate gemmæ, teeth reflexed, cilia 16 *O. Lyellii*,² 177.

Leaves not gemmiferous.

Teeth erect when dry, cilia 16 *O. Texanum*, 166.

Teeth reflexed when dry, cilia 8.

Cilia of two rows of cells, broad.

Leaf margins plane, cilia broad throughout *O. obtusifolium*, 177.Leaf margins revolute, cilia broad at base *O. sordidum*, 170.¹ *O. tetocarpum* B. & S.² See Appendix, no. 221.

Cilia of a single row of cells or double at the base.

Capsule immersed or emergent.

Teeth papillose, not vermicular striate *O. affine*, 168.

Teeth vermicular striate, not papillose *O. fastigiatum*, 214.

Capsule exerted.

Teeth not papillose *O. Blyttii*, 229.

Teeth minutely papillose *O. praemorsum*, 226.

c. *Capsule ribbed only near the mouth.*

Teeth erect when dry.

Papillae of leaves simple, minute *O. rupestre*, 167.

Papillae of leaves bifurcate, salient *O. Killiasii*, 225.

Teeth reflexed when dry.

Upper part of leaf bistratose, teeth transversely lineolate *O. Bolanderi*, 167.

Upper part of leaf unistratose, teeth very papillose.

Teeth when dry reflexed and applied to capsule *O. elegans*, 224.

Teeth when dry touching capsule only with tip *O. speciosum*,¹ 169.

Three little known and unimportant species from Greenland (Mac. Cat. 88): *OO. Breutelii* Hpe., *Barthii* Sendtn., and *Groenlandicum* Bergg., belonging to the division with superficial stomata are not included in the key. According to Venturi in Husn. Musc. Gall. 175, the first two are only forms of *O. Blyttii*.

II. *Stomata immersed.*

1. *Peristome simple, capsule costate.*

Leaves obtuse, capsule immersed or emergent *O. Jamesianum*, 177.

Leaves acute or acuminate.

Capsule long exerted, teeth erect when dry *O. anomalum*, 164.

Capsule half-emergent, teeth spreading when dry *O. cupulatum*, 165.

[*O. Hallii* may be sought here.]

2. *Peristome double.*

a. *Capsule smooth when dry.*

Cilia wider than teeth *O. exiguum*, 174.

Cilia narrow.

Capsule gradually narrowed to seta *O. pallens*, 175.

Capsule abruptly contracted to seta *O. pusillum* Mitt.² 173.

¹ See Appendix, no. 223.

² *O. psilocarpum* James.

b. *Capsule costate when dry.** *Leaves hyaline pointed.*

- Cilia of a single row of cells, teeth equidistant . . . *O. diaphanum*, 176.
 Cilia of two rows of cells, teeth bigeminate . . . *O. canum*, 176.

** *Leaves obtuse. (***)*

[*O. tenellum*, *pumilum*, *pallens* and *strangulatum* may also be sought here.]

- Capsule exserted . . . *O. cylindrocarpum*, 173.
 Capsule immersed.

Abruptly contracted to the seta, collum not evident.

- Teeth papillose . . . *O. eurphyllum*, 219.
 Teeth vermicular striate . . . *O. Hallii*, 170.

Gradually narrowed to seta with evident collum.

- Teeth 8, bigeminate, reflexed when dry.
 Punctulate, cilia of 2 rows of cells . . . *O. Ohioense*, 170.
 Papillose, cilia double at the base only.
 Cilia 8, shorter than the teeth . . . *O. Sprucei*, 215.
 Cilia 16, eight as long, eight shorter than the teeth *O. rivulare*, 176.
 Teeth 16, separate, erect when dry . . . *O. nudum*, 220.

*** *Leaves acute.*

Capsule exserted.

Abruptly contracted to seta, collum not evident.

- Cilia 16, single series of cells . . . *O. ulotaforme*, 225.
 Cilia 8, double row of cells . . . *O. consimile*, 173.

Gradually narrowed to the seta with evident collum,

- cilia 16, appendiculate . . . *O. pulchellum*,¹ 175.

Capsule immersed or nearly.

Leaves with salient furcate papillae.

- Teeth 16, separate . . . *O. urnigerum*, 216.
 Teeth 8, bigeminate.

Cilia of a double row of cells.

- Teeth papillose throughout, split along the median
 line . . . *O. pumilum Americanum*, 220.

- Teeth punctate, yellowish white . . . *O. Watsoni*, 168.

Teeth finely papillose below, paler above with long-

- itudinal sinuous lines, somewhat perforate *O. alpestre*, 168.

- Cilia double at the base, single above . . . *O. Schimperii*,² 171.

¹ See Appendix, nos. 233, 234.

² *O. fallax* Schimp.

Leaves with simple often weak papillae.

[*O. Schimperii* and *pumilum* may be sought here.]

Cilia of a single row of cells *O. Hendersonii*, 232.

Cilia of a double row of cells.

Teeth granulose, not papillose, capsule obovate, contracted below the mouth when dry *O. strangulatum*, 172.

Teeth papillose, capsule sub-cylindric, little contracted below mouth when dry.

Calyptra hairy, teeth pale brown *O. tenellum*, 172.
Calyptra naked, teeth reddish *O. Rogeri*, 231.

"Notes on North American Species of Orthotrichum" by E. G. Britton, Bull. Torr. Bot. Club 20: 393. 1893 and 21: 1, 137. 1894, have been followed in many points. *O. psilothecium* C. M. & Kind. is reduced to *O. strangulatum*; *O. Pringlei* C. Müll. to *O. Lyellii*; *O. brachytrichum* Schimp. to *O. Schimperii*; *O. stenocarpum* Vent. to *O. Macountii* Aust. *O. stellatum* Brid. and *O. anomalum Americanum* are rejected as doubtful.

MACROMITRIUM, p. 178.

Capsule plicate at mouth and base only *M. Sullivanii*, 178.
Capsule costate its whole length.
Lid conic, blunt, peristome wanting *M. Fitzgeraldi*, 178.
Lid subulate, peristome present *M. rhabdocarpum*, 179.
Capsule smooth *M. mucronifolium*, 179.

ENCALYPTA, p. 180.

I. Capsule spirally striate and sulcate when dry.

Capsule twisted to the right when dry, leaves with hyaline hair points, teeth glabrous *E. Selwynii*, 183.
Capsule twisted to the left when dry.
Leaves cucullate *E. cucullata*, 233.
Leaves not cucullate.
Leaves acute or apiculate, teeth papillose, median line *E. prosera*, 182.
Leaves mucicous, usually obtuse, teeth filiform, nodose, minutely papillose *E. streptocarpa*, 183.

II. *Capsule vertically striate and sulcate when dry, or smooth.*

Distinctly striate.

Leaves plane or slightly concave.

Oblong or lanceolate above, calyptra scabrous *E. rhabdocarpa*, 181.Short, often subspatulate, calyptra not scabrous *E. leiomitra*, 237.Leaves revolute all around, narrow *E. Alaskana*, 240.

Smooth or faintly striate.

Calyptra entire or lacerate at the base.

Smooth at the apex *E. commutata*, 180.

Papillose or spinose at the apex.

Costa long excurrent *E. subspathulata*, 236.

Costa percurrent or vanishing.

Capsule minutely wrinkled lengthwise *E. vulgaris*, 181.Capsule smooth when dry *E. lelocarpa*, 239.

Calyptra fringed at base, peristome present.

Leaves spiculate-acuminate *E. ciliata*, 183.Leaves mucous *E. Macounii*, 182.*E. apophysata* N. & H., *sec.* Schimper Synop. Musc. Europ. 345 [ed. 2] collected by Drummond is probably an error. See Appendix, no. 241.

CALYMPERES, p. 184.

Leaves oblong or broad-ovate.

Upper leaves very obtuse, often filamentose at apex *C. Richardi*, 184.Upper leaves acute, often filamentose in middle *C. disciforme*, 184.Leaves narrowly panduriform, obtuse or retuse *C. (?) crispum*, 184.

SYRRHOPODON, p. 185.

Leaf margins bilamellate upwards *S. Floridanus*, 185.Leaf margins single throughout *S. Texanus*, 185.

TETRAPHIS, p. 186.

Pedicel straight *T. pellucida*, 186.Pedicel geniculate at middle *T. geniculata*, 187.

DISSODON, p. 189.

Seta short (5 mm.), thick, capsule erect, chestnut

brown *D. Hornschuchii*, 189.

- Seta longer (1.5 cm.), plants 1-2 cm. high, capsule often inclined, orange **D. Froelichianus**, 190.
 Seta longer (3-4 cm.), plants 4-12 cm., capsule erect, orange **D. splachnoides**, 190.

TAYLORIA, p. 190.

- Teeth cleft along the middle line or split to the base.
 Columella long exserted, lid long conic **T. splachnoides**, 191.
 Columella scarcely exserted, lid suddenly obliquely beaked **T. acuminata**, 245.
 Teeth neither cleft nor split but sometimes lacunose.
 Columella scarcely exserted, rhizoids with gemmae **T. serrata**, 191.
 Columella long exserted, rhizoids without gemmae **T. tenuis**, 191.

TETRAPLONDON, p. 191.

- Leaves sharply serrate, narrowed to filiform point **T. angustatus**, 192.
 Leaves distantly incised-serrate, gradually acuminate **T. australis**, 192.
 Leaves entire, more or less abruptly filiform-apiculate.
 Costa sub-excurrent, empty sporangium constricted in middle **T. mnioides**, 192.
 Costa ceasing below point, empty sporangium not constricted in middle **T. urceolatus**, 193.

SPLACHNUM, p. 193.

- Apophysis ovate or subglobose.
 About the size of the sporangium.
 Costa excurrent, apophysis red **S. sphaericum**, 194.
 Costa ceasing below apex, apophysis at first green then brown **S. Wormskoldii**, 194.
 Greatly exceeding the sporangium **S. vasculosum**, 194.
 Apophysis pyriform, exceeding the sporangium **S. ampullaceum**, 194.
 Apophysis campanulate.
 Purple **S. rubrum**, 195.
 Yellow **S. luteum**, 195.

¹ See also Appendix, no. 254.

PHYSCOMITRIUM, p. 196.

Capsule or seta immersed.

[*P. Hookeri* may be sought here.]

- Capsule subglobose *P. immersum*, 196.
 Capsule pyriform when fresh *P. Coloradense*, 250.

Capsule and seta exserted.

Leaves entire or nearly so.

- Seta short, but little exceeding leaves *P. Hookeri*, 198.
 Seta much longer (5-20 mm.).
 Leaves very acute, bordered *P. acuminatum*, 198.
 Leaves oblong-lanceolate, more obtuse, not bordered *P. Californicum*, 253.

Leaves serrate at least above the middle.

Mouth of capsule bordered by 4-8 rows of cells.

Seta straight.

- Leaves oblong acuminate, annulus double *P. pygmaeum*, 197.
 Leaves lanceolate, annulus apparently single *P. Drummondii*, 251.
 Leaves ovate-acuminate, annulus apparently

single *P. Kellermanii*, 249.Seta curved *P. australe*, 252.

Mouth of capsule bordered by 8-12 rows of cells.

- Capsule turbinate, 1-2 mm., mouth flaring *P. turbinatum*,¹ 198.
 Capsule pyriform, 2-3 mm. mouth not flaring *P. megalocarpum*, 248.

The revision of the genus by Mrs. E. G. Britton, Bull. Torr. Bot. Club 21: 189. 1894, has been followed. *P. pyriforme* Brid. is a European species and does not occur in America; American specimens called *pyriforme* are referable to *P. turbinatum* Müll. *P. strangulatum* Kindb., Ott. Nat. 4: 62, is reduced to *P. turbinatum*. *P. platyphyllum* Kindb., Mac. Cat. 269, is rejected, as the species is founded upon such very immature specimens that it probably can not stand.

ENTOSTHODON, p. 199.

Leaves acute, capsule short-pyriform.

- Costa percurrent, teeth dark red, striolate *E. Drummondii*, 199.

Leaves acuminate, capsule long-pyriform.

- Costa reaching middle, teeth whitish, granulose *E. Bolanderi*, 199.
 Costa subpercurrent, teeth red, nodose, papillose *E. Templetoni*, 200.

¹ See also Appendix, nos. 245-247.

. Froelichianus, 190.

D. splachnoides, 190.

T. splachnoides, 191.

T. acuminata, 243.

T. serrata, 191.

T. teaulis, 191.

T. angustatus, 192.

T. australis, 192.

T. muioides, 192.

T. urceolatus, 193.

S. sphaericum, 194.

S. Wormskioeldii, 194.

S. vasculosum, 194.

S. ampullaceum, 194.

S. rubrum, 195.

S. luteum,¹ 195.

FUNARIA, p. 200.

Annulus wanting.

Leaves entire or nearly.

Capsule arcuate, leaves acuminate.

Costa excurrent *F. Americana*, 201.Costa vanishing *F. Mediterranea*, 201.Capsule erect, leaves acute *F. Californica*, 201.

Leaves sharply serrate.

Short-pointed, lid convex, mamillate *F. serrata*,¹ 201.Long acuminate, lid short conic *F. calcarea*, 201.

Annulus large, revoluble.

Capsule irregularly plicate and furrowed.

Leaves with involute margins *F. convoluta*, 202.Leaves with plane margins *F. flavicans*, 202.

Capsule distinctly striate-costate.

Leaves short-acuminate, lid large, spores 12-17 μ *F. hygrometrica*, 202.Leaves long-acuminate, lid small, spores 24-28 μ *F. microstoma*, 203.

BARTRAMIA, p. 203.

Capsule erect, peristome simple or none.

Leaves lance-subulate, ovate at the base.

Margin reflexed, capsule rugose when dry *B. Menziesii*, 204.

Margin plane, capsule furrowed when dry.

Capsule exserted, basal cells alike *B. subulata*, 204.Capsule immersed, basal cells shorter towards
the margin *B. breviseta*, 256.Leaves linear, gradually tapering to subulate apex,
capsule ribbed *B. stricta*, 205.

Capsule curved, lid oblique, peristome double.

Seta short (= capsule), fruit pseudo-lateral *B. Halleriana*, 206.

Seta exceeding stems.

Leaves smooth *B. Ederiana*,² 205.Leaves papillose only on upper surface *B. radicalis*, 206.

Leaves papillose on both surfaces.

Abruptly narrowed and bent above the hyaline
base *B. ithyphylla*, 205.

Gradually tapering to apex, not hyaline at base.

Leaves straight, autoicous.

Margined, borders revolute *B. pomiformis*, 206.¹ See also Appendix, no. 255.² *B. Ederiana minor* Kindb., Mac. Cat. 106, is *nomen nudum*.

- Not margined, sheathing *B. glauco-viridis*, 257.
 Leaves circinate, long sheathing *B. circinnulata*, 258.

PHILONOTIS, p. 208.

Leaves plicate lengthwise.

- Leaf cells linear, costa excurrent *P. Mohriana*, 210.
 Leaf cells rectangular or oval, costa percurrent or
 vanishing *P. seriata*, 262.

Leaves not plicate lengthwise.

- Leaf cells quadrate, slightly papillose *P. Macounii*, 208.
 Leaf cells oblong hexagonal, slightly papillose *P. glabriuscula*, 263.
 Leaf cells rectangular to linear.

- Cilia two, rudimentary *P. Muhlenbeckii*, 208.
 Cilia two, half as long as or equaling segments.

- Mouth of capsule with 8 rows of transversely
 elongated cells *P. fontana*,¹ 209.
 Mouth of capsule with 4 rows of transversely
 elongated cells *P. calcarea*, 209.

MEESIA, p. 212.

Leaves entire, margins reflexed or revolute.

- Synicous, costa very thick ($\frac{1}{2}$ leaf base) *M. uliginosa*, 212.
 Autoicous, costa narrow ($\frac{1}{2}$ leaf base) *M. Albertinii*, 213.
 Leaves entire, margins plane *M. longiseta*, 212.
 Leaves serrate *M. tristicha*, 213.

MIELICHHOFERIA, p. 214.

- Leaves lanceolate, sharply serrate at apex *M. nitida*, 214.
 Leaves broad ovate, entire, or slightly crenulate above *M. cuspidifera*, 264.

WEBERA, p. 215.

[*Bryum Froudei* and *B. angustirete* will be sought here.]

- I. *Leaves with a reddish border, distinct to apex* *W. Tozeri*, 222.

II. *Leaves not bordered, or indistinctly.*

¹ See Appendix, nos. 259-261.

A. *Annulus present.* (B on p. 213.)

1. *Segments and cilia of endostome imperfect, often only a lacinate membrane* *W. camptotrachela*, 278.

2. *Segments of endostome not widely open along the keel, cilia none or short (excl. W. longicolla).* (3)

Inflorescence autoicous *W. acuminata*, 216.

Inflorescence synoicous or dioicous.

Costa very broad, $\frac{1}{8}$ - $\frac{1}{4}$ of leaf base *W. Cardoti*, 265.

Costa narrow.

Plants less than 1 cm., seta 5-8 mm., capsule wide mouthed when dry *W. nudicaulis*, 220.

Plants small, seta longer, mouth of capsule constricted when dry *W. Bolanderi*, 220.

Inflorescence paroicous.

[*W. nudicaulis* may be sought here.]

Neck shorter than sporangium, cilia none *W. polymorpha*, 216.

Neck equaling sporangium, cilia more or less developed.

Tufts low, 1 mm.-2 cm. high *W. elongata*, 216.

Tufts higher, 2-5 cm. *W. longicolla*, 217.

3. *Segments of endostome split and gaping along keel, cilia well developed.*

Inflorescence paroicous or polygamous.

Capsule pendent, touching seta, not contracted under mouth *W. cucullata*, 218.

Capsule horizontal or pendent, not touching seta, contracted below mouth.

Costa excurrent, comal leaves revolute on the borders *W. canaliculata*, 271.

Costa vanishing below apex, comal leaves plane on the borders.

Tufts 1-2 cm. high, basal membrane of the endostome $\frac{1}{8}$ to $\frac{1}{2}$ height of teeth *W. nutans*,¹ 217.

Tufts 2-4 cm. high, basal membrane of the endostome $\frac{1}{4}$ height of teeth *W. cruda*,² 218.

¹ See Appendix, nos. 269, 270.

² See Appendix, no. 267.

Inflorescence dioicous.

Leaves with plane or slightly curved borders.

Antheridia in the axils of the perigonal leaves.

Upper leaves lance linear (1: 3-50) seta 3-4 cm. *W. sphagnicola*, 219.Upper leaves lanceolate (1: 4-5) seta 1-2 cm. *W. gracilis*, 275.Antheridia in a terminal cluster . . . *W. Lessuriana*, 221.

Leaves with mostly revolute borders.

Costa vanishing below apex.

Leaves of sterile shoots ovate, obtuse; comal leaves

broad pointed, margin entire . . . *W. Ludwigii*, 274.

Leaves narrow, linear lanceolate, sharp pointed,

plainly toothed near the apex.

Seta twisted to right, membrane of the inner peristome

 $\frac{1}{2}$ length of teeth, capsule oval, short necked.Annulus revoluble, costa red . . . *W. Columbica*, 279.Annulus remaining attached to lid, costa green *W. pulchella*, 222.Seta twisted to left, membrane $\frac{1}{4}$ length of teeth,capsule pyriform, longer necked . . . *W. commutata*, 220.

Costa reaching the apex.

Capsule oval, long necked, leaves serrate at apex *W. annotina*, 219.

Capsule obovate, short-necked, leaves denticulate

above . . . *W. pycnoderrens*, 277.B. *Annulus none.*Leaves nearly entire, cilia very short . . . *W. Drummondii*, 219.Leaves nearly entire, cilia 3 . . . *W. Bigelovii*, 223.

Leaves sharply serrate.

Stem rec., leaves glaucous-green . . . *W. albicans*,¹ 222.Stem and leaves green . . . *W. carnea*, 221.[*W. pulchella* may be sought here.]

A number of species have been described from sterile and immature specimens, whence it is impossible to determine their place in the key. They are *W. longibracteata* Broth., 268; *W. microapiculata* C. M. & Kindb., 281; *W. polymorphoides* Kindb., 266; *W. microcaulon* C. M. & Kindb., 272; *W. subcucullata* C. M. & Kindb., 273; *W. microdenticulata* C. M. & Kindb., 276. The first two are sterile, the remaining four are immature. The numbers attached refer to corresponding descriptions in the Appendix.

¹ See Appendix, no. 220.

BRYUM, p. 223.

Upper leaf cells rhombic to hexagonal (sublinear in *BB. Froudei* and *angustirete*).

Plants not from stolons.

Cilia none, or inappendiculate §I. *Cladodium*.

Cilia 2-4, appendiculate §II. *Eubryum*.

Plants from stolons §III. *Rhodobryum*.

Upper leaf cells linear (1:10-15), branches julaceous §IV. *Anomobryum*.

§I. *Cladodium*.A. *Autoicous*.

Leaves broad (1:2) costa vanishing *B. calophyllum*, 227.

Leaves ovate-lanceolate or long acuminate.

Cilia 2, long, smooth *B. Brownii*, 224.

Cilia 3, short *B. mamillatum*, 228.

Cilia none, or rudimentary.

Capsule symmetric, pyriform, collum $\frac{1}{2}$ sporangium.

Leaves faintly bordered, serrate above, slightly revolute *B. Warneum*, 226.

Leaves faintly bordered, margin entire, distinctly revolute *B. Edwardsianum*, 231.

Leaves very distinctly bordered, broadly revolute *B. Biddlecomiae*, 226.

Capsule usually unsymmetric, elongate, collum = sporangium *B. uliginosum*, 227.

B. *Synoicous, or heteroicous*. (C)

Costa long excurrent.

Endostome attached to peristome.

Spores verruculose *B. arcticum*, 224.

Spores smooth, about 30μ *B. pendulum*, 225.

Spores smooth, 20-25 μ .

Teeth orange red *B. angustirete*, 224.

Teeth pale *B. Roellii*, 225.

Endostome free, or slightly attached.

Seta 3-4 cm. long, capsule 1:2.5-3.

Upper leaf cells long hexagonal *B. inclinatum*,¹ 225.

¹ *B. stenotrichum* C. Müll. will be sought here; and I am unable from the description alone to discover any essential difference between it and *B. inclinatum*. See Appendix, no. 280.

- Upper leaf cells sublinear **B. Froudei**, 282.
 Seta 1 cm. long, capsule 1.2 : 2 **B. Archangelicum**, 287.
 Costa short excurrent or percurrent.
 Leaves not bordered **B. Knowltoni**, 292.
 Leaves bordered.
 Decurrent, seta 1 cm. long **B. brachyneuron**, 286.
 Decurrent, seta 2-3 cm. long **B. Labradorense**, 289.
 Not decurrent.
 Costa excurrent, leaves reddish, margin scarcely
 revolute **B. purpurascens**, 224.
 Costa vanishing or barely excurrent, margin
 strongly revolute **B. lacustre**, 226.

[*B. flexuosum* may be sought here.]

C. *Dioicous*.

- Collum smooth or rugose when dry.
 Endostome adherent to peristome, cilia none **B. flexuosum**, 227.
 Endostome free, cilia present.
 Costa percurrent, teeth bright purple **B. Californicum**, 237.
 Costa short excurrent, teeth orange **B. subpurpurascens**, 283.
 Collum longitudinally sulcate when dry.
 Leaf cells pitted, perichætical leaves costate, **B. cœneum**, 294.
 Leaf cells not pitted, inner perichætical leaves ecostate **B. fallax**, 293.

§II. Eubryum.

A. *Synœious*.

- Costa not excurrent.
 Leaf margin serrate above, apex straight **B. Oregonum**, 230.
 Leaf margin entire, apex recurved **B. utridulum**, 305.
 Costa excurrent into a smooth point.
 Margins recurved **B. torquescens**,¹ 230.
 Margins plane **B. microstegium**, 302.
 Costa excurrent into a serrate point.
 Leaves not decurrent, long cuspidate.
 Not bordered, entire **B. intermedium**,² 228.
 Not bordered, serrate or denticulate above **B. synœico-cœspiticium**, 316.
 Bordered **B. cirrhatum**,³ 228.

¹ See Appendix, no. 344. On account of the imperfect description I cannot place *B. sanguilentum* in the Key.

² See Appendix, no. 295.

³ See Appendix, no. 296.

Leaves decurrent.

With a broad border.

Leaves short-pointed *B. bimum*,¹ 229.Leaves long-cuspidate *B. cuspidatum*, 297.Without a border *B. lonchocaulon*, 229.*B. Polygamous.* (C)Leaves long cuspidate by excurrent costa, not bordered *B. provinciale*, 230.*C. Autoicous.* (D)

Leaves bordered, costa percurrent or excurrent.

Border wide, 5-6 rows of cells, distinctly revolute *B. pallescens*,² 231.Border narrow, faintly revolute *B. anæctanglaceum*, 323.

Leaves not bordered.

Costa excurrent *B. subrotundum*, 231.Costa vanishing below the apex *B. teres*, 306.*D. Dioicous.*1. *Costa not excurrent, or when excurrent forming a short point only*
(2 on p. 218.)a. *Leaves obtuse.*Distant, broadly ovate or oblong, rounded *B. cyclophyllum*, 237.

Thickly clothing stem, generally imbricate, narrower.

Not bordered.

Leaves strongly decurrent *B. obtusifolium*, 325.Leaves not decurrent *B. capitellatum*, 313.

Bordered, sometimes indistinctly.

Dull olive-green, margins strongly revolute.

Membrane of endostome $\frac{1}{2}$ length of peristome *B. Muhlenbeckii*, 233.Membrane of endostome low *B. rubicundulum*, 314.

Yellowish or dull green or purplish.

Costa vanishing below apex, tips of branches crimsoned.

Cells polygonal, thick walled *B. miniatum*, 233.Cells rhombic, sub-quadrate below *B. Atwateriae*, 234.Costa percurrent, red, branch tips green *B. percurrentinerve*, 312.¹ See Appendix, nos. 298, 299.² See Appendix, nos. 303, 304.

b. *Leaves pointed, costa percurrent or excurrent. (c)*i. *Capsule short (1:2), abrupt at base.*

Capsule blood-red or dark purple, teeth red at base *B. atropurpureum*, 232.
 Capsule pale, teeth pale throughout . . . *B. microglobum*, 307.

ii. *Capsule longer (1:3+), tapering at base.** *Blood red to dark purple.*

Plants short (5-15 mm.) in small lax, yellowish-green tufts.

Collum half length of sporangium . . . *B. erythrocarpum*, 232.
 Collum equaling or exceeding the sporangium in
 length . . . *B. micro-erythrocarpum*, 308.

Plants larger (3-5 cm.), in large tufts.

Tufts shining red or purplish . . . *B. alpinum*, 233.
 Tufts greenish, costa yellow, percurrent or excur-
 rent . . . *B. haematocarpum*, 311.

* * *Yellowish-brown.*

Slightly incurved.

Constricted below mouth . . . *B. meesloides*, 336.
 Not constricted . . . *B. pallens*, 237

Symmetric.

Strongly constricted below mouth.

Stems about 1 cm. high . . . *B. turbinatum*, 238.
 Stems 4-10 cm. high . . . *B. Schleicheri*, 239.

Slightly constricted below mouth.

Leaves long decurrent . . . *B. extenuatum*, 323.
 Leaves not decurrent or only slightly.

Leaf margin plane.

Pale on the borders . . . *B. Sawyeri*, 322.
 Red on the borders . . . *B. ernbescens*, 324.

Leaf margins revolute.

Margin serrate at the apex . . . *B. pseudotriquetrum*,¹ 238.
 Margin quite entire.

Cilia single . . . *B. acutiusculum*, 321.
 Cilia 2-3 . . . *B. crassirameum*, 329.

c. *Leaves pointed, costa vanishing.*

[*B. extenuatum* and *B. crassirameum* may be sought here.]

Leaves distinctly margined . . . *B. capillare*,² 235.

¹ See Appendix, nos. 330-332.

² See Appendix, nos. 317, 318, 344.

Leaves not margined or only indistinctly.

Closely appressed, imbricate.

- Broadly ovate or obovate, abruptly apiculate **B. argenteum**, 234.
 Ovate or lanceolate, not abruptly pointed **B. Blindii**, 309.
 Spreading, distant **B. Duvaillii**,¹ 238.

2. *Costa excurrent, leaves long-cuspidate.*

a. *Capsule short (1:2 or less).*

- Constricted between sporangium and collum **B. versicolor**, 233.
 Not constricted between sporangium and collum **B. coronatum**, 232.

b. *Capsule longer (1:3+).*

i. *Collum long, ½ sporangium or more.*

Leaves bordered with two or more rows of cells.

- Strongly twisted when dry, revolute **B. capillare**,² 235.
 Erect and straight when dry.

Capsule scarcely constricted under the mouth, collum costate, leaf margin revolute. **B. obconicum**, 236.

Capsule constricted under the mouth, collum smooth, leaf margin plane **B. elegans**, 319.

Leaves not or very indistinctly bordered.

Abruptly cuspidate, oblong spatulate or obovate lanceolate **B. Hendersoni**, 301.

Gradually cuspidate, lanceolate or ovate-lanceolate.

- Capsule constricted below the mouth **B. caespiticium**, 235.
 Capsule not constricted **B. Vancouveriense**, 315.

ii. *Collum short (¼ sporangium or less)* **B. occidentale**, 236.

§III. Rhodobryum.

- Costa stout, excurrent, leaf margins revolute $\frac{2}{3}$ to $\frac{3}{4}$ length **B. Ontariense**, 337.
 Costa vanishing, leaves plane **B. lucidum**, 338.

§IV. Anomobryum.

- Costa subexcurrent **B. concinatum**, 240.
 Costa vanishing below apex **B. bullatum**, 339.

The following species, described from sterile or immature specimens, are not included in the key, since the meager descriptions render it impossible

¹ See Appendix, nos. 334, 335.

² See Appendix, no. 317, 318, 344.

to determine their position. *B. leucolomatum* C. M. & Kindb., 300; *B. alpiniforme* Kindb., 310; *B. Floridanum*, R. & C., 320; *B. erythrophyllum* Kindb., 326; *B. erythrophyloides* Kindb., 327; *B. denticulatum* Kindb., 332; *B. hydrophyllum* Kindb., 333; *B. haematophyllum* Kindb., 340; *B. oligochloron* C. M. and Kindb., 341; *B. microcephalum* C. M. & Kindb., 342; *B. pygmaeo-alpinum* C. M. & Kindb., 343. Two other species, *B. Baueri* Hampe (California) and *B. Wrightii* Sull. (Behring Straits) are listed by Ren. & Card.

ZIERIA, p. 240.

Costa vanishing, collum twice sporangium *Z. julacea*, 240.
Costa excurrent, collum = sporangium *Z. demissa*, 241.

MNIUM, p. 241.

I. *Leaves serrate.*A. *Teeth of leaves single.*

Stems dendroid *M. Menziesii*, 240.
Stems simple or branched, not dendroid.
Basilar branches stoloniform.
Leaves acuminate, serrate to middle, lid convex or
mamilate, membrane of endostome lacunose *M. cuspidatum*,¹ 242.
Leaves acuminate, serrate to base.
Lid apiculate *M. medium*, 242.
Lid mammiform *M. affine*, 244.
Leaves rounded at apex, mucronate, lid rostrate *M. rostratum*, 243.
Basilar branches erect, or stems simple.
Capsule warty-papillose at base *M. venustum*,² 242.
Capsule smooth at base.
Leaves nearly entire not decurrent *M. affine rugicum*, 244.
Leaves serrate to base, long decurrent *M. insigne*,³ 244.
Leaves serrate above, entire below.
Border distinct, yellowish-brown or red.
Capsules clustered (2-3), leaf cells large and rect-
angular at the base *M. Drummondii*, 243.
Capsules solitary, leaf cells uniform *M. Blyttii*, 253.
Border none or faint *M. stellare*, 247.

¹ See Appendix, no. 345.² *M. macroclitäre* is insufficiently distinguished by Müller from this species. See Appendix, no. 346.³ See Appendix, no. 347.

B. *Teeth of the leaves in pairs.*[*M. Blyttii* may be sought here.]Costa vanishing below the apex *M. hornum*, 245.

Costa percurrent or excurrent.

Capsules solitary.

Synoicous.

Leaves decurrent *M. serratum*,¹ 245.Leaves not decurrent or only indistinctly so *M. Niagare*, 352.

Dioicous.

Costa excurrent in upper leaves, leaf cells 18-30 μ ,²plants 1.5-2 cm. high *M. pseudolycopodioides*, 351.

Costa percurrent.

Lid not rostrate.

Capsule with long neck, wide mouthed *M. inclinatum*, 350.Capsule oval, inclined *M. decurrens*, 349.

Lid rostrate.

Leaf cells small, about 15 μ ³ *M. orthorrhynchum*, 246.Leaf cells very large, 50-60 μ ⁴ *M. umbratile*, 243.Capsules clustered.⁵Dioicous, leaves strongly crispate, capsule horizontal
or inclined *M. spinosum*, 246.Synoicous, leaves not crispate, capsule pendent *M. spinulosum*, 247.II. *Leaves entire.*

Upper leaf-cells with long diameter oblique to costa.

Leaves bordered.

Costate to apex, dioicous, capsule oblong *M. punctatum*, 248.Costa vanishing, synoicous, capsule subglobose, leaves
distinct and few *M. subglobosum*, 248.Costa vanishing, dioicous, leaves more abundant and
closer *M. hymenophyllum*, 354.Leaves not bordered, costa vanishing, dioicous, capsule
ovate-oblong *M. cinctoides*, 248.Upper leaf-cells isodiametric, costa vanishing *M. hymenophylloides*, 249.*M. heterophyllum* Schw. is mentioned as occurring in America (Bryol. Eur. Mnium, p. 24), but no locality is given and for this reason it is not included in the key and descriptions.¹ See Appendix, no. 343.² As determined from Can. Musci, no. 197.³ Vide Husnot: Muscol. Gall. 235.⁴ Cells four times as large as *M. orthorrhynchum*, vide Mitten, Journ. Linn. Soc. 3: 30.⁵ *M. macroclitäre* may be sought here See footnote 2, p. 219.

CINCLIDIUM, p. 249.

- Leaf margin of 4-5 rows of cells, laminal cells irregularly disposed *C. stygium*, 250.
 Leaf margin of 2 rows of red cells, laminal cells in rows oblique to costa *C. subrotundum*, 250.

AULACOMNIUM, p. 252.

- Leaves coarsely serrate to middle, autoicous *A. heterostichum*, 253.
 Leaves serrulate near apex, acute or acuminate, dioicous.
 Stem leaves long acuminate, very roughly papillose *A. papillosum*, 253.
 Stem leaves acute.
 Stems commonly prolonged and gemmiferous, male
 flowers terminal, gemmiform *A. androgynum*, 252.
 Stems commonly not gemmiferous, male flowers discoid *A. palustre*,¹ 252.
 Leaves entire, obtuse *A. turgidum*, 253.
 [The leaves of *A. palustre* are entire when young, but soon become erose crenulate.]

TIMMIA, p. 254.

- Leaves larger above, sheath papillose on the back.
 Not hyaline at the insertion, monoicous *T. Megapolitana*, 254.
 Hyaline at the insertion, dioicous *T. Norvegica*, 253.
 Leaves uniform, sheath smooth on the back.
 Base hyaline, costa not toothed on back *T. Megapolitana Bavarica*, 257.
 Base orange, costa toothed above on the back *T. Austriaca*,² 254.

ATRICHUM, p. 255.

- Leaves margined, costa lamellose on upper side only.
 Lamellæ 2-6, entire, lamina with teeth on surface.³
 Lamellæ 4-6 cells high.
 Leaves serrate for $\frac{3}{4}$ length *A. undulatum*,⁴ 256.
 Leaves serrate above middle only.
 Teeth double, aculeate *A. angustatum*, 256.
 Teeth single, short *A. xanthopelma*, 257.

¹ See Appendix, nos. 355, 356.² See Appendix, no. 359.³ Excluding *A. xanthopelma*?⁴ See Appendix, no. 360.

- Lamellæ 9-13 cells high A. Selwyni, 256.
 Lamellæ 4-8, serrate A. Lescurii, 257.
 Lamellæ 1-3, 1-3 cells high, lamina smooth A. crispum, 257.
 Leaves not margined.
 Costa lamellose on both sides A. parallelum, 258.
 Costa lamellose on upper side only A. telephyllum, 261.
 Costa nearly smooth A. rosulatum, 262.

OLIGOTRICHUM, p. 258.

- Costa lamellose on both surfaces.
 Capsule rough when dry, leaf margin plane O. aligerum, 258.
 Capsule furrowed when dry, leaves nearly tubular by
 the incurved margin O. hercynicum,¹ 263.
 Costa only lamellose on upper surface O. Lyallii, 259.

POGONATUM, p. 260.

I. *Plants simple, mostly short, leaves straight when dry.*[*P. alpinum simplex* will be sought here.]

- Lamellæ with marginal cells smooth.
 Leaves entire P. brachyphyllum, 261.
 Leaves serrate P. brevicaulis, 260.
 Lamellæ with marginal cells papillose.
 Teeth of leaves very long, often reflexed, marginal cells
 of lamellæ subquadrate P. dentatum, 261.
 Teeth moderate, 2 rows of marginal cells of lamellæ
 transversely rectangular P. capillare, 261.

II. *Plants large (4-15 cm.), leaves twisted when dry.* (III)

- Leaves strongly contorted when dry, lamellæ 20-30.
 Less than 1 cm. long, short sheathing, capsule smooth.
 Leaves scarcely enlarged at the base, acute P. contortum, 262.
 Leaves enlarged at the base, abruptly pointed P. atrivirens, 262.
 More than 1 cm. long, not sheathing, capsule pap-
 illose P. erythrodontium, 265.
 Leaves spreading or patulose when dry, lamellæ
 about 60 P. Macounii, 267.

¹ See Appendix, no. 364.

A. Selwyni, 256.
A. Lescurii, 257.
A. crispum, 257.

A. parallelum, 258.
A. teleophyllum, 261.
A. rosulatum, 262.

III. *Plants usually robust (4-15 cm.), rarely small, often much branched above, leaves straight when dry.*

Capsule papillose, marginal cells of lamellæ round in section *P. urnigerum*, 264.
 Capsule smooth, marginal cells of lamellæ ovate in section *P. alpinum*,¹ 263.

POLYTRICHUM, p. 263.

Leaves entire, margins inflexed.

Obtuse at apex *P. sexangulare*, 269.

Aristate at apex.

Awn colored, short.

Leaves spreading when moist, subrecurved *P. juniperinum*,² 265.

Leaves erect-open, strict *P. strictum*, 265.

Awn hyaline, long *P. piliferum*, 264.

Leaves serrate.

Marginal cells of lamellæ like rest, oval, higher than broad in section.

Capsule ovate, obscurely angled, lid rostrate *P. gracile*, 264.

Capsule oblong, 4-6 angled, lid acutely conic *P. formosum*,³ 264.

Marginal cells of lamellæ enlarged, broader than high (2:1)

. *P. Ohioense*, 270.

Marginal cells of lamellæ semilunar, with two prominent papillæ at corners

P. commune,⁴ 266.

P. polare C. Müll., 273; *P. hyperboreum* R. Br.; *P. boreale* Kindb.; *P. leucopilum* Hpe., are not included in the key since no authentic material is at hand, and the descriptions are not sufficient to determine their position. The first three are arctic species and their omission is comparatively unimportant.

BUXBAUMIA, p. 267.

Outer peristome simple, spores 5-9 μ .

Capsule little longer than broad *B. aphylla*, 268.

Capsule much longer than broad (1:2-2.5) *B. Piperi*, 276.

Outer peristome quadruple or triple, spores 10-16 μ *B. induslata*, 275.

¹ See Appendix, no. 366.

² See Appendix, no. 374.

³ *P. conorhynchum* Kindb., evidently falls here. See Appendix, no. 368.

⁴ See Appendix, nos. 371, 372.

FONTINALIS, p. 268.¹I. *Leaves of branches unlike stem leaves (dimorphous).*

- Branch leaves 8-16 times as long as broad . . . *F. chrysophylla*, 333.
 Branch leaves 2.5-5 times as long as broad . . . *F. Kindbergii*,² 331.

II. *Leaves homomorphous.*³A. *Leaf cells rhombic-hexagonal (1:6 or less).*

Plants shining with golden or copper luster.

Stem robust, little branched . . . *F. antipyretica gigantea*,⁴ 269.

Stem soft, much branched . . . *F. California*, 269.

Plants dull, yellowish to dirty green.

Leaves with one edge reflexed near base . . . *F. antipyretica*, 268.

Leaves with margin plane.

Female flowers abundant, in most leaf axils . . . *F. Nova-Angliae*,⁵ 270.

Female flowers rare, at base of stems . . . *F. biformis*, 270.

B *Leaf cells long linear (1:7-30).*i. *Leaves auricled.*

[*F. Lescurii* and *F. Neo-Mexicana* may be sought here.]

Alar cells very large . . . *F. flaccida*, 333.

Alar cells moderately enlarged.

Leaves inflexed on the borders.

Gradually acuminate . . . *F. dichelymoides*, 335.

Cucullate and terminated by an acumen . . . *F. involuta*, 333.

Obtuse, neither acuminate nor cucullate . . . *F. Cardoti*, 333.

Leaves plane on the borders, acuminate.

Transverse bars of the endostome complete . . . *F. tenella*, 331.

Transverse bars of the endostome incomplete . . . *F. nitida*, 330.

¹ Revised by Cardot, Mem. Soc. Nat. d. Sci. Nat. et Math. de Cherbourg 28: 1-152. 1892.

² It has not been deemed advisable to maintain *F. Howellii* as a distinct species. Following the suggestion of Cardot, Monog. Fontin. 67, it is ranked as a variety of *F. Kindbergii*. See also Appendix, no. 382.

³ In *F. biformis* the summer leaves are unlike the vernal, so that specimens collected just as the vernal are falling might deceive.

⁴ See Appendix, nos. 377, 378.

⁵ See Appendix, nos. 386, 387.

2. *Leaves not auricled.** *Tubulose or sub-tubulose.*

- Apex obtuse and cucullate **F. Langloisii**, 397.
 Apex acute and not cucullate **F. filiformis**,¹ 271.

** *Concave and incurved on the borders.* (***)

Perichæatial leaves round-obtuse, entire or lacerate.

Teeth of 14-20 articulations, leaves oblong lanceolate **F. Delamarei**, 384.

Teeth of 28-32 articulations, leaves broadly oval . . . **F. mollis**, 385.

Perichæatial leaves abruptly pointed, entire . . . **F. Dalecarlica**, 270.

*** *Plane or concave but not incurved.*

Alar cells very large **F. Sullivantii**, 271.

Alar cells scarcely enlarged or only moderately.

Transverse bars of the endostome complete throughout.

Leaves decurrent, teeth not lacunose . . . **F. Neo-Mexicana**,² 269.

Leaves not decurrent, teeth generally lacunose.

Capsule contracted below mouth when dry . . . **F. hypnoides**, 272.

Capsule not contracted below mouth when dry . . . **F. Duriei**, 392.

Transverse bars of the endostome not complete.

Articulations of the teeth 12-20.

Endostome rudimentary, transverse bars not complete at any point **F. microdonta**, 394.

Endostome better developed, transverse bars complete above **F. disticha**, 272.

Articulations of the teeth 20-25 **F. Lescurii**, 271.

DICHELZYMA, p. 272.³

Costa percurrent or vanishing.

Capsule exceeding perichæatium **D. falcatum**, 273.

Capsule not exceeding perichæatium **D. pallescens**,⁴ 274.

¹ See Appendix, no. 396.

² *F. maritima* C. Müll., falls here, and is probably only a variety of *F. Neo-Mexicana*. See Appendix, no. 390; also no. 379.

³ Revised by Cardot, l. c.

⁴ Including *D. Novae-Brunswicæ* Kindb., and *D. obtusulum* Kindb. (Mac. Cat. 159), sec. Cardot, *ibid.* 143.

Costa excurrent.

- Erdostome a cancellate cone *D. uncinatum*,¹ 273.
 Erdostome of appendiculate cilia, united only at the
 tips *D. capillaceum*,² 273.

D. subulatum Myrin is *Brachelyma subulata* Sch. of which *Cryphæa inundata* Nees is also a synonym, sec. Cardot, *ibid.* 131.

CRYPHÆA, p. 275.

- Costa percurrent or excurrent *C. nervosa*, 277.
 Costa vanishing near middle.
 Costa of perichæial leaves excurrent into a thick point *C. glomerata*, 276.
 Costa of perichæial leaves vanishing in or below apex *C. pendula*, 276.
 Costa of perichæial leaves vanishing far below apex *C. Ravenelli*, 277.

LEPTODON, p. 278.

- Leaves ecostate.
 Leaf cells not pitted, capsule 2 mm. long *L. trichomitron*, 278.
 Leaf cells pitted, capsule 1 mm. long *L. Floridanus*, 414.
 Leaves costate.
 Leaf cells round-oval, capsule exerted, oblong-oval *L. Ohioensis*, 278.
 Leaf cells narrowly rhomboidal, capsule immersed, sub-
 globose *L. nitidus*, 279.

ALSIA, p. 279.

- Annulus none.
 Costa vanishing at middle, smooth(?), margins re-
 flexed *A. Californica*, 280.
 Costa vanishing near apex, dentate on back, margins
 plane *A. longipes*, 280.
 Annulus compound, revolute, leaves papillose at back *A. abietina*, 280.

NECKERA, p. 281.

- Leaves very obtuse.
 Plants slender (shoots 2 mm. wide), leaves loosely imbricate, rounded, concave *N. disticha*, 281.
 Plants robust (shoots 4 mm. wide), leaves densely imbricate, truncate, not concave *N. undulata*, 281.

¹ Including *D. cylindricarpum* Aust. as a variety sec. Cardot, *ibid.* 139.

² See Appendix, no. 998.

Leaves rounded, abruptly apiculate.

- Revolute at base on one side, capsule immersed . . . *N. Menziesii*,¹ 282.
 Not revolute, capsule exerted *N. complanata*, 283.

Leaves acute or acuminate.

Ecostate or nearly so.

Capsule immersed or half exerted.

- Shoots obtuse *N. pennata*, 282.
 Shoots attenuate to apex *N. oligocarpa*,² 283.

Capsule exerted.

- Segments as long as the teeth and similar . . . *N. Douglasii*,³ 283.
 Segments shorter, abruptly narrowed from a broad
 carinate plicate base *N. pumila*, 284.

Costate to the middle or beyond.

- Margins broadly revolute *N. Floridaana*, 284.
 Margins not revolute.
 Alar cells fawn-color, costa thin, percurrent . . . *N. Ludovicis*, 284.
 Alar cells opaque, costa vanishing *N. cymbifolia*, 284.

HOMALIA, p. 285.

- Costa single, vanishing above the middle, leaves serrate { *H. Macounii*, 403.
H. Jamesii,⁴ 285.
 Costa double, very short or none, leaves entire . . . *H. gracilis*, 286.

METEORIUM, p. 286.

- Leaves serrulate *M. pendulum*, 286.
 Leaves minutely crenulate *M. nigrescens*, 287.

LEUCODON, p. 287.

Capsule exerted.

- Leaves entire, open-erect, lid exactly conic . . . *L. sciuroides*, 288.
 Leaves serrulate at apex, squarrose, lid obliquely ros-
 trate *L. julaceus*, 288.

Capsule surpassed by perichæcial leaves, leaves secund *L. brachypus*, 288.

¹ See Appendix, nos. 399, 400.

² See Appendix, no. 401.

³ See Appendix, no. 402.

⁴ Including *H. trichomanoides* L. & J. Man. 235 (not Bruch. & Schimper), *H. obtusata* L. & J. (not Mitten). I am unable to detect from the descriptions alone any character which will separate *H. Jamesii* from *H. Macounii*. If type specimens show *H. Jamesii* to be the same its priority will make it the name for the type instead of *H. Macounii*.

PTERIGYNANDRUM, p. 288.

- Branch leaves acute, serrate or denticulate at the apex
 only *P. filiforme*,¹ 280.
 Branch leaves acute or acuminate, denticulate nearly all
 around *P. papillosum*, 406.

PTEROGONIUM, p. 289.

- Leaves broadly oblong-ovate or obovate, acute, smooth *P. gracile*, 290.
 Leaves broadly deltoid-ovate, narrowly acuminate, papil-
 lose *P. brachypterum*, 290.

ANTITRICHIA,² p. 290.

- Capsule oval (1:2-2.5), leaf cells fusiform . . . *A. curtispindula*, 291.
 Capsule cylindrical (1:6), leaf cells oval . . . *A. Californica*,³ 291.

HOOKERIA, p. 292.

- Leaves bicostate to middle (not papillose?) . . . *H. varians*, 292.
 Leaves bicostate to apex, papillose *H. cruceana*, 292.
 Leaves ecostate, entire (not papillose?) . . . *H. Sullivantii*,⁴ 293.

FABRONIA, p. 294.

- Leaves ciliate-dentate.
 Peristome of 16 teeth, costa none or very short . . . *F. pusilla*, 294.
 Peristome none, leaves costate to middle . . . *F. gymnostoma*, 294.
 Peristome of 8 geminate teeth, leaves costate nearly
 to middle *F. octoblepharis*, 295.
 Leaves serrate to subentire.
 Sharply serrate, teeth orange, spores about 11 μ . . . *F. Wrightii*, 295.
 Obscurely serrate, teeth brown, spores about 17 μ . . . *F. Ravenellii*, 295.
 Obscurely serrate, teeth with prominent articulations
 on back *F. Donnellii*, 295.

¹ See Appendix, no. 404.² A sterile species, *A. tenella*, Appendix, no. 406a, is probably referable to one of the two species here given.³ See Appendix, no. 406.⁴ "Die *Hookeria Sullivantii* mihi unterscheidet sich auch heute noch von *H. lucens* und ebenso von *H. acutifolia* aus Indien."—C. Müller, in litt. ad E. G. Britton, 18 Jul. 1888.

THELIA, p. 298.

- Papillæ of leaves simple.
 Horn shaped, curved T. hirtella, 299.
 Globose T. robusta, 299.
 Papillæ 2-4 furcate.
 Usually bifurcate, leaves ciliate T. asprella, 299.
 Usually 4-furcate, leaves not ciliate T. Lescurii, 299.
 Usually 3-4 furcate,¹ leaves ciliate T. compacta, 407.

MYURELLA, p. 300.

- Leaves serrulate, obtuse (rarely short apiculate) M. julacea, 300.
 Leaves serrulate, abruptly apiculate-acuminate M. apiculata, 300.
 Leaves spinulose-dentate, abruptly long-acuminate M. Careyana, 300.

LESKEA, p. 302.

I. *Costa reaching to or beyond the middle.*

- Percurrent L. nervosa, 302.
 Not percurrent.
 Leaves entire.
 Endostome divided into segments.
 Cleft between articulations, leaves bluntish.
 Leaves ovate at base, and narrowed above L. obscura, 301.
 Leaves oblong or broadly ovate L. subobtusifolia, 408.
 Not cleft, leaves acute L. polycarpa, 301.
 Endostome a short undivided membrane L. Austini, 303.
 Leaves crenulate L. tristis, 303.

II. *Costa very short or none.*

- Leaf cells linear oblong L. denticulata, 302.
 Leaf cells rotundate L. cyrtophylla, 409.

MYRINIA, App., no. 410.

- Leaves with faint traces of a costa at the base M. pulvinata,² 303.
 Leaves with broad costa, sometimes bifurcate, ceasing
 below apex M. (?) Dieckii, 417.

¹ No. 248 Canadian Musci shows this character.

² *L. pulvinata* Wahl., L. & J. Man. 303.

apex
 P. filiforme,¹ 280.
 y all
 P. papillosum, 405.

th P. gracile, 290.
 papil-
 P. brachypterum, 290.

A. curtipendula, 291.
 A. Californica,² 291.

H. varians, 292.
 H. cruceana, 292.
 H. Sullivantii,⁴ 293.

F. pusilla, 294.
 F. gymnostoma, 294.
 F. octoblepharis, 295.

F. Wrightii, 295.
 F. Ravenellii, 295.
 F. Donnellii, 295.

referred to one of the

noch von *H. lucens* und
 G. Britton, 13 Jul. 1888.

ANOMODON, p. 304.

Leaves not papillose A. Toccoe, 306.
 Leaves papillose.

Base with large fimbriate-papillose auricles.

Margins reflexed near apex, replicate below middle A. Californicus, 306.

Margins not at all reflexed A. apiculatus, 306.

Base not auriculate.

Leaves filiform acuminate.

Decurrent, paraphyllia broad A. heteroideus, 413.

Not decurrent, paraphyllia none A. rostratus, 305.

Leaves obtuse or apiculate.

Branches attenuate A. attenuatus,¹ 305.

Branches not attenuate.

Leaves open-erect, teeth nodose A. obtusifolius, 306.

Leaves secund, teeth not nodose A. viticulosus, 306.

PYLAISÆA, p. 308.

Segments free, split below, leaves quite entire or denticulate at apex.

Leaves ecostate or faintly costate at the base.

Plants glossy green P. polyantha, 308.

Plants pale yellowish green P. heteromalla, 308.

Leaves distinctly bicostate P. pseudoplatygyrium, 417.

Segments for $\frac{1}{2}$ or less adherent to the teeth.

Leaves ovate lanceolate, acuminate, margin not re-

curved P. intricata,² 309.

Leaves filiform acuminate P. filari-acuminata, 419.

Leaves short acuminate, one or both edges recurved P. Selwynii, 418.

Segments wholly adherent P. velutina, 309.

HOMALOTHECIUM, p. 309.

Costa short, simple or forking, vanishing below middle.

Teeth red, operculum rostrate H. subcapillatum, 310.

Teeth yellow, operculum short apiculate H. corticolum, 422.

¹ See Appendix, no. 412.

² From the description given I am unable to separate *P. Ontarioensis* C. M. and Kindb. See Appendix, no. 419a.

Costa narrow, vanishing at point.

Alar cells quadrate.

Leaves lanceolate, secund, carinate by the costa *H. Nevadaense*,¹ 332.

Leaves long subulate or filiform acuminate, neither

secund nor canaliculate *H. sericeum*,² 420.

Alar cells not differing from the rest *H. pseudosericeum*, 310.

CYLINDROTHECIUM,³ p. 310.

Capsules clustered (3 or 4) *C. Floridaum*, 312.

Capsules solitary.

Plants densely pinnately branched, leaves mucous *C. concinnum*, 313.

Plants loosely pinnately branched, leaves pointed.

Gradually narrowly acuminate *C. brevisetum*, 311.

Acute or abruptly acuminate-apiculate.

Almost entire, only alar cells quadrate or rectangular.

Leaves acuminate-apiculate, teeth with 14-17 articulations, capsule 1: 3.5-4 *C. cladorrhizans*, 310.

Leaves abruptly short apiculate, teeth with 6-8 articulations, capsule 1: 5-5.5 *C. seductrix*, 310.

Leaves not apiculate, teeth with 22-26 articulations, capsule 1: 2.5-3 *C. compressum*, 312.

Distinctly serrulate, all basal cells rectangular.

Annulus none, teeth obliquely striolate *C. Drummondii*, 312.

Annulus large, teeth vertically striolate *C. Sullivantii*, 313.

CLIMACIUM, p. 313.

Capsule straight, lid rostrate.

Ovate-oblong (1: 2.5-3), leaves slightly decurrent and hollowed at basal angles *C. dendroides*,⁴ 314.

Cylindric (1: 5-6), leaves long decurrent and broadly auriculate *C. Americanum*,⁵ 314.

Capsule arcuate, lid conic *C. Ruthenicum*, 314.

¹ *Hypnum* (*Camptothecium*) *Nevadense* L. and Man. 332. See also Appendix, no. 421.

² A closely allied but barren species is *H. sericeoides* C. M. and Kindb. See Appendix, no. 420a.

³ Three species belonging to this genus are described by Müller & Kindberg from barren specimens under the name *Entodon* Müll. *C. Macounii* is closely related to *C. Drummondii*; the others are *C. actulare* and *C. exallens*. See Appendix, nos. 423-425.

⁴ See Appendix, no. 426.

⁵ See Appendix, no. 427.

ORTHOTHECIUM, p. 315.

- Leaves lanceolate, long and narrowly acuminate . . . *O. rufescens*, 315.
 Leaves exactly ovate, apex flexuous, not plicate . . . *O. rubellum*, 315.
 Leaves lanceolate to ovate lanceolate, plicate, not acumi-
 nate *O. chryseum*, 316.
 Leaves long subulate *O. intricatum*, 428.

SEUDOLESKEA,¹ p. 319.

- Costa percurrent, leaves serrate at apex *P. rigescens*,² 320.
 Costa short, double, or none, leaves entire.
 Alar cells transversely elongated *P. tectorum*, 455.
 Alar cells quadrate *P. malacoclada*, 456.
 Costa single, vanishing below apex, leaves serrate and entire.
 Leaves long decurrent *P. falcicuspis*, 455.
 Leaves not decurrent.
 Lanceolate from an ovate base *P. atrovirens*,³ 319.
 Narrowly lanceolate *P. stenophylla*, 454.

P. oligoclada Lindb., has been reduced by Ren. & Card. to *P. atrovirens* Sch. *Hypnum radicosum* Mitt. is identified by them with *P. rigescens* Lindb. to which *P. sciuroides* Kindb. has also been added as a synonym. See Mac. Cat. 180, and Ren. & Card. Revue Bryol. 20: 15. 1893.

HETEROCLADIUM, p. 320.

- Leaves papillose.
 Coarsely toothed, mouth of capsule bordered by three
 rows of transversely elongated cells *H. heteropterum*, 437.
 Faintly serrulate, mouth of capsule not bordered *H. dimorphum*, 321.
 Denticulate above the middle.
 Leaves dimorphous and faintly papillose *H. Vancouveriense*, 438.
 Leaves homomorphous and strongly papillose *H. frullaniopsis*, 439.
 Leaves smooth.
 Auriculate and squarrose *H. aberrans*, 440.
 Not auriculate, divergent *H. procurrens*, 321.

¹ *P. catenulatum* Brid. has never been found in America; the locality given in L. & J. Man. 320, as Mt. Ingleborough, New York, is York, England.

² See also Appendix, no. 432.

³ See Appendix, nos. 429-431.

THUIDIUM, p. 321.

I. *Plants small (to 5 cm.), delicate, creeping, 1-2 pinnate.*Costa of stem leaves wide ($\frac{1}{2}$ leaf base).Branches papillate, leaf cells 6μ diam. *T. pygmaeum*, 322.Branches papillate, leaf cells 9μ , longer at margin *T. minutulum*, 322.Costa of stem leaves narrow ($\frac{1}{4}$ leaf base).Leaf cells with several minute papillae *T. scitum*,¹ 323.

Leaf cells with one papilla (rarely two).

Branch leaves roundish ovate, short acuminate *T. Virginianum*,² 324.Branch leaves ovate lanceolate, long acuminate *T. microphyllum*,³ 324.II. *Plants larger (to 10 cm.), creeping, 1-3 pinnate, forming extensive flat mats. (III)*

Perichaetial leaves ciliate, costa of stem leaves not reaching point

T. delicatulum, 325.

Perichaetial leaves not ciliate.

Costa of stem leaves filling point *T. recognitum*, 325.Costa of stem leaves $\frac{2}{3}$ length *T. Philiberti*, 443a.[*T. paludosum* may be sought here.]III. *Plants large (to 10 cm.), erect or ascending, 1-pinnate, in wide tufts.*

Leaf cells roundish or oblong (1:1-2), strongly papillose

both sides *T. abletinum*, 326.

Leaf cells long rhombic to linear (1:3-6), smooth above.

Stem leaves soft, sub-clasping, decurrent *T. Blandovii*, 326.Stem leaves rigid, plicate-striate, sub-decurrent *T. paludosum*,⁴ 330.

This genus has been revised by G. N. Best, Bull. Torr. Bot. Club 28: 78-90. 1896, and we have been able to avail ourselves of his suggestions by recalling MS. *T. tamariscinum* Hedw. and *T. remotifolium* Grev. are not N. American; *T. erectum* Duby = *T. delicatulum* L.; *H. calyptratum* Sull. = *T. microphyllum* (Sw.) Best, an earlier name for *H. gracile* B. & S., and to it *T. lignicola* Kindb. is referred as a variety; *T. Alteni* Aust. is probably a sterile form of *T. delicatulum* L.

¹ See Appendix, no. 441. Not recognized by Best.² *T. Virginianum* (Brid.) Lindb. = *Hypnum gracile* Lancastrise S. & L., L. J. Man. 324.³ *T. microphyllum* (Sw.) Best = *Hypnum gracile* B. & S., L. & J. Man. 324. See also Appendix, no. 442.⁴ *T. paludosum* (Sull.) Rau & Hervey = *Hypnum paludosum* Sull., L. & J. Man. 330. See also Appendix, no. 443.

O. rufescens, 315.
O. rubellum, 315.
 imi-
O. chryseum, 316.
O. intricatum, 428.

P. rigescens,² 320.

P. tectorum, 436.
P. malacoclada, 436.
 entire.
P. falceuspsis, 435.

P. atrovirens,³ 319.
P. stenophylla, 434.


Card. to *P. atrovirens* with *P. rigescens* been added as a synonym. 20: 15. 1893.



three
H. heteropterum, 437.
H. dimorphum, 321.

H. Vancouveriense, 438.
H. frullianlopsis, 439.




H. aberrans, 440.
H. procurrens, 321.

locality given in L. & J.





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1981

CLAOPODIUM, p. 327.

- Seta smooth *C. leuconeurum*, 328.
 Seta rough.
 Perichæatial leaves costate *C. Whippleanum*, 328.
 Perichæatial leaves ecostate.
 Bright green, leaves open, loosely imbricate *C. laxifolium*, 329.
 Dirty or yellowish green, leaves subfalcate secund,
 closely imbricate *C. crispifolium*,¹ 329.

TRIPTEROCLADIUM, p. 330.

- Leaves shortly bicostate.
 Obscurely denticulate at the apex *T. leucocladulum*, 330.
 Distinctly denticulate all around *T. compressulum*, 331.
 Leaves ecostate *T. rupestre*, 444.

CAMPTOTHECIUM, p. 331.

- Plants regularly pinnate.
 Stems erect, stout (to 15 cm.) *C. megaphyllum*, 334.
 Stems prostrate.
 Leaf base entire, alar cells abundant.
 Capsule oblong, segments split below *C. aureum*,² 447.
 Capsule long cylindrical, segments split throughout *C. Amesii*, 448.
 Leaf base denticulate, alar cells few *C. Nuttallii*,³ 332.
 Plants irregularly branched, stout, in extensive mats.
 Seta rough.
 Leaves ovate-lanceolate (1:3), cilia 3, as long as the
 segments *C. saeum*,⁴ 331.
 Leaves long lanceolate (1:5), cilia 1-2, long or short,
 operculum rostrate *C. lutescens*, 331.
 Seta smooth *C. nitens*, 333.
 Plants irregularly branched, slender, cilia 1-2, short oper-
 culum obtusely conical *C. arenarium*, 333.

¹ Including *H. ramulosum* Hpe., L. & J. Man. 328.² *H. pinnatifidum* S. & L., L. & J. Man. 338, is reduced by Ren. & Card. to a variety of *C. aureum* Sch. Hedw. 39: 255. 1893.³ *C. hamatidens* Kindb. Mac. Cat. 189, sec. Ren. & Card., *ibid.* 254. See also Appendix, no. 449.⁴ See Appendix, no. 446.

BRACHYTHECIUM, p. 334.

I. *Seta smooth.*[*B. senecipes* and *B. mirabundum* may be sought here.]* *Cilia rudimentary or none.*

- Perichæstrial leaves costate *B. acuminatum*,¹ 336.
 Perichæstrial leaves ecostate.
 Inflorescence synoicous *B. Utahense*, 339.
 Inflorescence dioicous or monoicous.
 Capsule oval, cilia none *B. Donnellii*, 338.
 Capsule cylindric oblong.
 Leaf costate to middle, basal cells very small *B. biventrosus*, 338.
 Leaf costate $\frac{3}{4}$ the length, basal cells dilated *B. laevisetum*, 469.

** *Cilia well developed, appendiculate.* (***)

- Perichæstrial leaves abruptly long pointed *B. latum*,² 335.
 Perichæstrial leaves gradually long pointed *B. Mildeanum*,³ 337.

*** *Cilia well developed, not appendiculate.*

- Annulus compound *B. collinum*, 339.
 Annulus simple, narrow.
 Leaves acute or acuminate, serrulate.
 Leaves straight when dry, capsule short (1-1.5), monoicous *B. salebrosum*,⁴ 336.
 Leaves twisted flexuous when dry, capsule short, (1: 1.5-2) *B. Thedenii*, 338.
 Leaves acute or acuminate, entire or slightly serrate at apex only.
 Open, alar cells oblong.
 Monoicous *B. salebrosum*,⁴ 336.
 Dioicous *B. glareosum*, 468.
 Appressed imbricate, alar cells quadrate *B. albicans*,⁵ 337.

¹ See Appendix, nos. 455, 456.² See Appendix, nos. 450-452.³ *B. Mildeanum* Sch. = *B. salebrosum palustre* L. & J. = *B. acutum* Mitt., sec. Ren. & Card. Rev. Bry. 30: 17, 1903.⁴ See Appendix, nos. 461, 462, 463.⁵ See Appendix, no. 459.

- Capsule with evident collum *B. glaciale*, 481.
 Cells of basal angles distinct, small or dilated, quadrate
 or oblong rectangular.
- Leaves decurrent.
- Broad ovate, faintly striate, dioicous *B. platycladum*, 475.
 Triangular ovate, deeply plicate, monoicous.
- Seta long (1.5 cm.), faintly rough, teeth ser-
 rulate *B. leucoglancum*, 484.
 Seta long (2 cm.), very rough, teeth not ser-
 rulate *B. Columbo-rutabulum*,¹ 487.
 Seta short (5-8 mm.), faintly rough *B. mirabundum*, 489.
- Leaves not decurrent.
- Capsule short or roundish oval.
- Horizontal, pedicel arcuate above *B. Bolanderi*, 341.
 Oblique, pedicel straight *B. gemmascens*, 483.
- Capsule oblong cylindrical.
- Inner perichaetial leaves very abruptly
 pointed *B. trachypodium*, 473.
 Inner perichaetial leaves gradually acuminate *B. asperrium*,² 343.

The following species are not included in this key, since the descriptions are taken from sterile or immature specimens or are otherwise so incomplete as not to offer sufficient characters for their proper location: *H. Coloradense* Aust., Man. 412; *B. Fitzgeraldi* C. Müll., 454; *B. Realii*, R. & C., 457; *B. harpidioides* C. M. & K., 460; *B. pseudoalbicans* Kindb., 464; *B. spurio-acuminatum* C. M. & K., 465; *B. pseudo-collinum* Kindb., 468; *B. latifolium* (Lindb.) R. & C., 472; *B. pseudo-Starkei* R. & C., 473; *B. spurio-rutabulum* C. M. & K., 476; *B. Villardi* R. & C., 490; *B. cirrhosum* Sch., 491.

SCLEROPODIUM, p. 346.

- Cilia solitary, rudimentary *S. Krausel*, 493.
 Cilia 2-3, as long as the segments.
- Seta rough throughout.
- Capsule erect or oblique, stem leaves gradually ac-
 minate *S. caespitosum*,³ 346.
 Capsule horizontal, leaves abruptly short acuminate
 with point recurved *S. Illicebrum*, 347.
 Seta rough and reddish above, smooth and yellow
 below *S. Californicum*, 346.

¹ From description I cannot distinguish *B. lamprochryseum*. See Appendix, no. 488.

² *Hypnum vallium* S. & L. is identical *sec.* Sull. Ic. Muse. Suppl. 100.

³ See Appendix, no. 492.

ISOTHECIUM, p. 347.

Leaves papillose on back.

Cilia solitary, margin of stem leaves reflexed I. *spiculiferum*, 348.Cilia 2-3, margin of stem leaves not reflexed I. *stoloniferum*, 348.

Leaves smooth on the back.

Perichaetial leaves costate.

Upper branch leaves acute or acuminate, entire I. *apocladum*, 350.

Branch leaves acute or acuminate, serrate all

around I. *myosuroides*, 347.

Perichaetial leaves ecostate.

Seta rough I. *lentum*, 350.

Seta smooth.

Alar cells dark yellow or orange, quadrate or rect-

angular I. *Cardoti*, 494.

Alar cells not colored, quadrate or round quadrate

or not differing.

Alar cells obscure, costa vanishing in middle.

Cilia equaling segments, capsule inclined I. *acutiusculis*, 349.Cilia shorter, capsule erect I. *Brewerianum*,¹ 349.Alar cells distinct, costa $\frac{3}{4}$ length of leaf I. *myurellum*, 495.

EURHYNCHIUM, p. 351.

I. *Seta smooth.*Points of leaves twisted, plants golden yellow E. *Boselli*, 352.

Points of leaves straight.

Leaves not decurrent, cilia not appendiculate.

Spreading, branches attenuate E. *strigosum*,² 351.Appressed, branchlets short, julaceous E. *diversifolium*, 352.Leaves long decurrent, cilia appendiculate E. *substrigosum*, 498.II. *Seta rough.*

Leaves with filiform points.

Stems short, with erect fasciculate branches, stoloni-

ferous E. *Vaucheri*, 414.

Stems long, prostrate, irregular, branched, not radicu-

lose E. *piliferum*, 353.¹ Leaves not papillose, *teste* M. A. Howe *in litt.* Including *H. aggregatum*, Mitt., L. & J. Man. 350 sec. R. & C. Rev. Bryol. 20: 20. 1893.² See Appendix, nos. 496, 497.

Leaves not filiform.

Leaves serrulate all around.

Decurrent, excavate at basal angles.

Perichætal leaves spreading *E. Stokesii*, 354.Perichætal leaves reflexed *E. Oregonum*, 355.

Not decurrent nor excavate.

Leaves ovate lanceolate, acuminate, segments
split *E. Sullivantii*,¹ 353.Leaves broad ovate, acute, segments perforate *E. praelongum*, 353.Leaves broad ovate, acute, segments split *E. hians*, 354.

Leaves entire at the base.

Decurrent *E. semiasperum*, 502.

Not decurrent.

Lid not half as long as the capsule *E. colpophyllum*, 342.Lid nearly as long as the capsule *E. hians*, 354.

E. Dawsoni Kindb., App. 501, and *E. crassinervium laxirete* Kindb., App. 500, are described from sterile specimens; hence they are not included in this key.

RAPHIDOSTEGIUM, p. 355.

I. *Operculum long subulate rostrate.*

Leaf margin entire.

Ovate or ovate lanceolate, monoicous *R. demissum*, 355.Long subulate, dioicous *R. subdemissum*, 504.

Leaf margins serrate or denticulate.

Filiform acuminate.

Cilia none or rudimentary *R. laxepatum*, 358.Cilia 2, stout, nearly equaling segments *R. recurvans*, 356.

Acute or short acuminate.

Decurrent *R. Roellii*,¹ 508.

Not decurrent.

Capsule short, oval or oblong (1:2) *R. microcarpum*, 357.Capsule longer, cylindric (1:4) *R. subadnatum*, 506.II. *Operculum short rostrate or conic.*

Leaves filiform acuminate.

Cilia none, annulus none, capsule cylindric *R. cylindrocarpum*, 356.Cilia 1, annulus simple, large, capsule oblong *R. Jamesii*, 357.¹ See Appendix, no. 503.

Leaves acute or short acuminate.

- Diclicous, leaves obovate or subrotund *R. Novæ-Cesareæ*, 356.
 Monocous, leaves broadly ovate to ovate or oblong lanceolate.
 Leaves decurrent *R. Roellii*,¹ 508.
 Leaves not decurrent.
 Margin strongly reflexed *R. Kegellanum*, 507.
 Margin plane *R. micans*,² 365.

RHYNCHOSTEGIUM, p. 358.

- Pedicel rough *R. curvisetum*, 360.
 Pedicel smooth.
 Costa single, reaching half way or more.
 Leaves apparently 2-ranked, plants of dry woods *R. serrulatum*, 359.
 Leaves spreading every way *R. rusciforme*, 359.
 Costa very short or none or double.
 Bicostate, annulus large *R. geophilum*, 358.
 Uni- or ecostate, annulus none *R. deplanatum*, 359.

Hypnum Caloosiense Aust., *H. Royæ* Aust., and *H. Brandegei* Aust., insufficiently established and of uncertain relationships, are not included in this key.

THAMNIUM, p. 361.

- Branch and stem leaves apparently 2-ranked, complanate.
 Capsule oblong, without collum *T. Holzingeri*, 511.
 Capsule oval, with a distinct collum *T. Bigelovii*, 362.
 Branch and stem leaves equally spreading.
 Perichaetial leaves reflexed.
 Cilia equaling teeth *T. Leibergii*, 510.
 Cilia short *T. neckeroides*, 362.
 Perichaetial leaves erect.
 Teeth with a hyaline, dentate margin *T. alopecurum*, 509.
 Teeth not hyaline margined and toothed only at the articulations *T. Alleghaniense*, 362.

¹ *R. Roellii* is included under both heads since the character of the operculum is unknown to me.

² *R. micans* E. & C. Rev. Bryol. 20: 21. 1898 = *Hypnum micans* Sw. L. & J. Men. 365. See also Appendix, no. 505.

PLAGIOTHECIUM, p. 362.

I. *Leaves complanate.** *Lid rostrate.*

- Leaves transversely undulate, serrulate at the apex *P. undulatum*, 369.
 Leaves not undulate, quite entire *P. sylvaticum*, 368.

* * *Lid conic or convex.*

- Capsule pendent or sub-pendent, seta arcuate *P. elegans*, 366.
 Capsule suberect, inclined or horizontal, often arcuate.
 Sulcate and constricted below the mouth when dry *P. turfaccum*, 366.
 Smooth when dry.
 Leaves serrulate, capsule sub-cylindric *P. Sullivantiae*, 368.
 Leaves quite entire.
 Capsule obovate, campanulate when dry *P. Muellerianum*, 367.
 Capsule oblong, constricted under the mouth when dry.
 Costa bipartite, vanishing half way to the middle *P. denticulatum*,¹ 367.
 Costa simple, short, or none *P. brevipungens*, 515.

II. *Leaves equally spreading, straight.* (III)* *Alar cells abruptly enlarged.*

- Leaves acute or short acuminate.
 Flat, point broad and straight *P. membranosum*, 514.
 Concave, point needle-like and recurved or patent *P. acicularipungens*, 516.
 Leaves filiform acuminate.
 Cilia none *P. piliferum*, 364.
 Cilia 2-3.
 Branches erect, leaves serrate *P. Muhlenbeckii*, 370.
 Branches intricate, leaves nearly entire *P. Fitzgeraldi*, 370.

* * *Alar cells scarcely different, quadrate or oblong, not abruptly enlarged.*+ *Capsule suberect, smooth when dry.*²

- Dioicous, cilia none, costa obsolete.
 Inner perichastial leaves ovate lanceolate *P. latebricola*, 363.
 Inner perichastial leaves abruptly acuminate *P. Passaicense*, 363.

¹ See Appendix, nos. 512, 513.² *P. pseudo-latebricola* has an inclined capsule.

Autolcous, cilia 2-3, costa double.

Costa thick, ascending to the middle *P. geminum*, 365.

Costa thin, reaching half way to the middle *P. denticulatum*, var. 367.

+ + *Capsule inclined or pendent, sulcate when dry.*¹

Leaves oval, narrowly acuminate, monolcous *P. pseudo-Silesiacum*, 370.

Leaves ovate lanceolate, diolcous.

Suddenly tapering to a long filiform point *P. pseudo-latebricola*, 519.

Short acuminate *P. decursivifolium*, 517.

III. *Leaves secund.*

Costæ 2, thick, reaching the middle *P. geminum*, 365.

Ecostate or shortly bicostate.

Leaves entire, seta straight *P. pulchellum*, 364.

Leaves serrulate above, seta arcuate *P. Silesiacum*, 518.

P. bifariellum Kindb., App. 520, and *P. attenuatirameum* Kindb., App. 521, described from barren specimens, are not included in the key.

AMBLYSTEGIUM,² p. 371.

I. *Leaves ecostate or with obscure traces of a nerve.*

Cilia none.

Perichaetial leaves long and narrow acuminate, entire *A. subtile*, 372.

Perichaetial leaves oval or oblong lanceolate, entire at the apex *A. Sprucei*, 373.

Cilia 1-2.

Plants minute, filiform (1-2 cm.).

Leaves ovate, long acuminate *A. confervoides*, 372.

Leaves long lanceolate *A. minutissimum*, 371.

Plants large, in wide flat tufts *A. adnatum*, 375.

II. *Leaves plainly costate.*

A. *Leaves with a distinct border* *A. Lescurii*, 376.

¹ *P. pseudo-latebricola* has a smooth capsule.

² The assistance of PROF. L. S. CHENET, who is engaged upon a revision of this genus is gratefully acknowledged.

B. *Leaves not bordered.** *Costate to the apex.*

- Leaves acuminate, basal cells abruptly enlarged *A. irriguum*, 374.
 Leaves acuminate, basal cells not enlarged.
 Serrate throughout *A. compactum*, 375.
 Entire or obscurely serrulate above.
 Branches mostly erect, leaves loosely spreading when
 dry *A. orthocladon*, 374.
 Branches mostly spreading, tufts intricate, leaves im-
 bricate when dry *A. varium*,¹ 373.
 Leaves not acuminate, ovate to oblong lanceolate *A. fluviatile*, 375.

* * *Costa ceasing at the middle or above.*[*A. compactum* may be sought here.]

- Cells near middle of leaf 1: 10-15.
 Leaves long acuminate, apex sharp *A. riparium*,² 376.
 Leaves acute, apex blunt *A. vacillans*, 377.
 Cells near middle of leaf less than 1: 8.
 Inner perichaetial leaves short acuminate ($\frac{1}{4}$ length).
 Leaves .65-.85 \times .26-.36 mm. *A. serpens*,³ 373.
 Leaves 1.2-1.6 \times .5-.7 mm. *A. Kochii*, 535.
 Inner perichaetial leaves subuliform acuminate, cells
 vermicular *A. porphyrrhizon*, 527.

Kindberg has described four species belonging to this genus from barren specimens. *A. fenestratum*, App. 522, is probably not an Amblystegium. *A. speirophyllum*, App. 523, may be doubtfully referred from specimens in Can. Musci to *A. irriguum*. *A. distantifolium*, App. 529, is very near *A. irriguum*. *A. dissitifolium*, App. 530, and *A. subcompactum* C. M. & Kindb., App. 531, are undoubtedly *A. compactum* Müll. *A. Holzingeri* R. & C. is *Hypnum* (Limnobium) *Closteri* Aust. *A. Floridanum* R. & C., App. 534, is so poorly characterized that I am unable to separate it from *A. riparium*. *A. Juratzkanum*, App. 525, *A. hygrophilum* (an Amblystegium?), App. 526, and *A. Schlotthaueri*, App. 528, are subspecies of *A. serpens*, sec. Cardot, Hedwigia 82: 270, 271. 1893. *A. homalostegium* Jgr. & Sauerb., App. no. 536, is so doubtful as an Amblystegium (*ex descr.*) that we omit it.

¹ *Hypnum radtoale* L. & J. Man. 373.² See Appendix, no. 532, 533.³ See Appendix, no. 524-526, 528.

HYPNUM.¹

I. *Leaves spreading, or complanate, not secund.* (II on p. 247.)

A. *Leaves costate half way or more.* (B on p. 245.)

1. *Leaves acute or acuminate.*

Margins denticulate.

Decurrent and auricled, suddenly acuminate *H. chloropterum*, 562.

Neither decurrent nor auricled, gradually acuminate.

Cells uniform throughout *H. Columbiæ*, 542.

Alar cells dilated, distinct *H. Columbiæ-palustre*, 588.

Margins entire.

Leaves acute or short acuminate.

Alar cells few.

Costa $\frac{1}{2}$ length or double and short *H. palustre*, 398.

Costa vanishing just below apex *H. polare*, 589.

Alar cells more numerous, reaching costa,
large *H. pseudo-montanum*, 593.

Leaves long acuminate.

Erect or spreading *H. polgamum*,² 379.

Reflexed squarroulose.

Decurrent *H. decursivulum*,³ 541.

Not decurrent.

Plants in dense tufts *H. uniconstatum*,⁴ 540.

Plants in loose, intricate tufts *H. chrysophyllum*,⁵ 378.

2. *Leaves obtuse,⁶ entire.*

* *Cells enlarged at the basal angles.*⁷ (* * on p. 245.)

Costa sub-percurrent.

Monoicous, sparingly branched, alar cells gradually enlarged *H. cordifolium*, 402.

¹ Including subgenera XVII to XXVI of L. & J. Man. 377-406.

² See Appendix, nos. 544, 545.

³ The separation of these three species is very unsatisfactory. *H. chrysophyllum* is a very variable species and from the inspection of barren material of *H. decursivulum* and *H. uniconstatum* and comparison with the characters as given by Kindberg, it seems probable that they can not be regarded as distinct species but only as forms of *H. chrysophyllum*.

⁴ Sometimes apiculate in *H. sarmentosum*.

⁵ *H. torrentis* C. M. & Kindb., Appendix, no. 586, falls here.

Dioicous, profusely branched.

- 5-10 cm. long, variegated or dark purple, stolons
 green *H. sarmentosum*, 403.
 15-30 cm. long, bright to yellowish green *H. giganteum*, 403.

Costa reaching middle.

- Branches irregularly pinnate, leaves spreading *H. Richardsoni*, 404.
 Branches few, leaves imbricate *H. stramineum*,¹ 405.

* * *Cells not enlarged at the basal angles.*

Leaf margin serrulate above, leaf cells short *H. occidentale*, 598.

Leaf margin entire, leaf cells vermicular, bordered by a
 row of short cells.

- Leaves open *H. arcticum*, 400.
 Leaves closely imbricate *H. trifarium*, 405.

B. *Costa very short or none or double.*

1. *Alar cells abruptly enlarged (often inflated or colored).**
 (2 on p. 246.)

* *Operculum short rostrate.*

- Leaves entire *H. Haldanianum*,² 397.
 Leaves sharply serrate *H. nemorosum*, 398.

* * *Operculum convex or conc.*

Leaves falcate.

[*H. palustre hamulosum* may be sought here.]

- Scarcely costate, alar cells orange *H. eugyrium*,⁴ 401.
 Costa reaching middle, alar cells hyaline *H. ochraceum*,⁵ 401.
 Leaves not falcate.

Gradually filiform acuminate, alar cells orange.

- Plants irregularly branched *H. Sommerfeltii*,⁶ 537.
 Plants sub-dichotomously branched *H. stellatum*,⁷ 379.

¹ See Appendix, no. 596-597.

² Three barren species described by C. M. & Kindb. belong to this division: *H. flaccum*, Appendix, no. 580; *H. subflaccum*, Appendix, no. 581; *H. pseudo-drepanum*, Appendix, no. 582.

³ See Appendix, no. 579.

⁴ See Appendix, nos. 590, 591.

⁵ See Appendix, no. 592.

⁶ See also Appendix, no. 538.

⁷ See Appendix, no. 543.

Acute or short apiculate, alar cells few, large . . . *H. palustre*, 398.
 Obtuse, entire, alar cells hyaline *H. cuspidatum*, 403.

2. *Alar cells scarcely different or quadrate or rectangular, not abruptly enlarged.*

* *Leaves thin, glossy, open; plants mostly small, prostrate or with ascending branches.*

Leaves squarrose, acuminate.

Sub-serrulate all around, lid apiculate *H. hispidulum*, 378.

Entire except the acumen, lid obtuse *H. Macounii*, 539.

Leaves loosely imbricate, obtuse or acute.

Serrulate, at least above.

Nearly as broad as long, obtuse or apiculate.

Alar cells small, not forming distinct auricles,

others 1:8-10 *H. molle*, 399.

Alar cells large, forming distinct auricles, others

much longer than preceding *H. dilatatum*,¹ 601.

Nearly twice as long as broad, acute, point often half

twisted *H. alpestre*, 399.

Entire.

Leaves close set, ovate (1:2) *H. Norvegicum*, 587.

Leaves more distant, rotund *H. Gouardii*, 585.

** *Leaves firm; plants very large, mostly 1-2 pinnate, erect or ascending.*

Paraphyllia none.

Capsule smooth when dry.

Leaves obtuse.

Olive or grayish green, 1-2-pinnate, leaves open *H. Schreberi*, 404.

Dirty green to dark brown, almost simple, leaves

closely appressed *H. trifarium*, 405.

Leaves abruptly apiculate, plants pale green *H. purum*, 594.

Capsule plicate when dry, plants dark green to reddish

brown *H. scorpioides*,² 406.

Capsule unknown; plants dark yellow and greenish, branches

julaceous, few, fastigiate, leaves short apiculate *H. turgescens*, 406.

Paraphyllia present *H. Alaskanum*, 405.

¹ *H. circullifolium* C. M. & Kindb., Appendix, no. 583, belonging to this division and described from sterile specimens, is related to *H. dilatatum* Wils.

² See Appendix, no. 599.

II. *Leaves secund.*A. *Costa single, reaching to the middle or beyond.* (B on p. 248.)

1. *Leaves transversely rugose and longitudinally plicate* *H. aduncum gracillescens*, 381.

2. *Leaves not rugose, often plicate.** *Paraphyllia abundant (rarely few).*

Leaves plicate *H. commutatum*, 387.

Leaves not plicate.

Smooth *H. filicinum*,¹ 386.

Very papillæ *H. decipiens*, 561.

* * *Paraphyllia none.*²+ *Annulus none.*

Leaves quite entire, short acuminate *H. palustre*, 398.

Leaves denticulate, subulate acuminate *H. fluitans*,³ 393.

+ + *Annulus present, often large.*

Leaves acute or bluntish *H. ochraceum*, 401.

Leaves subulate by the long excurrent costa *H. capillifolium*, 551.

Leaves gradually long acuminate, costa often entering point but not long excurrent.

Broad (1-2 mm.), crumpled and plicate when dry *H. lycopodioides*,⁴ 385.

Narrower (0.5-1 mm.), not crumpled when dry.

Auricles none or indistinct.

Leaves usually plicate and denticulate.

Capsule oblique, incurved, subcylindric *H. uncinatum*, 392.

Capsule symmetric, erect, cylindrical *H. uncinatum symmetricum*, 552.

Leaves entire.

Plicate, plants pale green or shining yellow *H. vernicosum*, 385.

Smooth, plants brown or purple or blackish *H. revolvens*, 384.

¹ See Appendix, no. 560.

² *H. constatum* C. M. & Kindb., Appendix, no. 558, belonging to this division, is not sufficiently described to enable me to determine its position.

³ Including *H. ezannulatum* Gnemb., sec. Renauld in Hensot's Muscol. Gall. 379. See also Appendix, nos. 553-557.

⁴ Including *H. Wilsoni* Seb. (*H. Sendtneri* Wilsoni Seb.) as a sub-species, sec. Renauld, *ibid.* 375.

Auricles distinct.

Leaves usually denticulate *H. uncinatum*, 382.

Leaves entire or obtusely sinuate.

Falciform, costa 80-120 μ broad, capsule cylindric *H. Sendtneri*, 381.Various, costa 30-60 μ broad, capsule oblong *H. aduncum*,¹ 380.B. *Costa double and short, or none.*1. *Plants regularly pinnate.* (2 on p. 249.)* *Capsule costate and arcuate when dry.*Alar cells short, yellow, thick walled *H. curvifolium*, 396.Alar cells inflated, hyaline, thin walled *H. Patientiae*,² 372.* * *Capsule not costate when dry.*+ *Leaves quite entire.*[*H. imponens* and *H. subimponens* may be sought here.]Alar cells dilated, hyaline or yellowish.³Cilia 2, nodulose *H. Dieckii*, 577.Cilia 3 *H. callichroum*, 393.

Alar cells not dilated.

Cells uniform throughout leaf *H. Watsoni*, 386.Alar cells subquadrate, rest elongated *H. complexum*, 396.+ + *Leaves serrate or denticulate all around.* (+ + +)Capsule long cylindric, suberect or slightly incurved *H. imponens*, 393.

Capsule ovate, oblong or obovate, inclined or arcuate.

Paraphyllia subulate, cilia short *H. Canadense*, 562.Paraphyllia ovate-lanceolate, cilia equaling the segments *H. molliuscum*, 389.+ + + *Leaves serrate or denticulate only above the middle.*Capsule arcuate or incurved cernuous, stem leaves plicate, plants large (to 15 cm.) *H. Crista-castrensis*, 389.

Capsule incurved cernuous, stem leaves not plicate.

Plants large (to 10 cm.) *H. cypressiforme tectorum*, 395.¹ See Appendix, nos. 546-550.² See Appendix, nos. 573-575.³ *H. Renaultii* Kindh., Appendix, no. 571, falls here, but characters given do not allow it to be separated in the key.

Plants small (usually less than 5 cm.).

Perichætical leaves plicate.

Inner perichætical leaves costate.

Alar cells not distinct¹ *H. fertile*, 391.

Alar cells large, hyaline *H. Waghornel*, 565.

Inner perichætical leaves ecostate *H. hamulosum*, 391.

Perichætical leaves not plicate.

Alar cells not enlarged *H. subimponens*, 393.

Alar cells quadrate or vesicular, enlarged.

Leaves narrowly acuminate, faintly bicostate *H. depressulum*, 391.

Leaves long subulate, ecostate *H. circinale*, 392.

2. *Plants irregularly branched.*

[*H. callichroum* may be sought here.]

* *Leaves quite entire.*

Alar cells not different *H. pseudoarcticum*, 584.

Alar cells enlarged, oblong rectangular.

Teeth hyaline margined, cilia appendiculate *H. arcuatiforme*, 576.

Teeth not hyaline margined, cilia nodose *H. pratense*,² 397.

Alar cells quadrate.

Plants erect, fastigiately branched *H. Bambergeri*, 397.

Plants prostrate, irregularly pinnate *H. incurvatum*, 600.

** *Leaves serrulate or denticulate above.*

[*H. pseudo-arcticum* and *H. incurvatum* may be sought here.]

Perichætical leaves costate, and plicate or sulcate.

Costa geminate, reaching middle *H. reptile*, 390.

Costa geminate, short.

Middle leaf cells long (1:12-15) *H. cupressiforme*,³ 394.

Middle leaf cells shorter (1:6-8) *H. Vaucheri*,⁴ 570.

Costa single, or wanting.

Inner perichætical leaves long subulate (1:12) *H. Moseri*, 559.

Inner perichætical leaves lanceolate (1:5) *H. fastigiatum*, 564.

¹ "Blattflügelsellen sehr gross, aufgeblasen, wasserhell bis gold gelb," sec. Mildt, Bry. Silles. 369. Not according to Bryol. Eur. pl. 591, from which this character is taken.

² See Appendix, no. 573.

³ See Appendix, no. 569.

⁴ The character of the perichætical leaves is unknown to me, but since the species is so closely related to *H. cupressiforme*, it is placed in this position.

Perichaetial leaves ecostate.¹

- Plicate *H. plicatille*, 394.
 Not plicate.
 Foliage leaves strongly revolute *H. revolutum*,² 506.
 Foliage leaves concave *H. Sequoieti*, 392.

HYLOCOMIUM,³ p. 409.

Leaves secund.

- Sharply serrate at apex, alar cells quadrate *H. rugosum*, 388.
 Subserrate at apex, alar cells scarcely different *H. robustum*, 388.

Leaves equally spreading.

Paraphyllia none.

Leaves sulcate.

- Ecostate, leaf cells all alike *H. loreum*, 410.
 Bicostrate, leaf cells enlarged at the base *H. triquetrum*,⁴ 409.
 Leaves not sulcate *H. squarrosum*,⁵ 409.

Paraphyllia present.

- Leaves with a long double costa and deeply sulcate *H. umbratum*, 407.
 Leaves obscurely bicostrate.

Paraphyllia pinnate, branches 2-3 pinnate *H. splendens*, 407.

Paraphyllia minute, branching irregularly pinnate *H. brevirostre*, 408.

Leaves unicostrate to middle, coarsely serrate *H. Oakesii*, 408.

¹ *H. pseudo-fastigiatum* C. M. & Kindb., Appendix, no. 563, and *H. pseudo-pratense* Kindb., Appendix no. 578, both described from sterile specimens, fall here.

² See also Appendix, no. 567.

³ Including *Pleurostium* L. & J. Man. 407.

⁴ See Appendix, no. 603.

⁵ See Appendix, no. 602. *H. catvescens* (Wils.) Lindb. is closely related to *H. squarrosum* and is ranked as a variety by Husnot: Muscol. Gall. 425.

BIN.

H. plicatile, 394.

H. revolutum,³ 566.

H. Sequoietl, 392.

H. rugosum, 388.

H. robustum, 388.

H. loreum, 410.

H. luetrum,⁴ 409.

H. marrosum,⁵ 409.

H. umbratum, 407.

H. splendens, 407.

H. revirostre, 408.

H. Oakesii, 408.

H. pseudo-pratense
near.

ated to *H. squar-*

APPENDIX.

DESCRIPTIONS OF SPECIES AND VARIETIES

published since the issue of Lesquereux and James' Manual of the Mosses of North America in 1884, and before January 1, 1896.

1. *Sphagnum Bolanderi* Warnst.—Plants resembling delicate forms of *S. fimbriatum*: stem leaves small, lingulate, mostly delicately fimbriate by resorbed cell membranes; border narrow, equal throughout or broader at base: hyaline cells of upper part abundantly fibrillose, with half-elliptic pores along commissures: branches 3 in a fascicle, two divergent, one pendent: branch leaves very small, ovate lanceolate, narrowly bordered, apex obtuse and toothed, margins involute, densely imbricate and when dry not shining; hyaline cells fibrillose, inner surface in superior half with single large round pores becoming more numerous in marginal regions, outer surface near apex with medium sized round or semi-elliptic pores gradually increasing in size towards base and arranged along commissures; chlorophyllose cells exposed on inner side and not on outside, or only occasionally. Hedwigia 30: 173. 1891.—California.

2. *Sphagnum Russowii* Warnst.—Plants usually tall and strong; tufts loose and high or compact and low: stem leaves large, broad linguliform, with somewhat undulate margins, only in middle of broad rounded apex dentate or somewhat fimbriate; border much widened below; hyaline cells in upper part of leaf large, broad, rhombic, mostly without cross partitions, but with delicate membrane plaits, all with membrane thinnings, which rarely at edges towards apex change into isolated pores, mostly without fibers and pores but rarely fibrillose near apex: fascicles 4 or 5 branched, distant or crowded; 2 or 3 stouter branches spreading, recurved horizontal, curving upward or erect, longer or shorter; pendent branches very long and closely appressed to stem: branch leaves closely or loosely imbricated, mostly with a somewhat spreading (more rarely nearly squarrose) tip, very seldom almost secund, lanceolate, narrow bordered, the upper margins involute, and at the transversely or roundly truncate apex dentate, two or

three plaits near the base: dioicous, rarely monoicous: perigonal leaves resembling the branch leaves; perichaetial leaves as in *S. Girgensohnii*; fruit rare. Bot. Gaz. 15: 130. 1890. Hedwigia 25: 225. 1886.—Newfoundland; Labrador; Canada; New Brunswick; Maine; New Hampshire; Rocky Mountains; Washington.

3. *Sphagnum Warnstorffii* Russ.—Tufts mostly loose; plants usually delicate, slender and graceful, and at the same time firmly erect: stem leaves small to medium sized, mostly linguiform, from base very gradually narrowed and then rather abruptly contracted into a roundish pointed dentate or entire apex; border narrow, much widened downwards; hyaline cells of upper half of leaf rhombic to elongate rhombic, mostly divided but non-fibrillose: fascicle of 3-5 branches of which 2-3 are spreading: leaves of the latter ovate in basal half, involute above and subulate, truncate and 3-5 toothed, often regularly 5 ranked, sometimes secund, always with their points diverging from each other; hyaline cells from basal half of spreading branches with numerous pores on outer surface: dioicous: perichaetial leaves large, ovate-lanceolate, in lower part consisting of chlorophyllose cells only; hyaline cells of upper part non-fibrillose: capsule comparatively large, dark reddish brown. Bot. Gaz. 15: 138. 1890.—Damp or wet birch swamps, margins of elevated bogs when adjacent to birch-covered wet meadows, or in springy swamps: Newfoundland; Labrador; Massachusetts; New Hampshire; Connecticut; Minnesota; Montana; Rocky Mountains; Alaska.

4. *Sphagnum Vancouveriense* Warnst.—Stem leaves without fibrils and pores, rounded apex suddenly narrowed to a short often obtuse and toothed acumen, border up to 10 cells broad and very much broadened toward base; membrane of hyaline cells sometimes very thin or (particularly in upper part) resorbed on both sides: branch leaves when dry curved, erect spreading, on inside in region of margins with many large round pores, on outer face with medium sized to large pores along commissures. Hedwigia 33: 308. 1894.—Vancouver Island.

5. *Sphagnum tenellum* Klingg.—Tufts soft; plants generally quite slender: stem leaves larger or smaller, linguiform, usually cucullate incurved at apex, and sometimes at sides, and afterwards by spreading out flat becoming lacerate dentate or delicately fimbriate; hyaline cells with or without fibrils in upper part of leaf: fascicles of 3-4 branches, 2 diverging: branch leaves loosely or densely imbricate, frequently secund, ovate to ovate-lanceolate, small, dentate at broad rounded apex, margin involute, apical half of inner surface with numerous small pores, especially in upper and lower cell angles, and larger ones in broader part of leaf, especially near margins, outer surface of leaf very porose, pores strongly ringed near apex: dioicous, rarely monoicous: perichaetial leaves large, ovate, above abruptly contracted to a narrow truncate emarginate involute point; hyaline cells

without fibrils or pores. Bot. Gaz. 15: 135. 1890.—In elevated bogs: N. E. United States; Newfoundland; eastern Canada.

6. *Sphagnum fuscum* Klingg.—In extensive, dense or loose, often cushion-shaped patches; stems usually slender and delicate: stem leaves usually small, linguiform, often at rounded apex abruptly contracted to a small cucullate point, which is generally somewhat fimbriate, border broader below; hyaline cells nearly always without fibrils and pores, 2-4-divided by obliquely transverse walls, and with delicate longitudinal plaits in membrane: fascicles of 3-4 branches: leaves small, nearly lusterless when dry, densely or loosely imbricated, from an ovate base extending to a comparatively short round-truncate dentate involute tip; border 3-4 cells wide; hyaline cells on inner side of leaf in upper part with numerous usually ringless pores, especially in upper and lower cell angles, pores in middle of cell, near margin and base, outer surface with numerous pores: dioicous: perichaetial leaves large, ovate, slightly emarginate at rounded apex, hyaline cells without fibrils and pores: fruit rare. Bot. Gaz. 15: 133. 1890.—In elevated bogs: N. United States; Newfoundland; Canada.

7. *Sphagnum quinquefarium* Warnst.—Plants slender and delicate like *S. Warnstorfi* or strong and robust like *S. Russowii*: stem leaves from a broad base deltoid, not linguiform, above at often abruptly contracted apex truncate, dentate, usually involute and bordered at edge, border wider below; hyaline cells in whole middle part of leaf broad, in upper half almost rhombic, often once or several times divided by oblique transverse walls and with membrane plaits, without fibrils and pores oftener than with them: fascicles of 5 branches, 3 spreading; branches 5-angled by 5-ranked leaves: branch leaves loosely or closely imbricate, never secund, small, ovate-lanceolate, at usually round-truncate involute-edged and dentate apex bordered by 2-3 rows of narrow cells, upper part of inner surface porose, pores small, ringed, in upper and lower cell angles, in broader part of leaf and near margins pores larger and not ringed, outer surface with numerous pores along commissures: commonly monoicous, more rarely dioicous: perichaetial leaves large, ovate, above drawn out to a longer or shorter emarginate involute-edged apex: fruit not rare: spores fine, smooth. Bot. Gaz. 15: 189. 1890.—Newfoundland; Canada; E. United States.

8. *Sphagnum tenerum* (Aust.) Warnst.—Stem leaves large, isosceles-triangular, mostly with cut edges and a quite long obtuse toothed acumen with incurved edges; border suddenly broadened at base; hyaline cells often divided by oblique walls and mostly fibrillose to base, inner side porose, pores large, round, outer side with half elliptic pores along commissures: fascicles generally 4-branched: branch leaves quite large, ovate-lanceolate, regularly imbricate; apex obtuse, toothed, margins involute; hyaline cells on inside with large round pores near sides and small pores

near apex in upper and lower cell angles, pores of outside half-elliptic, in rows on commissures. Hedwigia 29: 194. 1890.—New Jersey; Connecticut.

9. *Sphagnum subnitens* Russ. & Warnst.—Plants when dry very soft and with more or less of a metallic luster: stem leaves large, elongated, isosceles-triangular, broad at base, not rarely with undulate margins in middle, above abruptly narrowed into a longer or shorter broad-truncate dentate and involute-edged point; border broad, much widened downwards and formed of very narrow pitted tubular cells; hyaline cells in middle of base wide and large, above rather shorter, rhomboidal, at margins narrow, mostly without fibrils and pores, rarely with rudiments of fibrils and pores at apex, longitudinally plaited: branches 3-4 in a fascicle, one or two pendent: leaves of spreading branches larger or smaller, densely or loosely imbricate, often curved, erect-spreading, seldom somewhat secund or squarrose, never distinctly 5-ranked, from an ovate base narrowed upward to a rather long dentate transversely or roundly-truncate involute-edged apex; hyaline cells with numerous fibrils, pores on inner surface almost all near margins, on outer surface more numerous: mostly monocious, more rarely dioicous: perichæatial leaves large, ovate, edge very broadly bordered in upper part and emarginate at rounded truncate apex: fruit very common. Bot. Gaz. 15: 194. 1890.—Newfoundland; Labrador; Miquelon Is.; Nova Scotia; Maine; New Hampshire; Massachusetts; New Jersey; Connecticut; Virginia; Indiana; California.

10. *Sphagnum microphyllum* Warnst.—Plants very delicate, very similar to small slender forms of *S. fimbriatum* Wils.: stem leaves large, about $2\frac{1}{2}$ times as long as broad, above narrowed into a distinctly truncate toothed apex; border narrow and broadened slightly below; hyaline cells elongated-rhomboidal, without cross walls in apical half, in upper $\frac{2}{3}$ to $\frac{3}{4}$ abundantly fibrillose, on inner side with many large round ringless pores in middle of cell, on outside with half-elliptic pores along commissures: fascicles 3-branched, 1 pendent: leaves of spreading branches small, ovate lanceolate, densely imbricate or with apex erect-spreading or almost squarrose, apex obtuse, toothed, in upper part margin involute, narrowly bordered, when dry not shining; hyaline cells fibrillose, abundantly porose on inside, pores round, ringless, outside with half-elliptic ringed pores in rows along commissures. Hedwigia 80: 172. 1891.—California.

11. *Sphagnum Labradorensis* Warnst.—Stem leaves medium sized, linguulate spatulate from a narrowed base, border narrow, broadened at base, margin broadly involute on one or both sides, finely toothed at obtuse apex; hyaline cells broadly rhombic, near base narrower and longer, divided with one to four oblique walls, inner surface with large pores, outer surface with very thin membranes, resorbed in apical part, without fibrils or in apical part just the beginnings: fascicle of 3-4 branches, two spreading: branch leaves ovate, about as large as stem leaves, apex broad, obtuse,

with 4-6 large teeth; margin faintly bordered, toothed above, involute far down, loosely erect spreading; hyaline cells broad rhombic to rhomboidal, inner surface near apex with small faintly ringed pores, towards middle in side regions with few large pores, outer surface with numerous narrow elliptic pores gradually larger towards the leaf base and margins, fibrils very numerous. *Hedwigia* 81: 174. 1892.—Newfoundland; New Jersey.

12. *Sphagnum Floridanum* Card.—Differs from *S. macrophyllum* by more numerous pores of stem leaves (4-10 in each cell); shorter pointed branch leaves, rounded obtuse, not tubulose, canaliculate and cucullate at summit; hyaline cells narrower, pores small and very numerous, 40-60 upon each face in large cells of middle, biseriate, entirely at one end of the cell, or uniseriate; fruit unknown. *Cardot, Rév. des Sphaignes de l'Amérique du Nord* 22. 1887.—Florida; Louisiana.

13. *Sphagnum riparium* Aongstr.—Cortex of stem wanting; stem leaves very large, reflexed, triangular lingulate, apex rounded, deeply two cleft by resorption of membranes, always without fibrils, borders entirely of green cells with pits: fascicle of 4-5 branches: leaves narrowly bordered, when dry somewhat crispate, middle ones with squarrosely reflexed points, chlorophyllose cells exposed on both faces, pores on concave leaf surface quite large and numerous along the commissures, membranes resorbed at upper end of cells on convex surface for entire breadth of leaf in middle or side regions, in other cases with 2-4 large pores instead: perichetial leaves with the lower $\frac{2}{3}$ composed mostly of pitted chlorophyllose cells. *Limpr. Laubm.* 1: 133.—New Hampshire; New Jersey; Canada; Greenland; Alaska; Behring Sea.

14. *Sphagnum Dusenii* Jensen.—Strong to robust, green or yellowish-green: wood body green or yellow, formed of quite thick-walled cells; cortex of 5 layers of cells with weakly thickened walls: stem leaves triangular lingulate, apex with few large teeth, generally fibrous in upper half, border broad: branches 4 in each fascicle, 2 divergent often long and attenuate: leaves crowded or loosely imbricate, seldom secund, large (2×1 mm.). broad oval lanceolate, above with incurved margin; hyaline cells long and narrow, below $15-20 \times 2.5-3\mu$, strongly fibrous, on convex side with numerous round or oval pores ($5-7\mu$ diam.) with weakly thickened edges, on concave side with single rather distant round pores along sides of broad part, all pores somewhat distant from chlorophyll cells, often forming 1 or 2 rows: dioicous; σ branches acute, with yellowish brown bracts, ρ branches short or elongated, their leaves broadly oval with hyaline cells at base and fibrous in upper half: capsule brown; spores 2μ diam., yellow and finely papillose: seldom fruiting. *S. majus* Russow; Jensen in *De danske Sphagnum-Arter, Festskrift bot. Foren. Copenh.* 50-aarsfest 106. 1890. Anticosti; Maine; New Hampshire; New York; Wisconsin.

15. *Sphagnum Mohrianum* Warnst.—Stem leaves large, narrow at

base, broadening towards middle and then narrowing into an obtuse fine toothed point with involute margins, accordingly nearly lanceolate; border narrow; hyaline cells mostly divided by an oblique wall, abundantly fibrillose to base, inner surface of basal half with large round ringless pores in middle of cell wall, basal cells with a very large opening, in upper half with single pores in the cell angles: fascicle 3-branched: branch leaves ovate to oblong-ovate, apex broadly obtuse and toothed, narrowly bordered margins broadly involute; hyaline cells on inner surface abundantly fibrillose, sparingly porose on both sides, only with single pores in cell angles. Hedwigia 31: 179. 1892.—Mobile, Alabama.

16. *Sphagnum obesum* Wils.—Stem leaves large, reflexed, oblong triangular, apex cucullate or simply obtuse, fringed, border narrow, without fibrils below: branches swollen, curving downward, never twisted: leaves soft and loosely placed, in water somewhat pinnately spreading, broad to narrow ovate-lanceolate, broader bordered, apex several toothed, both sides of apical half with small pores in almost all cell angles, especially in upper and lower, rarely almost wanting. *Sphagnum contortum obesum* (Wils.) Limpr. Limpr. Laubm. 1: 121.—New Hampshire; Massachusetts; Virginia; Connecticut.

17. *Sphagnum dasyphyllum* Warnst.—Stem leaves quite large, linguulate, concave, border narrow, involute for greater part, apex cucullate, split when spread out; hyaline cells below narrow and long, towards apex rhomboidal, all divided by 1 or 2 oblique walls, upper $\frac{3}{4}$ abundantly fibrillose, then a space without fibrils and again fibrillose at base; inner surface rarely porose, outer surface with pores at apex only: fascicles of 2-3 spreading and 2 pendent branches: branch leaves quite large, base narrow and greatly broadened towards middle, then suddenly narrowed into a short obtuse finely toothed point, accordingly round-ovate, very concave, border narrow and involute to base; hyaline cells broad rhomboidal, upper half of inner surface with single large pseudo-pores, outer surface with more numerous pseudo-pores and single very small strongly ringed pores in the cell angles. Hedwigia 31: 176. 1892.—New Haven, Connecticut.

18. *Sphagnum microcarpum* Warnst.—Stem leaves crowded, very large, broad ovate, apex broadly rounded obtuse, finely toothed; margin broadly involute and narrowly and equally bordered to base; hyaline cells narrow, rhomboidal, abundantly fibrillose to base; pores almost wanting on inner surface, outer surface with very numerous small pores in uninterrupted chains along commissures, decreasing in number towards base: branches single, not fasciculate: branch leaves loosely placed, quite large, ovate to oblong-oval, border narrow, involute for greater part; areolation very similiar to that of stem leaves. Hedwigia 30: 170. 1891. New Jersey; Florida; Alabama; Mississippi; Louisiana.

19. *Sphagnum platyphyllum* Sulliv.—Stems lax, quite robust, irregu-

larly branched, sometimes without branches; branches 1-3 in a fascicle, mostly short and thick: stem and branch leaves very similar in form, stem leaves large, oval to obovate, very concave, apex rounded, slightly fringed, below narrowly bordered, fibrillose to base but with very small pores along commissures only in upper half; branch leaves appressed, loosely placed, large, very concave, ovate, rounded above, scarcely toothed, narrowly bordered all around, strongly fibrillose to base, upper $\frac{3}{4}$ of outer surface with very small pores along commissures. *Limpr. Laubm.* 1: 122. 1890.—Massachusetts; New Jersey; Virginia.

20. *Sphagnum plicatum* Warnst.—Stem leaves quite large, lingulate, upper margins frequently involute, apex hyaline margined, bordered, border weak, of 4-6 rows of cells, equally broad to base; hyaline cells in apical half about rhombic, in basal portion broader and longer, frequently divided by one or two oblique walls, abundantly fibrillose in upper $\frac{3}{4}$; pores on inner surface in upper part of leaf in all cell angles, towards margins more numerous; outer surface with pores in often interrupted rows along commissures, towards base almost exclusively in upper cell angles: fascicles mostly 4-branched, 2 spreading: leaves of spreading branches quite large, ovate-lanceolate, coarsely toothed at narrowly obtuse apex, border narrow, involute at apex only or sometimes farther down; hyaline cells broad, rhomboidal, not divided, pores on inner surface only in side regions, outer surface with numerous pores in rows along commissures. *Hedwigia* 80: 169. 1891.—Granville, Mass.

21. *Sphagnum Orlandense* Warnst.—Stem leaves relatively small, lingulate, apex rounded, scarcely toothed, not cucullate, border narrow, involute on one side nearly to base; hyaline cells generally divided by an oblique wall, fibrillose to base, outer surface with small faintly ringed pores in interrupted rows along commissures: fascicles mostly four-branched, 2 spreading: leaves of spreading branches quite large, roundish-oval, finely toothed at obtuse apex, very concave, border narrow, margins broadly involute; hyaline cells broad rhomboidal, abundantly fibrillose, outer surface towards apex with pores in cell angles, in middle side regions with small pores in rows, outer surface with numerous pores in interrupted rows along commissures. *Hedwigia* 81: 177. 1892.—Florida; New Jersey.

22. *Sphagnum Mobileense* Warnst.—Stem leaves relatively small, lingulate, narrowly bordered to base, in upper half one margin sometimes involute; hyaline cells of lower part of leaf not divided or with a single oblique wall, without fibrils or only basal cells with few delicate fibrils, in upper part with parallel oblique cross walls and abundantly fibrillose and porose, towards apex chlorophyll cells ceasing and hyaline cells forming a broad border which is sometimes resorbed: fascicles 4-5-branched, 2 spreading: leaves of spreading branches large, round-ovate, finely toothed on scarcely obtuse apex, narrow bordered, very concave, margins broadly

involute; hyaline cells abundantly fibrillose, not divided or somewhat divided towards margins, inner surface with small pores in almost all cell angles particularly in upper half, outer surface with pores in rows along commissures, less numerous towards base. Hedwigia 31: 190. 1892.—Mobile, Alabama.

23. *Sphagnum simile* Warnst.—Stem leaves small, lingulate; hyaline cells at apex only occasionally divided by a cross wall, hyaline border of apex partly resorbed and fimbriate, upper part of leaf fibrillose; branch leaves oval to oblong-ovate, both sides abundantly porose, inner surface with small pores in almost all cell angles, outer surface with pores along commissures; chlorophyll cells in cross section parallel-trapezoidal and free on both faces. Hedwigia 33: 326. 1894.—Madison, Wisconsin.

23a. *Sphagnum Waghornel* Warnst.—Chlorophyll cells of the branch leaves in cross section broadly isosceles-trapezoidal, the longer side on the inner face, exposed on both faces, walls uniformly thickened; hyaline cells smooth within where they adjoin green cells: wood body dark brown to almost black: cortical cells with spiral fibrils and outer wall with 1 (rarely 2) large pores: otherwise as in all *CYMBIFOLIA*. Hedwigia 33: 329. 1894.—Newfoundland.

24. *Sphagnum rufescens* Bryol. Germ.—Seldom completely submersed: stem leaves large, triangular-linguiform, lateral margins narrowly and uniformly bordered down to base; hyaline cells fibrillose from apex far downward, often quite to base, and with small pores on both sides, on inner side especially in cell angles, on outer side mostly in interrupted rows on commissures: branch leaves large to very large, ovate-lanceolate with numerous small pores on inner side, sometimes in rows near margins, still more numerous on outer side, in rows, like strings of pearls, on commissures; rings strong. Bot. Gaz. 15: 246. 1890. Hedwigia 33: 326. 1894.—From Newfoundland, Labrador and Canada to Connecticut; also Washington and California.

24a. *Sphagnum medium* Limpr.—Stem leaves as in *S. cymbifolium*, larger, plicate, rarely in upper half with a few fibrils and few large pores; branch leaves involute all around, pores more numerous on the outer surface, especially in cell angles: dioicous. Laubm. 1: 104. 1890.—From Newfoundland and Labrador through Canada to Florida.

25. *Sphagnum Ludoviclanum* (Ren. & Card.) Warnst.—Stem leaves very large, broad-lingulate, apex cucullate and the margins there involute, narrowly bordered all around; fibrillose to base, inner surface with pseudo-pores where three cell angles meet, near margin with large or small pores, outer surface with narrow elliptic pores along commissures, larger below; or stem leaves narrow at base and broadened to middle, then narrowed into a broad rounded cucullate apex with involute margins: fascicles 3-4-branched 2 spreading: branch leaves large, ovate, apex hyaline

bordered, cucullate, margins more or less involute; hyaline cells abundantly fibrillose, pores few on inner surface, on outer surface more numerous especially towards apex. Hedwigia 80: 161. 1891.—New Jersey; Florida; Mississippi; Louisiana.

26. *Andreaea parvifolia* Müll.—Dioicous: small, slender, filiform, sparingly dichotomous; stem leaves erect-imbricate or slightly secund, minute, when moist from an erect deeply ventricose oval base extended into a spreading oblong more or less roundish-obtuse and recurved more deeply ventricose lamina; ecostate; cells angular, thick walled, fuscous, strongly hyaline-papillose; perichætal leaves convolute into a very narrow short cylinder, larger, broader, rather broadly oblong from a narrow base, ligulate-acuminate, somewhat obtuse, strongly papillose; capsule short pedicellate, minute. Flora 70: 219. 1887.—Alaska.

27. *Andreaea papillosa* Lindb.—Plants elongate (1-2 in. high) rather stout, branched, slightly tufted, dark brown to black; leaves crowded, spreading in all directions, somewhat shining, ovate with short narrow acute point, cells large (3 times as large as in *A. petrophila*¹ while leaves are as short as in that species), coarsely papillose. Hartman, Skand. Flora 122. 1871.

28. *Andreaea alpestris* Sch.—Monoicous: leaves much smaller than in *A. petrophila*, crowded from the sheathing base when moist, ovate, oblique, obtuse, margin hyaline, faintly papillose on the back, ecostate; perichætal leaves with long papillæ, cells smaller than in *A. petrophila*, round-hexagonal to oval in upper part of leaf, basal cells elongated rectangular, pits few or absent. Limpr. Laubm. 1: 142.—On damp rocks: Nova Scotia, near Halifax; Greenland.

29. *Andreaea obovata* Thed.—Densely pulvinate-caespitose, tufts blackish above, fuscous below: plants tall, robust, much branched, branches fastigate; leaves densely crowded, spreading from the imbricate base, apex inclined upward, when dry closely imbricate; from ovate-panduriform gradually lanceolate, rather obtuse; minutely papillose or almost smooth; shining, margin very entire; blackish, under a lens golden yellow; cells very thick walled, rhombic above, sinuate-elongate below, the cell cavities really roundish above, rectangular-hexagonal below: male fls. numerous, on special slender branches; antheridia large, paraphyses long; perichætal leaves surpassing the upper cauline leaves, convolute, the outer broadly ovate-lanceolate, internal broadly elongate; apex obtuse, not rarely hyaline or premorse. Schimper Syn. Musc. Eur. 814. 1876.—Disco Is., Greenland.

30. *Andreaea Huntii* Limpr.—Monoicous: tufts dark, somewhat glossy; leaves strongly falcate-secund, short, broadly ovate, strongly concave base gradually narrowing into a long subulate point; base of lamina made up

¹ *A. petrophila* has cells 10-14-17 μ in diameter sec. Limpricht, Die Laubmoose 1: 140.

of one layer of cells, upper part of leaf two-layered; costa strong, about one-third width leaf; leaf cells in the lower part round-quadrate, at the base near costa rectangular: conial leaves shorter with the costa often disappearing below apex, margin sometimes irregularly toothed: inner perigonal and perichaetial leaves ecostate, margin erenulate, convex surface strongly papillose. *Limpr. Laubm.* 1: 145. On rocks, altitude 5,500 ft.—Vancouver Island.

31. *Andreaea Blyttii* Schimp.—Very small slender and densely pulvinate, brownish black to deep black: stem very slender, rather rigid, brittle, fastigiate-branched, rooting: leaves divaricate-homomallous, those near the tips falcate-secund, from an oval or ovate base suddenly subulate, nearly smooth, shining, brittle; costa semi-terete, occupying almost all the subula, terete at apex; cells near apex round, at base all rectangular: ♂ plants gregarious, more slender than ♀, perigonal lvs. 6, the 3 inner united into a bud, acuminate, ecostate; antheridia 5-8, long pedicellate, with longer filiform flexuous paraphyses: perichætium large, far exceeding stem leaves, external leaves erect, costate to apex, internal subconvolute, elongate-obovate, short apiculate, ecostate: capsule very small, conc-ovate. *Sch. Syn. Musc. Eur.* 821. 1876.—Greenland.

32. *Andreaea Macounii* Kindb.—Dioicous: tufts soft, dark brown or brick red, not glossy: stems about 5 cm. high: leaves narrow, suddenly long acuminate from broader base, distinctly but faintly papillose, more or less curved, uppermost falcate, when dry sub-crispate; cells uniform sub-quadrate; costa narrow, linear, nearly smooth, in the upper leaves long excurrent and denticulate, sometimes to middle. *Bull. Torr. Bot. Club* 17: 82. 1890.—In cushions and flat mats on inclined faces of rocks on mountains and by brooks flowing from perpetual snow, Gold Range, B. C.

33. *Andreaea nivalis* Hook.—Dioicous: tufts broad, thick, soft, dirty greenish brown, reddish brown or black, not glossy: stem 4-10 cm. long: leaves more or less falcate, crispate when dry, uniformly elongated into a sharp point from a somewhat enlarged or auricled base; lamina 1-layered, papillose on both surfaces, irregularly toothed near apex; costa quite strong, uniform in width even to apex; leaf cells quadrate or short rectangular: perigonal leaves ecostate, broadly oval, suddenly long pointed; perichaetial leaves quite similar to stem leaves. *Limpr. Laubm.* 1: 152.—Mt. Hood, Oregon.

34. *Archidium Hallii* minns R. & C.—Smaller, leaves shorter, costa percurrent or short excurrent. *Bot. Gaz.* 19: 237. 1894.—Louisiana.

35. *Phascum subexsertum* Hook.¹—Stemless, cespitose: leaves linguulate oblong, spreading, apiculate: capsule spherical, sub-exserted, lid acuminate. *Mac. Cat.* 12.—Northwest Territory.

¹This species is not sufficiently known (Ren. & Card). Wilson states that this is perhaps only a remarkable variety of *P. cuspidatum*.—*Hook. Journal Bot.* 3: 433. 1841.

costa strong, about
d-quadrata, at the
the costa often dis-
oothed: inner peri-
ate, convex surface
altitude 5,500 ft.—

and densely pulvi-
rather rigid, brittle,
allous, those near
suddenly subulate,
pying almost all the
se all rectangular:
lvs. 6, the 3 inner
3, long pedicellate,
large, far exceeding
rnal subconvolute,
small, conic-ovate.

ft, dark brown or
narrow, suddenly
y papillose, more or
cells uniform sub-
upper leaves long
l. Torr. Bot. Club
aces of rocks on
Gold Range, B. C.
, thick, soft, dirty
em 4-10 cm. long:
elongated into a
lamina 1-layered,
apex; costa quite
te or short rectan-
long pointed; peri-
lbm. 1: 152.—Mt.

ves shorter, costa
—Louisiana.
ose: leaves lingu-
sub-exserted, lid

ates that this is per-
Bot. 3: 433. 1841.

36. *Pleuridium alternifolium* Howe Ren. & Card.—Differs from the typical form in the leaves being entire or slightly denticulate at the apex. *Revue. Bryol.* 19: 64. 1892.—California.

37. *Microbryum Floerkeanum* Henrici R. & C.—Differs from the typical form in the green color of the plant, and the excurrent costa often hyaline at the point. *Bot. Gaz.* 14: 91. 1889.—Sandy ground: Saline Co., Kansas.

38. *Bruchia longicollis* Eaton.—Plants densely clustered, 7-10 mm. high: stem leaves with a broadly ovate clasping base suddenly narrowed a into long excurrent awn like costa, which is bordered below by leaf margin: perichetial leaves lanceolate, somewhat tubulose, gradually acuminate, costa excurrent: flowers monoicous: capsule exserted on a stout seta, orange yellow, collum very long, exceeding sporangium, stomatose beak slender. *Bull. Torr. Bot. Club* 17: 100. *pt. 101.* 1890.—Decayed logs in swamp: New Hampshire.

39. *Bruchia fusca* Britt.—Plants gregarious, 2-3 mm. high: leaves few, three to six, erect appressed, short, clasping, often broader than long and tricuspidate, entire or subserrulate, with a narrow border of small retuse cells, apex obtuse, acute or cucullate; costa faint, vanishing below apex or absent in lower leaves, basal cells lax: seta immersed or slightly exserted, straight or curved; capsule large and broad, 1-1.5 mm. long, entirely exserted, ovoid-pyriform, suddenly apiculate; collum shorter than but occasionally equaling the sporangium; calyptra smooth, deeply lobed, half covering the capsule; spores small, brown, angled and pitted. *Bull. Torr. Bot. Club* 21: 361. 1894.—Growing around quartz pebbles in sandy soil: Maryland; North Carolina.

40. *Bruchia Carolinæ* Aust.—Plants gregarious, in brown patches, 1-2 mm. high: stems naked and radiculose at base, leaves crowded at the summit, more or less secund, subulate from a broader base; costa channeled, filling the entire or serrulate apex, faintly papillose on the back; basal cells smooth, irregular, upper with thickened walls: seta shorter than the capsule, both immersed, or the capsule occasionally exserted laterally, pyriform, yellow or brown, conic apiculate; collum large, truncate, stomata immersed; calyptra broad lobed, papillose at the apex; spores small, pitted: flowers monoicous. *Bull. Torr. Bot. Club* 21: 365. 1894. *Bruchia Ravenelii* Wils., var. *mollis*, L. & J. Man. 49.—South Carolina.

41. *Astomum Drummondii* Kindb.—Plants cespitose, almost stemless: leaves linear subulate, very entire: seta as long as leaves: capsule globose, lid conic rostrate. *Mac. Cat.* 12.—Plains of the Saskatchewan.

42. *Voitia* Hech.—Tufts thick, densely cespitose, below brown tomentose: stems simple or dichotomously branched: leaves somewhat separated, or densely imbricate, long or short, ovate or elongated elliptical, cuspidate; costa thin and weak; cells large, thin-walled, hexagonal or sub-quadrata,

hyaline near the base, chlorophyllose above: flowers monoicous: capsule erect, oval or ovate; apiculate lid not deciduous; calyptra large, cucullate, nearly covering the capsule. Musc. Gall. 69.

43. *Vetitia hyperborea* Grev. & Arnott.—Smaller and more slender than the European *V. nivalis*: leaves densely imbricate, erect-appressed, broadly oval or sub-rotund, short acuminate, strongly concave, margin entire; costa vanishing in apex: perichæstrial leaves long cuspidate, costa excurrent: vaginule long, membranaceous: pedicel short; capsule oval-globose, base subangulate, apex slightly curved; spores minute, smooth. Müller Syn. Musc. Frond. 1: 35.—Greenland.

44. *Gymnostomum platyphyllum* Kindb.—Dioicous: plants green, not glossy: stem not papillose, sparingly radiculose: leaves decurrent, distinctly papillose, short, obtuse, oblong, lanceolate, the comal greater, one border often recurved all around; costa greenish yellow, very papillose. Bull. Torr. Bot. Club 17: 84. 1890.—On the face of a small waterfall, near Kamloops, B. C.

45. *Anacetangium compactum* Schw.—Dioicous: stem fragile, red tomentose: leaves erect spreading, when dry appressed, spirally arranged or with the twisted points incurved, linear-lanceolate, margin plane, faintly crenulate; costa strong, percurrent or vanishing below apex; basal cells short rectangular, the remainder round-hexagonal, papillose on both sides: perichæstrial leaves sheathing, outer ones small, inner larger, smooth, costa vanishing far below apex: seta straw-colored, twisted to the right; capsule obovate, smooth, neck short, indistinct; two rows of quadrate annular cells which sometimes remain with the capsule and sometimes with the lid. Limpr. Laubm. 1: 244. fig. 92.—Greenland.

46. *Eucladium* B. & S.—Genus closely related to *Trichostomum* and *Gymnostomum*: plants caespitose, erect, dichotomously branched, base of the old innovations radiculose: leaves erect-spreading, when dry straight or slightly curved, lanceolate or linear-lanceolate, margin plane, toothed near the middle; costa strong, round: dioicous: capsule generally erect, oval or long oval, with a short indistinct neck; lid obliquely rostrate; annulus simple; peristome of sixteen lanceolate or linear-lanceolate teeth, entire, lacerate or perforated; spores small, smooth.—Bryol. Eur.

47. *Eucladium verticillatum* B. & S.—Tufts thick, 1-4 cm. rarely 14 cm. long, bluish-green, below light or yellowish brown: leaves linear-lanceolate; costa percurrent or rarely excurrent; cells of the lower part of leaf thin-walled, hyaline, long rectangular, marginal cells smaller, the remainder richly chlorophyllose, round quadrate and also rectangular, papillose on both sides: perichæstrial leaves scarcely different: calyptra cucullate, covering $\frac{1}{2}$ the capsule. *Mollia verticillata* Lindb., Braith. Brit. Moss Flora 1: 241.—Santa Ana Cañon, California.

48. *Welsia convoluta* C. M. & Kindb.—Tufts dense, dark green: stem

slender, branching, densely foliate: leaves very patent when moist, flexuous, from a broader oblong base narrowed into the very narrow and long often piliform acumen, margins entire and erect; cells small, quadrate, chlorophyllose, opaque; costa very thin, vanishing in the acumen: perichæatial leaves broader, vaginant at base, shorter than the others, densely appressed involving the pedicel: capsule cylindric, very narrow, faintly curved; lid obliquely rostrate; pedicel yellowish, elongate, slender, spirally twisted, narrowed to the capsule. Mac. Cat. 14.—Mountains near Silver City, N. W. T., altitude 7,700 ft.

49. *Welsia viridula nitida* Ren. & Card.—Leaves shorter, capsule narrow, sub-cylindrical, shining as though varnished and distinctly sulcate when dry. Bot. Gaz. 14: 91. 1889.—On sandy ground: Florida; Louisiana.

50. *Welsia viridula rugellana* Ren. & Card.—Calyptra larger, operculum longer rostrate. Revue Bryol. 19: 73. 1892.—Georgia.

51. *Cynodontium subalpestre* Kindb.—Tufts green, about 1 cm. high or lower: leaves crisped, from the narrowly ovate-oblong base attenuate-subulate, acute, nearly smooth as also on the costa, entire or distinctly denticulate above; borders recurved to middle at one side; cells pellucid, mostly quadrate, alar much wider, rectangular, hyaline: perichæatial leaves with a broader base, inner ones much shorter: capsule small, subcylindric-clavate, wide-mouthed, faintly striate when dry; teeth pale, bifid, not papillose; annulus wanting; lid conic, short rostrate; pedicel yellow, finely bright red. Mac. Cat. 17, 257.—Rocks near water: Labrador; Quebec; Lake Huron; Lake Superior; Lake Nepigon.

52. *Cynodontium strumulosum* C. M. & Kindb.—Tufts bright green, 1-2 cm. high: leaves crisped, sub-linear, blunt or sub-acute, recurved at the basal margins, crenulate above, nearly pellucid; basal cells linear, the others short, angular; costa very papillose at the back, not percurrent: perichæatial leaves short acuminate: capsule small, oblique, oblong-ovate, inclined, when dry furrowed, distinctly strumose; teeth not papillose, cleft to the middle, reddish also in the upper part; annulus distinct; pedicel light red or yellow, straight, 1 cm. long. Mac. Cat. 16.—Limestone rock along a torrent near Hector, B. C.

53. *Dichodontia pellucidum fagimontanum* Sch.—Plants shorter, more dense, with shorter branches: leaves shorter, more obtuse, scarcely recurved; capsule smaller, less inclined. Braithw. Brit. Moss Flora 1: 163.—Hector, B. C.

54. *Dichodontium Olympleum* Ren. & Card.—Differs from much more robust *D. pellucidum* thus: leaves strongly papillose, minutely denticulate almost all around: capsule strumose at base: plants delicate, scarcely 1 cm. high. Bot. Gaz. 17: 296. 1892.—Olympic Mts., Wash.

55. *Trematodon brevicollis* Hsch.—Monoicous: tufts small, stems 2-4 cm. high: leaves imbricate, lower smaller and loosely appressed, long and

pointed, upper aggregated, larger, broad ovate, concave, margins plane, netire, suddenly narrowed into a short lance-like point; costa percurrent, broad and flat; cells thin walled, densely chlorophyllose, below rectangular, above rhombic or rhomboidal: perichætal leaves larger, sheathing, short-pointed, costa narrower, vanishing near apex: seta twisted to right when dry; capsule long, neck equaling sporangium, slightly curved, gradually narrowed into seta, when dry faintly costate; calyptra inflated, cucullate; lid $\frac{1}{2}$ as long as capsule, obliquely long rostrate from a broad base; annulus compound, of two rows of cells; teeth lanceolate, undivided, often perforated along middle; spores large, round or oval, papillose. Limpr. Laubm. 1: 416. *fls.* 139, 140.—Greenland.

56. *Dicranella Schreberi lenta* (Wils.) Limpr.—Tufts thick, about 4 cm. high: leaves larger and broader, plainly toothed on margin. Limpr. Laubm. 1: 318.—Moist earth, Washington.

57. *Dicranella parvula* Kindb.—Allied to *D. varia*: dioicous: plants very short, densely cespitose in small tufts, dark green: leaves crenulate all around, sub-ovate; cells short, oblong quadrate, basal linear; costa very thick and brown, excurrent, denticulate above, forming at least half acumen: perichætal leaves denticulate above: capsule obovate, sub-erect or inclined, without a neck, not striate; beak very short and thick; annulus scarcely loosed; teeth orange, pale at apex, partite, without basilar membrane, not papillose; pedicel pale red yellow, 1 cm. long or shorter, straight or curved only at apex. Mac. Cat. 21.—On the earth: Cathedral Mountain, B. C.

58. *Dicranella Langloisii* Ren. & Card.—Cespitose, pale or yellowish green: stems short, 4-7 mm. long: leaves small, crowded, erect spreading when moist, appressed when dry, from an oblong base shortly acuminate, blunt at denticulate apex, strongly revolute on one side, slightly reflexed or almost plane on other; costa stout, broad, percurrent, rounded at back; leaf cells short, rectangular or sub-rectangular below, elongated, 4-7 times longer than broad above: perichætal bracts longer, longer acuminate; costa short excurrent: pedicel purple; capsule suberect, oblong, incurved, reddish-brown, constricted under orifice when dry; lid large, highly convex-conic, with an oblique beak; peristome purple, high, teeth bifid to above middle; annulus none. Bot. Gaz. 15: 39. *pl.* 5. 1890.—On the ground: Saint Martinville, Louisiana.

59. *Dicranella leptotrichoides* Ren. & Card.—Resembling in habit *Leptotrichum tortile*: dioicous, small, loosely cespitose, green: stems short, erect, simple: leaves erect or sub-secund, lanceolate, acuminate, sub-acute or rather obtuse at the apex, quite entire, borders plane or partly revolute; costa stout, percurrent; leaf cells oblong or sub-linear, shorter, looser, and sub-rectangular below: perichætal bracts scarcely different, a little longer: seta long, yellowish when young, afterwards red-

dish; capsule small, erect, symmetric, ovate-oblong, scarcely constricted under the orifice when dry; lid equaling sporangium, long and obliquely subulate rostrate; peristome purple, teeth trabeculate, striolate lengthwise, cleft to below the middle into 2-3 subulate legs, granulose and partly connected; annulus very broad, deciduous, 2-3 rows of cells: male plants unknown. Bot. Gaz. 19: 237. pl. 21. 1894.—On the ground: Louisiana; Mobile, Alabama.

60. *Dicranella Fitzgeraldi* Ren. & Card.—Cespitose; yellowish or dirty green: stems long, simple or bipartite: leaves crowded, sub-secund, from a lanceolate base gradually narrowed into a long subulate canaliculate point, denticulate at the apex; costa broad, occupying $\frac{1}{3}$ of leaf-base and nearly all of point; cells rectangular or sub-rectangular, firm, more or less elongated: perichaetial leaves from a dilated, sinuate-denticulate base abruptly subulate, denticulate at apex: seta yellowish, long, twisted; capsule erect, symmetric, oblong, not constricted under orifice, when dry slightly plicate, brown or yellowish; lid convex, obliquely long rostrate; teeth purple or orange, densely trabeculate, striolate lengthwise, faintly granulose. cleft to below middle into two very long subulate legs, quite free or partly connected: male flowers unknown. Bot. Gaz. 18: 197. pl. 15. 1888.—On sandy ground: Palatka, Florida.

61. *Dicranella Howei* Ren. & Card.—Closely related to *D. varia*: stem longer: leaves more lax, longer, more flexuous, margin plane throughout; costa broader, cells narrower: capsule longer and more slender, shining green. Revue Bryol. 20: 30. 1893.—California.

62. *Dicranella laxiretis* Ren. & Card.—Closely related to *D. debilis*: capsule narrower: leaves narrower and more flexuous, with a denticulate apex; cells larger and shorter. Revue Bryol. 20: 30. 1893.—Louisiana.

63. *Dicranella cerviculatula* Kindb. (Labrador) and *D. polaris* Kindb. (Alaska) are *nomina nuda*.

64. *Dicranum hyperboreum* (Gunn.) Smith.—Resembling *D. fulvulum* but more robust, cespitose: leaves patent and sub-secund, when dry incurved, subflexuous, lanceolate and lanceolate-subulate from the lanceolate base, acumen nearly smooth; areolation dense: perichaetial leaves narrowly aristate from the long sheathing base: capsule erect or subcernuous, ovate or oval, neck inflated, substriate, when dry and empty cyathiform and plicate; teeth of the peristome broader, short subulate, distinctly strigoe, 2-3 divided and perforated to the middle. Mull. Syn. Musc. 1: 372.—On rocks: Mt. Hood, Oregon.

65. *Dicranum hyperboreum papillosum* Ren. & Card.—Leaves shorter, deeply canaliculate, papillose on back; costa strongly rugose. Bot. Gaz. 14: 91. 1889.—Disco, Greenland.

65a. *Dicranum falcatum Hendersoni* Ren. & Card.—Pedicel purple

below, yellow above. Bot. Gaz. 15: 39. 1890.—Moist sunny rocks: Mt. Hood, Oregon.

66. *Dicranum molle* Wils.—Tufts large, dense, 6-15 cm. high, yellow green or olivaceous above, fuscous below: stem eradiculose: leaves erect-patent, straight, oblong lanceolate, subulate, entire, very concave or semi-tubulose above by incurved margin, auricled at base; costa narrow, percurrent or excurrent; alar cells large, orange, quadrate, remainder narrow linear: perichætal leaves oval-oblong, sheathing, suddenly subulate, imperfectly denticulate at apex: capsule oblong-cylindric, cernuous incurved, substrumose, not striate, fuscous; annulus simple; lid short beaked; peristome purple: monoicous, male inflorescence close to perichæcium. Braithw. British Moss Flora 1: 144. pl. 20.—Greenland.

67. *Dicranum longifolium compactum* Ren. and Card.—Plants small, in very compact tufts: stems 1-2 cm. long: leaves half as large as type, erect, straight or slightly flexuous; costa very large, $\frac{1}{2}$ or $\frac{3}{4}$ the width of leaf at base, canaliculate and very rough on back: sterile. Fl. Miq. 42.—Miquelon.

68. *Dicranum Santeri pachytrichum* Kindb.—Stem very tomentose in its whole length: leaves nearly entire; costa very much narrower than type: pedicel short, about 1 cm. long. Mac. Cat. 260.—On sticks in a brook: Selkirk Mountains.

69. *Dicranum Greenlandicum* Brid.—Dioicous, male plants unknown: appearance of *D. elongatum*: tufts thick, about 6 cm. high, bright yellowish green, slightly radiculose: leaves rigid, when dry densely appressed with slightly bent points, never secund, linear-lanceolate, tubulose near apex, blunt, margins entire; costa weak, $\frac{1}{3}$ greatest width of leaf, percurrent or vanishing below apex; alar cells large, distinct, the remainder elongated, thick walled, pitted: inner perichætal leaves ecostate or with a weak costa: capsule erect, small, about cylindrical, somewhat curved, short and narrow-necked, faintly striate; annulus of one or two rows of cells; lid longer than capsule, obliquely rostrate; peristome orange, teeth divided to below middle, above pale and almost smooth. Limpr. Lævöm. 1: 364.—Miquelon Island; Mansfield Island, Hudson strait; Jupiter River, Anticosti; Labrador; Greenland.

70. *Dicranum fuscescens* Turn.—Dioicous: tufts dense, 1-6 cm. high, mostly brownish green, rarely dark green, not shining, more or less brownish radiculose: stem erect, fasciculately branched, leaves irregularly spreading, rarely secund, when dry somewhat crispate, very long, concave subulate from a narrow lanceolate base, margin toothed above; costa strong, $\frac{1}{2}$ - $\frac{3}{4}$ leaf base; leaf cells not or only sparingly pitted, mamilllose on under side, elongated rectangular below, rectangular in middle and quadrate above: inner perichætal leaves suddenly narrowed above middle into a

long bristle point: seta straw colored, finally red, twisted to right below and to left above; capsule slightly inclined, obovate, curved, six costate, neck distinct, inflated; annulus compound, of 2-3 rows of cells; lid often longer than capsule, obliquely long rostrate from a broad base; peristome teeth dark purple with yellow thickly papillose points, broad and approximate, bi- or trifold for one-third length; spores brown, papillose. Limpr. Laubm. 1: 359. *D. fuscescens longirostre* Schimp., and *angustifolium* Schimp., Lesq. and James Man. 72.—Common on old logs across the continent: United States; Canada; Greenland.

71. *Dicranum fuscescens falcifolium* Braith.—Densely tufted, deep green, fastigate; leaves all falcate-secund, flexuous-cirrhate toward apex, shorter and less attenuated to point. Braith. Brit. Moss Flora 1: 153.—Washington.

72. *Dicranum Miquelonense* Ren. and Card.—In small, compact, yellowish-green tufts: stems dichotomous, radiculose below, 1-3 cm. long; leaves small, short, erect imbricate or slightly incurved, oblong-lanceolate, acute or obtuse, concave, entire or minutely sinuate denticulate at apex, 2-3 mm. long; costa vanishing near apex, smooth or scarcely rugose at back; cells smooth, small, short, quadrate or irregularly angular in upper half, rectangular, 1-3 times longer than broad and thin walled towards base, alar lax, large quadrate or sub-hexagonal, brown or yellowish: fruit unknown. Bot. Gaz. 14: 93. 1899.—On the ground and on rocks, Miquelon Island.

73. *Dicranum neglectum* Juratz.—Dioicous, male plants small and delicate, densely tomentose: intermediate in habit and characters between *D. scoparium* and *D. Muehlenbeckii*: tufts thick, 5 cm. high, faintly shining, sparingly radiculose: stem densely foliate, leaves erect spreading, when dry somewhat curved and twisted, scarcely crispate, brittle, from ovate concave base lanceolate-subulate and tubulose, margin entire, obscurely toothed at apex; costa weak, $\frac{1}{2}$ leaf base, percurrent or excurrent, smooth on under side; cells smooth, nearly as in *D. scoparium* but very thick-walled and profusely pitted, lower elongated-rectangular or linear, upper elongated mixed with quadrate cells: perichaetial leaves tubulose, abruptly narrowed into a subulate point: capsule, lid, annulus, and peristome as in *D. Muehlenbeckii*. Limpr. Laubm. 1: 353.—On rocks: Oregon; Mt. Niblock, Mt. Aylmer and near Hector, B. C.

74. *Dicranum Muehlenbeckii cirrhatum* (Sch.) B. & T. MS.—Similar in appearance to the deep tufted forms of *D. congestum*: tufts to 10 cm. high, slightly radiculose, easily falling apart: leaves not so curved, erect-spreading strongly curved; costa $\frac{1}{2}$ greatest width of leaf; lamina continued to apex, margin and under side of costa toothed; alar cells very small, well-defined, all leaf cells smaller than in *D. Muehlenbeckii*, slightly thickened, lower rectangular, somewhat porose, above very irregular: seta shorter than in

D. Muehlenbeckii; capsule shorter, cernuous; peristome very low, pale purple, within scarcely barred; teeth divided to middle, legs narrowed. *D. Muehlenbeckii brevifolium* Lindb., *Limpr. Laubm.* 1: 355.—Godhavn, Greenland; Digges Island, Hudson Bay.

75. *Dieranum dipteroneuron* C. Müll.—Tufts rather dense, radiculose, an inch high: stems dichotomous, flexuous: leaves laxly erect, rather broad, small, yellowish green, shining, stiffish, smooth, flexuous, when moist rather strict, broadly lanceolate, stoutly acuminate, involute, coarsely and unequally serrate near apex; costa thicker at base, gradually narrowed, with two parallel serrulate low wings on back above; cells large, elliptical above, longer at base, walls thick, pitted, alar cells numerous, brownish: perichaetial leaves similar: seta erect, scarcely an inch long, slender, slightly twisted, red at base, yellowish above; capsule arcuate, very narrow, small, obliquely long rostrate; calyptra very narrow, smooth. *Flora* 70: 219. 1887.—Alaska.

76. *Dieranum pallidum* Bry. Eu. (not Müll.).—In compact green or yellowish tufts: stems 2-4 cm. long, simple or dichotomous, covered below with ferruginous tomentum: leaves erect-patent or subsecund above when moist, flexuous and somewhat crispate when dry, oblong-lanceolate, long narrowed acuminate, generally somewhat inflexed at margins, subcanaliculate, smooth or papillose at back, serrulate above; costa percurrent or excurrent, generally papillose and denticulate at back above, sometimes nearly smooth; cells short, angular, irregular in the upper part, linear, elongated towards base, the alar, large, lax, quadrate or sub-hexagonal, orange brown: outer perichaetial leaves from a broad base suddenly constricted into a more or less elongated denticulate point, the inner convolute sheathing, abruptly subulate from the rounded apex: seta yellowish; capsule pale, cernuous or sub-erect, oblong or subcylindric, arcuate, slightly plicate when old, neck small, strumose; lid conic, long subulate rostrate; annulus distinct; teeth purple, cleft to the middle or below into 2-3 free or coherent legs. *D. sabuletorum* R. & C., *Bot. Gaz.* 14: 91. 1889. *D. spurium* var. *condensatum* L. & J., not *D. condensatum* Hedw., L. & J. *Man.* 76. In dry sandy places on hills especially in southern district: Florida; S. Carolina.

77. *Dieranum scoparium compactum* Ren.—Tufts very compact: leaves slightly homotropous, often broken at point, dentate, acumen short: sterile. *Fl. Miq.* 44.—Miquelon Island.

78. *Dieranum scoparium flexicaule* Ren. & Card.—Stems long, slender, decumbent, then erect, flexuous, innovations slender, reaching or exceeding the capsule: leaves erect incurved, slightly flexuous, somewhat long acuminate, entire or sinuate-crenulate; costa vanishing below apex; cells with delicate sinuate walls not or but slightly pitted: capsule reddish, irregularly rugose-plicate when dry. *Fl. Miq.* 44.—Miquelon Island.

79. *Dicranum scoparium sulcatum* Ren. & Card.—Tufts yellowish, habit more slender: leaves erect or spreading, sometimes flexuous or even a little twisted, narrower, longer and more slenderly subulate, teeth more salient and pointed; cells delicate, generally chlorophyllose, less pitted: seta slenderer, strongly twisted to the right, pale; capsule dark red, distinctly plicate when dry. Fl. Miq. 44.—Miquelon Island.

80. *Dicranum scoparium spadiceum* Boul.—Nearly identical with European plant: leaves erect, entire or sub-entire, acumen smaller, sub-obtuse, costa vanishing below apex; cells sinuous and pitted. Fl. Miq. 44. *D. spadiceum* Zett.—Miquelon Is.; Rocky Mountains.

81. *Dicranum scoparium orthophyllum* Brid.—Leaves rigidly erect, when dry appressed, coarsely dentate. Limpr. Laubm. 1: 352.—Vancouver.

82. *Dicranum scoparium curvulum* Brid.—Stems arcuate ascendent; leaves falcate-secund, narrower, yellowish-green; capsule shorter. Husn. Muscol. Gall. 35.—Oregon and Washington.

83. *Dicranum scoparium crispulum* De Not.—Plants low: leaves crispate when dry.—Washington.

84. *Dicranum consobrinum* Ren. & Card.—Densely caespitose, yellowish green: stems erect, simple or dichotomous, tomentose, 5-8 cm. long: leaves rather crowded, secund or erect spreading, narrowly lanceolate-subulate, serrate in the upper half; costa serrate at the back toward the apex; cell-walls porose, scarcely thickened: perichaetial bracts sheathing, truncate or emarginate at the apex, sometimes mucous, generally tipped with a short or little elongated subula: seta yellow, sub-flexuous; capsule cernuous or horizontal, narrowly cylindrical, curved, not sulcate, long attenuate below, rufescent when old; lid long subulate: male plants gemmaceous, nidulant in the tomentum of the female stems. Bot. Gaz. 15: 39. 1890.—Minnesota.

85. *Dicranum Bonjeani Schlotthaueri* Barnes.—Mostly olive green: stem very short, 1-2 cm. high: leaves shorter and broader, margin entire or with a few indistinct teeth at the apex, often somewhat revolute. Bot. Centralbl. 44: 386. 1890.—Oregon; Washington; Idaho; Wyoming.

86. *Dicranum Bonjeani Rollii* Barnes.—Plants robust, in deep extended quite lax tufts, 4-8 cm. high, below pale, above yellowish green: leaves crowded, not undulate, lanceolate, entire, acuminate; costa vanishing in the acumen, only faintly canaliculate on the back and not toothed. *Loc. cit.*—Vancouver.

87. *Dicranum Bonjeani alatum* Barnes.—Dark green, 4-6 cm. high: leaves lanceolate, very sharply toothed; costa 2-3 cells thick, with 2-3 sharply toothed lamellæ; laminal cells shorter and broader, less thickened and less strongly pitted. *Loc. cit.*—Chicago, Ills.

88. *Dicranum brachycaulon* Kindb.—Allied to *D. spurium*: stems short, only 1 cm. high: the leaves smaller and shorter, oblong-ovate, acute,

not acuminate, entire, not papillose at the back; costa elevate, percurrent and smooth, alar cells brown: capsule small, pedicel 1 cm. long. Mac. Cat. 34.—On dry rocks in the open woods: Yarmouth, N. S.

89. *Dicranodontium Virginicum* Britt.—Dioicous: plants bright glossy green: stems matted below by a red tomentum, leafy nearly to apex, denudate roughened above, with a few leaves at summit: leaves erect or secund, straight or curled and twisted, narrowly subulate from a short thick base; caducous ones with a long slender smooth point; persistent ones serrate, blade inflexed, cells densely chlorophyllose, filled with oil globules, those of basal angles clear: perichaetial leaves from a short base suddenly subulate, dentate at apex: pedicels lateral by innovations, pale, glossy yellow, twisted in two directions, very slender, arcuate when young, becoming erect before mature; capsules cylindric, ribbed only at mouth, beak straight or curved, shorter than capsule; peristome bright red, not deep set, teeth split unequally to middle, striolate at base, pale and granulose above; annulus none, spores small, calyptra cucullate. Prelim. Cat. Flora W. V. 488.—On sandstone boulders: Monongalia, W. Va.

90. *Dicranodontium Millspaughii* Britt.—Dioicous: plants light yellowish green, silky, cespitose: stems matted with rufous tomentum at the base, a few denudate-roughened by fragments of slightly caducous leaves: leaves secund or erect-spreading, narrowly subulate from a broad base, becoming tubular above by inrolled margins, basal angles not auricled, filled by large hyaline cells to base of broad brown vein, those of blade oblong or square next the vein, becoming spindle shaped and prosenchymatous towards margin; costa thick, excurrent into a slender dentate tip, rough on back: perichaetial leaves sheathing half their length, tapering to a long slender obscurely serrate tip, outer shorter, abruptly subulate, more sharply serrate: pedicels recurved, burying capsule among leaves, erect when old, stout and twisted in two directions; capsules pyriform-cylindric with a distinct neck; lid as long as capsule, straight beaked; calyptra cucullate, entire; peristome red, connivent, teeth deep set, slender, split to middle or perforate, striolate below, granulose above; annulus none. Prelim. Cat. Flora W. Va. 488.—On sandstone boulders in deep woods: Tibbs Run, W. Va.

91. *Campylopus Schimperii* Milde.—Dioicous: tufts dense, compact, interwoven with rufous tomentum, fastigiata: stems slender, cuspidate, light silky green above, fuscous below, dichotomous with alternate innovations which are easily detached: leaves appressed when dry, erect-spreading, straight, rigid, narrowly lanceolate subulate, slightly toothed at apex, channelled in lower part, becoming tubulose above from incurved wings; costa broad, $\frac{2}{3}$ width of leaf base; cells of the extreme base brown and vesicular, above hyaline, very narrow at margin, elongate rectangular towards the nerve, upper small and elliptic: perichaetial leaves sheathing,

suddenly narrowed into a long subula; capsules pale, ovate, striate; annulus broad; lid half length of capsule, beaked; peristome small; teeth cleft to middle; spores large. Braithw. Brit. Moss Flora 1: 130.—Greenland.

92. *Campyopus Henriei* Ren. & Card.—Cespitose, yellowish green: stems very short, without tomentum: leaves slightly secund, lanceolate-subulate and semitubulose from an oblong base, upper generally tipped with a short hyaline denticulate and often broken point; basal cells rectangular (3-4:1), those of angles sometimes rather soft and yellowish, not forming distinct auricles, upper elongated, straight, linear; costa broad, $\frac{1}{2}$ width of leaf: male flowers small, gemmiform, placed near tips of stems: female flowers and capsule unknown. Bot. Gaz. 18: 198. 1888.—Sandy ground: Saline county, Kansas.

93. *Fissidens incurvus brevifolius* Ren. & Card.—Leaves broader and shorter: border of the vaginant lamina widening less at the base. Bot. Gaz. 14: 94. 1889.—On the ground in woods: Baton Rouge, La.

94. *Fissidens pusillus* Wils.—Heteroicous: plants very small, simple or branched at base: stems short, inclined: lower leaves very small, larger above and mostly secund, narrow lanceolate, becoming smaller towards apex, sharp pointed, margin quite entire, border narrow, vanishing below apiculus, costa lost at point; vaginant lamina not reaching middle of leaf, inferior lamina semi-lanceolate, rapidly narrowed at base and lost before reaching stem; cells oval or rounded: capsule on a pale seta, very small, erect or inclined, oval-cylindric, strongly contracted below mouth when dry; lid conic, somewhat obliquely rostrate; peristome deep red, arising below orifice, teeth deeply cleft, legs subulate, filiform, rough; spores brown, smooth. Braithw. Brit. Moss. Flora 1: 68.—Abundant on damp flat limestone rocks in woods: Ontario.

95. *Fissidens obtusifolius Kausanus* Ren. & Card.—Leaves with a broad border of elongated cells on the margins of vaginant lamina, and a narrow more or less distinct border on dorsal wing. Bot. Gaz. 15: 40. 1890.—Saline county, Kans.

96. *Fissidens taxifolius Langloisii* Ren. & Card.—Habit a little more robust, leaves subobtuse, apiculate, dorsal lamina suddenly rounded at base as if auriculate. Revue Bryol. 19: 78. 1892.—Louisiana.

97. *Fissidens adiantoides brachyphyllus* Kindb.—Leaves very short. Mac. Cat. 37.—On boulders: Belleville, Ont.

98. *Fissidens falcatus* Ren. & Card.—Very small, gregarious, yellowish green: stems rather rigid, plumulose: leaves 4-8 pairs, falcate secund and rigid when dry, linear lanceolate, acute or sub-apiculate; vaginant lamina about one-half length, narrowly bordered, dorsal lamina not bordered, tapering below, apical lamina without a border, subentire or minutely crenulate at apex; costa pale, subpercurrent; cells hexagonal, pellucid: fruit unknown. Bot. Gaz. 19: 237. 1894.—On the bark of trees: Louisiana.

99. *Fissidens pauperculus* Howe.—Dioicous: minute, loosely gregarious, flavescent; stems decumbent or ascending; leaves 3-5 pairs, increasing in size upward, lowest minute, upper oblong to obliquely spatulate-oblong, acute or shortly acuminate, margin slightly serrulate crenulate; border none; costa stout, vanishing below apex; vaginant lamina $\frac{1}{6}$ to $\frac{1}{2}$ length of leaf, unequal; inferior lamina ending at about middle of vaginant lamina in upper leaves or reaching base in lower; cells of vertical lamina mostly hexagonal, smaller and oval at margin, enlarged and oblong rectangular next to costa, those of vaginant lamina becoming longer and narrower towards base: seta terminal, flexuous, pale yellow; capsule yellowish, oval or oval-oblong, inclined or cernuous, slightly arcuate when dry; lid conic-rostellate, about equaling capsule; peristome red, normal; annulus pale, of 2-3 rows of deciduous cells. Erythea 2: 97. 1894.—On moist banks in company with *F. limbatulus* Sulliv., Marathon county, Cal.

100. *Ceratodon conicus* Hpe.—Dioicous: tufts caespitose, dull yellow green above, fuscous below: stems short, rather slender, dichotomously branched: leaves erect appressed, deep green, smooth, crowded at tip into a small closed coma, straight wet or dry, rather broadly ovate lanceolate, margin quite entire, revolute to apex, nerve thick, excurrent into a long point; cells all small, regularly quadrate, basal larger and pellucid: perichaetial leaves convolute-vaginant, obtusate with an excurrent nerve, laxly areolate: capsule on a pale red seta, erect, ovate-elliptic, rather wide, fuscous, when dry and empty but little altered, sulcate in upper part, not strumulose; lid purple, short conic; teeth pale, red at base, yellowish above, erect with fewer articulations, scarcely bordered externally. Braithw. Brit. Moss. Flora 1: 175.—At the base of a stump, Spence's Bridge, B. C.

101. *Ceratodon heterophyllus* Kindb.—Agrees with *C. purpureus*, in shape of capsule, stem leaves, not excurrent costa and revoluble annulus, but capsule often more curved and distinctly strumulose; agrees with *C. conicus* in peristomial teeth having few articulations: differs from both in blunt perichaetial leaves, and is also very peculiar in short concave sub-oval leaves of long shoots. Ott. Nat. 5: 179.—On earth: St. Paul Island, Behring Sea.

102. *Trichodon flexifolius* Ren. & Card.¹—Loosely caespitose, green: stems erect, simple, short: lower leaves small, lanceolate subulate, upper larger, patulous, very flexuous, from an oblong base gradually narrowed into a long linear subulate canalliculate subtubulose point, sinuate at margins, toothed at apex, and with a broad obscurely excurrent costa; basal cells rectangular, elongated (1:2-4), others small, quadrate, very chlorophyllose, with transverse walls slightly prominent: perichaetial leaves similar, but dilated and sub-sheathing at base: male flowers on same plant,

¹The authors of this species are not certain as to its generic position, but state that by its vegetative system it is somewhat allied to *T. cylindricus*.

below female, small, gemmiform. Bot. Gaz. 14: 94. 1899.—Sandy ground: Florida.

103. *Ditrichum*¹ *montanum* Leiberg.—Plants caespitose, fastigiate branching, above more or less dichotomous: stem leaves erect or slightly curved, channeled and subulate above with a narrow lamina of two to three rows of cells, subserrulate, margin inflexed, apex coarsely toothed; costa broad, strong, vanishing below apex; leaf cells above and in middle hyaline or chlorophyllose, thick walled, quadrate rectangular below: plants monocous, seldom synlocous: outer perichastial leaves similar, inner broadly sheathing: seta pale yellow, slightly twisted when dry; capsule narrowed at mouth, erect, elliptical; when dry laterally compressed and longitudinally wrinkled; teeth cleft to base, legs equal, semiterete, minutely papillose, obscurely and distantly articulate, attached to a short basilar membrane; annulus double, narrow, dehiscent; lid long, conical. Bull. Torr. Bot. Club 20: 112. 1893.—On the broken soil, upturned tree roots, etc in mountain regions, at all elevations up to 8,000 feet, Idaho.

104. *Ditrichum* *ambiguum* Best.—Dioicous: tufts loosely caespitose, stem rather stout, arcuate-erect with one or more innovations: leaves pale yellow, shining, crispate when dry, accrescent upwards, lanceolate-subulate, patent-subsecund, flexuose; lower erect at half clasping short base, lanceolate; upper with oblong erect bases, long lanceolate subulate, concave, the slightly thickened involute margins sinuate dentate; cells linear, oblong, indistinct above; walls thick, tortuous, striate; costa percurrent and dentate on back: perichastial leaves with longer sheathing bases not abruptly narrowed: seta long flexuose; capsule cylindrical, narrow, straight or slightly curved; lid conical-rostrate, blunt, nearly or quite erect, about $\frac{1}{2}$ length of capsule; teeth reddish, long, straight, papillose, nearly or quite split to broad basal membrane; legs filiform except at slightly flattened connate bases, equal and regular: annulus large, adherent. Bull. Torr. Bot. Club 20: 117. 1893.—Moist banks: Mason co., Washington.

105. *Ditrichum* *flexicaule brevifolium* Kindb.—Leaves very small, from an ovate oblong base, contracted to a short point; costa not excurrent: barren. Mac. Cat. 46.—On rocks: National Park; Rocky mountains.

106. *Ditrichum* *Macounii* C. M. & Kindb.—Tufts dusky green, very dense, compact, radiculose below: stems 2-3 cm. high: leaves patent or subfalcate from a short, suddenly narrowed, sheathing base, entire or with a few small teeth at apex; cells short subquadrate; costa sub-percurrent: barren. Mac. Cat. 40.—On banks subject to inundation: Columbia river, B. C.

107. *Seligeria campylopoda* Kindb.—Agrees with *S. recurvata* in shape of capsule and arcuate pedicel: differs in leaves broader, very much shorter, sub-linear obtuse, rarely short acuminate and subacute; costa not

¹*Leptotrichum* of L. & J. Man. 106.

excurrent: perichaetial leaves not ovate-oblong, thin costate: peristome darker red: male flower fixed on side of female. Mac. Cat. 41.—Damp and shaded limestone rocks: Owen Sound, Ont.

108. *Blindia acuta flexipes* Ren. & Card.—Pedicel flexuous, distinctly geniculate. Revue Bryol. 19: 79. 1892.—Oregon.

109. *Pottia heimloides* Kindb.—Nearly allied to *P. Heimii*: differs in leaves shorter, the lower obtuse; costa sometimes excurrent: capsule longer and narrower, cylindrical; seta golden yellow; peristome present but rudimentary. Mac. Cat. 43.—On earth: National Park; Rocky mountains.

110. *Pottia intermedia* Förn.—Scarcely distinguished from *P. truncatula* with which it agrees in inflorescence, structure of stem and costa: plants larger, stem erect, longer, lax leaved below, dense above: leaves pale green, accrescent upwards, upper leaves long lanceolate, acuminate, margin revolute from base to middle; costa excurrent into a long yellowish-green point; cells in upper half quadrate or hexagonal, faintly papillose or perfectly smooth: capsule obovate to almost cylindrical, constricted below mouth when dry and with short, irregular folds; calyptra smooth, cucullate, covering half of capsule; lid about equaling capsule, obliquely rostellate; annulus compound, 2-3 rows of cells; peristome rudimentary; spores finely papillose. Limpr. Laubm. 1: 531.—On earth: North West Territory.

111. *Pottia littoralis* Mitt.—Autoicous; resembling *P. intermedia*, pale below, green or bluish green above: leaves much longer, more erect, sheathing at base, more obtuse, with nerve excurrent in a short point, lower smaller with a long excurrent costa; upper cells smaller, quite smooth, with wall much more incrassate, basal pellucid, very narrow and elongated: seta pale orange red, elongated; capsule oblong oval, slightly narrowed at mouth, ferruginous; calyptra smooth, annulus adherent, of one row of cells; lid rostrate, slightly twisted, oblique; spores ferruginous, scarcely rough. Braithw. Brit. Moss Flora 1: 198.—On earth: Yale, B. C.

112. *Didymodon Canadensis* Kindb.—Differs from *D. rubellus* principally in perichaetial leaves thinner, from the ovate base abruptly attenuate to a short subulate acumen; borders not reflexed; basal cells very long; costa thinner: dioicous. Mac. Cat. 44.—On perpendicular rocks: Rocky mountains.

113. *Didymodon Baden-Powellii* Kindb.—Dioicous: tufts compact: leaves revolute nearly all around, distinctly dentate: short acuminate, the lower pale brown: perichaetial leaves longer acuminate or subulate, entire: capsules more or less curved; pedicel pale red; lid blunt conic, very short (scarcely $\frac{1}{2}$ capsule). Mac. Cat. 262.—St. Paul Island, Behring sea.

114. *Didymodon Hendersoni* Ren. & Card.—Tufts compact, yellowish above, ferruginous below: stems erect, branched, 1-2 cm. long: leaves crowded, patulous when moist, subincurred, erect-imbriate when dry, ovate or oblong-lanceolate, entire; apex rounded-obtuse or minutely apic-

ulate or subacute, borders revolute but flat below point; costa stout, rufescent when old, vanishing at or below apex; cells small, distinct, thick-walled, irregular, roundish quadrate, minutely papillose, lower rectangular, rather elongated towards costa, quadrate or transversely dilated on borders; perichaetial leaves not sheathing, oblong-lingulate, obtuse at apex; seta reddish, twisted to left above; capsule erect, cylindrical, badius when old, lid obliquely rostrate; peristome unknown. Bot. Gaz. 15: 40. 1890.—Crevices of rocks: Oregon.

115. *Didymodon rufus* Lorentz.—Dioicous: tufts compact or loose, dark reddish brown, brownish green above; stems 2-4 cm. high, erect or geniculate, forked or fasciculately branched; leaves crowded, spreading and recurved above middle, when dry erect appressed or slightly twisted, gradually acuminate from an ovate lanceolate base, margin entire, revolute on border; costa percurrent or vanishing below apex, round, rough on both sides; lamina of one layer of cells, long mamilllose-papillose; cells equally and strongly thickened, lumen round or transversely oval, at base near costa elongated rectangular, faintly pitted, and several marginal rows quadrate; fruit unknown. Linpr. Laubm. 1: 559.—Greenland.

116. *Leptodontium Hampe*.—Leaves squarrose-reflexed, complicate-concave, flexuous, the margin serrulate or crose-crenulate; capsule leptodermous, narrow, erect; lid conic; peristome of 32 filiform straight erect smooth teeth, unequal or anastomosing here and there in pairs, or connascent; calyptra cucullate.

117. *Leptodontium Canadense* Kindb.—Tufts loosely cohering, nearly without rhizoids, dark green, blackish below; stem about 1 cm. high, nearly simple; leaves squarrose, flexuose, undulate and not recurved at borders, when dry curled, sublingulate, acute, entire below middle, coarsely and unequally serrate above, not margined; lower basal cells rectangular and subpellucid, upper at middle quadrate, uppermost rotundate, slightly papillose; costa finally brown, short excurrent; dioicous, only female plants found. Mac. Cat. 45.—On stones: in the Sydenham river, Owen Sound, Ont.

118. *Trichostema nitidum* Schimp.—Dioicous: tufts compact, pulvinate, 1-2 cm. high, olive-green, within reddish brown and radiculose; stems rigid and dense-leaved; leaves accrescent upwards, fragile, spreading when moist, when dry arcuate-incurved, with faintly crenulate incurved margins, glossy shining on the back, linear lanceolate, mostly obtuse, short pointed by excurrent costa; hyaline cells at base forming $\frac{1}{2}$ leaf, hyaline cells continued upwards as a border, but less sharply marked off from chlorophyllose cells, latter round-quadrate, with simple or geminate papillae on both sides; perichaetial leaves half sheathing, gradually diminishing in size, almost subulate; capsule erect, nearly cylindrical, faintly curved, when old somewhat furrowed; lid $\frac{1}{4}$ capsule, obliquely rostrate;

annulus indistinct; teeth of peristome rudimentary, yellowish-red, truncate, papillose (?). Lampr. Laubm. 1: 581.—United States, without locality, collected by James and communicated by Bescherelle. Bot. Gaz. 14: 90. 1889.

119. *Trichostomum Vancouveriense* (Broth.) R. & C.—Dioicous: loosely caespitose, fragile, bright green, shining; stem simple, 3 mm. high, long radiculose below, densely foliate above: leaves twisted when dry, arcuate-inflexed, margin involute; when moist straight, spreading, nearly plane, linear lanceolate from short base, acute, margin slightly undulate, remotely and obtusely denticulate from below middle upward; costa green, excurrent; lamina bistratose, chlorophyllose cells round-quadrate, basal cells elongated, hyaline: perichaetial leaves scarcely different: capsule erect or suberect, cylindrical, shining, striate, short-necked; annulus broad, triple, revolute; peristome simple, pale, tubulose base short, teeth erect, strongly papillose. *Timmia Vancouveriensis* Broth. Bot. Centralbl. 44: 387. 1890.—On slightly moist ground: Victoria, Vancouver Island.

120. *Desmatodon subterrescens* C. M. & Kindb.—Nearly allied to *D. atrovirens* Smith (*D. nervosus* B. & S.), but tufts compact: stems higher (about 1 cm.): leaves green, not dingy, very short, suboval or sub-spathulate, smaller, less opaque; costa neither excurrent nor broader above: capsule cylindric. Mac. Cat. 48.—On earth on exposed cliffs: Gaspé coast, Can.

121. *Desmatodon camptothecus* Kindb.—Habit of *D. cernuus*: plants densely caespitose: leaves long, narrow-lingulate, sub-obtuse, entire or obscurely crenulate, faintly papillose, marginate and at base revolute; mucronate by excurrent greenish costa: capsule cylindric, arcuate; teeth short, cut into 2 or 4 partly coherent segments; annulus distinct; lid obliquely short beaked; seta flexuous, reddish yellow, spores large. Mac. Cat. 49.—Rocks: Gaspé coast, Can.

122. *Desmatodon cernuus xanthopus* Kindb.—Leaves less chlorophyllose, costa virescent: capsule larger; teeth more united; pedicel yellow. Mac. Cat. 48.—On earth: Manitoba.

123. *Leptobarbula* Sch.—Plants small, delicate, gregarious: leaves minute, linear, sparse, strongly verruculose, areolation dense, not carinate: flowers dioicous, male gemmiform, terminal: perichaetium sheathing: calyptra long, narrowly cucullate; annulus compound, broad, revolute; teeth of peristome perfect, loosely one-half twisted; spores minute, smooth. Syn. Musc. Eur. 181. 1876.

124. *Leptobarbula berica* Sch.—Dioicous: plants gregarious or forming small tufts, bright green: stems very low (1-5 mm.), mostly simple, radiculose only at base: leaves when dry crispate, when moist spreading and somewhat reflexed, thickly warty papillose on both sides and costa; lower leaves distant, lance-subulate, upper gradually longer (.85 mm.),

uppermost (1.5 mm.) and perichæstial leaves (3 mm.) from a sheathing base ($\frac{1}{2}$ of leaf) gradually or suddenly narrowed into an acute or obtuse recurved subula; lamina with flat margins, very narrow in subula, crenulate by papillæ; cells of base elongate-rectangular, thick walled, the upper small quadrate (6-8 μ); seta erect, 6-13 mm. long; capsule erect, cylindric-oval; lid $\frac{1}{2}$ the sporangium, obtuse; annulus of 3-4 rows of large cells, revolvable; peristome tube .04 mm. high, teeth filiform, not nodose. Linpr. Laubm. 1: 598.—On earth: Revelstoke, B. C.

125. *Barbula macrorhyncha* Kindb.—Habit of *B. brevirostris*: leaves round-ovate, obtuse, involute all around even at the apex; costa not excurrent; capsule oblong-ovoid; annulus revolvable; teeth long, twisted at least twice, beak of lid more than $\frac{1}{2}$ length of capsule; seta pale red or yellowish; dioicous; calyptra unknown. Different from *B. rigida* in very much shorter leaves and longer lid. Mac. Cat. 50.—On earth: Ontario.

126. *Barbula Henriei* Rau. (§ CHLORONOTÆ).—Dioicous (?): plants short, branched, closely cespitose, canescent from white excurrent costæ: leaves spreading when moist, imbricated when dry, concave, short spatulate; costa keeled, filamentose near apex of leaf, excurrent portion as long as leaf, hyaline, serrate; cells quadrate and chlorophyllose above, hyaline and elongated toward base: fruit unknown.—On rocks: Saline Co., Kansas.

127. *Barbula Mannii* C. Müll.—Slender, small, bright green: stems short, simple: leaves minute, few; base erect, slender, loosely reticulate, concave; lamina for the plant elongate, narrow, oblong; apex roundish obtuse; margin obscure, quite entire, revolute; cells minute, hexagonal, costa rather thick, with a few narrow longitudinal lamellæ and deciduous gemmæ above: inner perichæstial leaves very narrow, linear-oblong; seta long, red, flexuous; capsule small, erect, cylindraceous, brown; lid with slender oblique beak; annulus persistent, rather large; peristome membrane pale, short, teeth slender, red. Flora 70: 222. 1887.—On calcareous soil: Colorado.

128. *Barbula Egelingi* Schlieph.—Dioicous: very small, slender, simple: leaves aggregated into a minute tuft, few, small, imbricate; lamina oblong lanceolate, very entire, from a more pellucid loosely reticulate broader base deeply carinate-concave; margin above convex, scarcely revolute, rather broad, of larger yellow cells; costa slender, excurrent; cells minute, roundish hexagonal, thick, obscure: seta very slender, long, flexuous, red; capsule erect, minute, very narrowly cylindric; lid very short conic, oblique, spirally twisted; annulus narrow, simple, persistent; peristome with a short membrane. Flora 70: 222. 1887.—Memphis, Tenn.

129. *Barbula carnifolia* C. M. & Kindb.—Stem very short, 2-3 mm. high; leaves lingulate, not margined, blunt; margins recurved below and above the middle at one side; cells finally subpellucid, upper and median subquadrate, lower basal near costa large rectangular and hyaline, margi-

nal shorter and narrower; costa red, thick, and subpercurrent or rarely short excurrent: inner perichætical leaves smaller and shorter: capsule longer than the straight lid: probably dioicous. Mac. Cat. 52.—*Sine loco*.

180. *Barbula subcarnifolia* C. M. & Kindb.—Smaller than *B. carnifolia*, differing principally in leaves shorter, suboblong, short apiculate, upper pellucid at flat margins, costa green: probably monoicous. Differs from *B. amplexa* in capsule longer than oblique lid. Mac. Cat. 52.—On earth at the base of trees: Pelee Island, Lake Erie.

181. *Barbula brachyphylla angustifolia* C. M. & Kindb.—Tufts compact, sparingly radiculose, rusty-red, bright-green at summit, 1-2 cm. high: leaves shortly ovate-lanceolate, acute or subobtuse, indistinctly papillose, densely disposed, when dry appressed, borders reflexed nearly all around; cells small, round-quadrate; costa broad, reddish and percurrent: barren. Mac. Cat. 55, 264.—On calcareous rocks where water drips in spring: Columbia river, B. C.

182. *Barbula decursivula* Kindb.—Tufts rusty red with green tops, about 2 cm. high: stem radiculose below, not rigid, branches long and filiform: leaves small, when dry loosely appressed, open when moist, very broad at the base, subovate or ovate-oblong, acute or subobtuse, nearly smooth or faintly papillose at back, loosely disposed, long decurrent; borders reflexed nearly all around; cells pellucid, subquadrate, only the alar elongate; costa reddish, broad and percurrent: barren. Mac. Cat. 264.—On earth and rocks: Hector, B. C.

183. *Barbula sparsidens* C. M. & Kindb.—Loosely tufted: plants small, green above, ferruginous below, nearly simple: leaves small, when dry incurved, when moistened subsquarrose, arcuate, ovate-lanceolate, distinctly papillose; borders reflexed nearly all around; lower basal cells rectangular, hyaline; costa pale, rough at back, percurrent: inner perichætical leaves shorter than outer; costa not or very short excurrent: capsule small, oblong cylindrical, reddish, longer than oblique beak; peristome and seta red: dioicous. Allied to *B. fallax*, but differing principally in peristomial teeth spreading, loosely and not spirally twisted. Mac. Cat. 54.—On earth: British Columbia.

184. *Barbula Dieckii* Broth.—Dioicous: tufts densely caespitose but loosely cohering, above tawny to dark olivaceous: stem erect, dichotomously branched, fastigiate, loosely foliate: leaves when dry imbricate, when moist spreading, apex erect, uniform, concave, broadly ovate lanceolate, acuminate, acute, papillose, margin entire, revolute to a little beyond middle; costa green, subterete, strong, above weaker, vanishing at apex; lamina unistratose, cells thick walled, rotund-quadrate, strongly chlorophyllose, papillose on both sides, basal cells scarcely larger, subquadrate: perichætical leaves similar to others, but long acuminate, base loosely retic-

ulate, inner ones smaller, margin plane: sterile. Hedwigia 82: 205. 1893.
—Growing on rocks: Washington.

185. *Barbula subgracilis* C. M. & Kindb.—Densely tufted; plants small, about 6 mm. high, green above, reddish brown below, nearly simple: leaves when dry subcrispate, when moistened squarrose, curved from ovate base, narrow, long subulate, nearly smooth; borders reflexed nearly all around; lower basal cells rectangular, more pellucid; costa reddish, percurrent or short excurrent: inner perichaetial leaves broad, blunt, with a long excurrent costa, others acuminate, all faintly crenulate: capsule oblong cylindrical, red-brown, longer than obliquely subulate lid; seta red: dioicous. Allied to *B. gracilis*. Mac. Cat. 53.—On rocks: Yale, B. C.

186. *Barbula subgracilis viridior* Kindb.—Plants higher, about 1.5-2 cm., green and branching above, decolorate-brown below: leaves when dry crispate, more papillose; costa green: capsules not found. Mac. Cat. 263.
—On rocks: Sydenham river, Ont.

187. *Barbula sublemnophila* C. M. & Kindb.—Laxly tufted; plants about 1 cm. high, pale brown, nearly simple: leaves when dry loosely appressed, when moistened subsquarrose or patent, nearly straight, short, ovate-lanceolate, acute, faintly papillose; borders slightly reflexed at base, inflexed at upper part, papillose crenulate; cells nearly uniform, roundish or subquadrate, apical often larger and pellucid; costa pale, percurrent: perichaetial leaves from a short ovate base long-acuminate sublinear; costa dark brown, long excurrent, filling nearly whole acumen: capsule oblong-cylindrical, finally blackish; pedicel dark brown: dioicous. Mac. Cat. 53.—
On dry rocks, covered at high water: Yale, B. C.

188. *Barbula melanocarpa* C. M. & Kindb.—Laxly tufted: fertile stem nearly simple, about 2 mm. high, the barren branching, 4 mm.: leaves when dry appressed, not twisted, when moistened open, olive green, short ovate, lanceolate acute, slightly papillose; borders reflexed below middle; cells nearly uniform, roundish or subquadrate; costa brown, short excurrent: perichaetial leaves from ovate oblong base long acuminate, costa filling whole linear acumen: capsule oblong-cylindrical, straight, finally blackish, obliquely short beaked; annulus not distinct, pedicel red: dioicous. Mac. Cat. 54.—On rocks close to the water: Yale, B. C.

189. *Barbula spadicea* Mitt.—Dioicous: resembling *B. fallax*, but more robust, in looser thicker tufts, dull brownish green above, fuscous below: stems simple or branched: leaves when dry incurved and imbricated, when moist spreading from the base, recurved, from a broadly ovate base elongated lanceolate, channelled, margin recurved in lower half, folds more distinct, nerve strong, distinct to apex; cells incrassate and rounded quadrate from base, only lowest elongated oval, obscure above, papillose: perichaetial leaves lanceolate, recurved from a long, lax-celled base: seta red; capsule erect, cylindrical, slightly curved, castaneous with a red mouth;

annulus of 3 to 5 rows of small cells; lid short rostrate, nearly half length of capsule; peristome short, teeth red, on a very short orange basal membrane, scarcely twisted. Braithw. Brit. Moss Flora I; 263.—On earth by a brook: Lytton, B. C.

140. *Barbula pseudo-rigidula* Kindb.—Agrees with *B. rigidula* in tufts fuscous: leaves from a short erect base, patent or recurved, contorted when dry, long lanceolate, revolute below, nearly smooth; basal cells pellucid and rectangular; costa brown, nearly filling narrow acumen: differs in leaf borders not thickened, inner perichaetial leaves from an oblong base, narrow acuminate with a long excurrent costa: peristome pale and distinctly contorted. Mac. Cat. 264.—On earth by a brook: Lytton, B. C.

141. *Barbula circinnatula* C. M. & Kindb.—Nearly allied to *B. cylindrica* or rather intermediate between *B. elata* and *B. virescens*: differs in upper leaves green, circinnate-twisted when dry: peristome paler and annulus simple: leaf cells more distinct. Mac. Cat. 56.—On earth on rocks: Agassiz, B. C.

142. *Barbula horridifolia* C. M. & Kindb.—Densely tufted: plants about 3 cm. high, green above, rusty red below, more or less branching: leaves crispate when dry, when moist falcate, spreading at all sides, very long and narrow, ovate-lanceolate, carinate above, loosely disposed and long-decurrent, nearly smooth; borders reflexed at ovate base, basal cells pellucid, short rectangular, inner larger, upper cells small and obscure; costa red-brown, broader below, narrow above, vanishing in apex: barren. Mac. Cat. 57.—On damp rocks: Columbia River, B. C.

143. *Barbula robustifolia* C. M. & Kindb.—Very nearly allied to *B. tortellifolia*: differs in dark brown or olive-green color of whole plant: leaves more crowded, shorter, with an ovate oblong base, borders reflexed nearly to the middle on one side; upper cells larger and subpellucid, basal ones hyaline, and subquadrate; costa thicker and not excurrent, linear and distinct to apex: barren. Mac. Cat. 56.—Abundant on rocks: Vancouver; British Columbia.

144. *Barbula tortellifolia* C. M. & Kindb.—Very much resembling *B. horridifolia* in habit: generally more robust (often 4-5 cm.) and finally sometimes quite red: stem more divided: leaves broader, less distinctly decurrent, comal ones larger and crowded, short-pointed by excurrent costa: perichaetial leaves suddenly acuminate, costa faintly excurrent capsule large, subcylindric, reddish, twice longer than beak; peristome long, once loosely contorted, whitish: dioicous. Mac. Cat. 56.—On wet rocks: Vancouver and B. C.

145. *Barbula subcylindrica* Broth.—Dioicous: tufts densely caespitose, pulvinate, dark colored above: stems up to 6 cm. high, erect, flexuous, dichotomously branched, branches fastigate, densely foliate: leaves when

dry crispate, when moist recurved from the erect base, carinate-concave, comal longer, lanceolate subulate from lanceolate base, obtuse, minutely papillose, margin entire, revolute, plane; costa reddish, apex terete, slightly narrower, excurrent; lamina unistratose, cells round-quadrate and transversely oval, basal quadrate and short rectangular: perichaetial leaves similar, entire, base laxly areolate: fruiting specimens unknown. Bot. Centralbl. 44: 387. 1890.—Washington.

146. *Barbula pl. jaebra* C. M. & Kindb.—Dioicous: tufts compact, semipulvinate, light brown: stems scarcely 1 cm. high: leaves suberect, when dry imbricated and contorted, short, ovate or ovate-oblong, subacute, papillose; margins strongly revolute; basal cells short rectangular, others subquadrate; costa thick, short excurrent or percurrent: barren. Idac. Cat. 52.—On dry rocks: British Columbia.

147. *Barbula convoluta obtusata* Kindb.—Leaves broader, generally obtuse, rarely subacute, not apiculate. Mac. Cat. 265.—On limestone rocks: Owen Sound, Ont.

148. *Barbula chrysopoda* C. M. & Kindb.—Differs from *B. convoluta* in short nearly indistinct stem: leaves sublingulate, very obtuse: perichaetial leaves rounded or truncate at the apex, long exserted: lid spirally contorted. Mac. Cat. 57.—On earth in burnt woods: Revelstoke, B. C.

149. *Barbula inclinatala* C. M. & Kindb.—Dioicous: tufts laxly coherent, yellowish green: stem indistinct, not branching: leaves cirrhate-crisped and rigid when dry, suberect and nearly straight when moist, not undulate, from a thin ovate-oblong base attenuate, nearly sublinear, acute or suddenly pointed, very papillose; margins involute, cucullate above; costa yellow, pellucid, excurrent in an often denticulate point: perichaetial leaves much broader and longer than others, whitish, subulate from a long lanceolate base; cells narrow, linear above also: capsule narrow cylindrical, nearly straight, suberect; peristome long, several times convolute. Mac. Cat. 50.—On earth and gravel bars: Illicillewaet river, B. C.

150. *Barbula subulata longifolia* Kindb.—Intermediate between *B. subulata* and *B. angustata*: leaves long and narrow, acuminate and acute, distinctly denticulate above, papillose, yellow, bordered; costa long excurrent but shorter in perichaetial leaves. Mac. Cat. 53.—On earth: British Columbia.

151. *Barbula alpina* Bruch & Schimp.—Closely related to *B. levispila*: leaves not emarginate at summit, ending in a short point or simply mucronate: monoicous: tube of peristome half as long, scarcely one-sixth whole length; teeth once twisted. Husnot, Musc. Gall. 114.—On rocks: Quesnel, B. C.

152. *Barbula ruralis subintermedia* Ren. & Card.—Differing from related *B. intermedia* in leaf margin revolute to apex and more dentate. Ren. & Card. Revue Bryol. 19: 84. 1892.—Arizona.

153. *Barbula ruraliformis* Besch.—Plants robust: stems 3-5 cm. high, branching: leaves squarrose spreading when moist, erect-contorted when dry, acuminate, lamina extended along each side of point as a scarious and denticulate membrane; point reddish in lower part, dentate: perichæstrial leaves strongly plicate: capsule and peristome as in *B. ruralis*. Husnot, Musc. Gall. 115.—Montana; Washington.

154. *Barbula brachyangia* C. M. & Kindb.—Dioicous: plants brown ferruginous, 2-3 cm. high: leaves small, upper close, curved, short ovate-oblong, obtuse or subacute, upper part scarcely longer than sheathing base; margins slightly recurved in middle; broad cells hyaline, nearly uniform, costa rough at back; hair point long, denticulate, greater part pale red: capsule small and short, oblong cylindric, sub-symmetric; pedicel red. Mac. Cat. 59.—On rocks: Becscie River.

155. *Barbula faeviscula* Kindb.—Tufts brown, radiculose at base of leaves, 3-4 cm. high: leaves narrow lingulate, carinate, nearly smooth or indistinctly papillose at recurved borders; lower marginal cells of leaf-base hyaline, short rectangular, in 1-2 rows, median yellow and porose, in 2-3 rows, inner larger, longer and hyaline; costa red, indistinctly papillose at back, denticulate and hair line near long hispid hair-point: capsule sub-cylindric, attenuate at base to pedicel; peristome tube very much shorter than pale teeth.—Mac. Cat. 235.—On rocks: Rocky Mountains, B. C.

156. *Barbula papillinervis* C. M. & Kindb.—Allied to *B. ruralis*: upper leaves subacute, leaf-base narrowly margined, with pale yellow cells: calyptra very much prolonged below capsule. Mac. Cat. 60.—On rocks: British Columbia; Labrador.

157. *Barbula intermedia* Brid.—Stems 1-2 cm. high, erect, bifurcate, in compact olive green tufts, brownish below: leaves erect-spreading, not squarrose, oblong spatulate, rounded or emarginate at apex, slightly concave, not carinate, revolute on borders as far as middle; point shorter and less dentate than in *B. ruralis*: perichæstrial leaves acute: pedicel and capsule shorter; peristome shorter, teeth once twisted. Husnot, Musc. Gall. 116.—Montana; Washington.

158. *Barbula aciphylla* Bruch & Schimp.—Aspect of *B. ruralis*: leaves spreading squarrose when moist, erect contorted when dry, oblong lanceolate, very concave, revolute at border, acuminate, lamina prolonged on each side of nerve, which is extended into a reddish hair furnished with fewer teeth and more erect: flowers dioicous: perichæstrial leaves not folded: perigonal leaves not always ecostate, sometimes distinctly nerved: pedicel strongly twisted to left at base, and to right in upper part; capsule oval or oval-oblong; peristome shorter. Husnot, Musc. Gall. 115.—On rocks: Disco Island, Greenland; Gold Range, Selkirks, and Rocky Mountains, British Columbia.

159. *Barbula leptotricha* C. M. & Kindb.—Differs from *B. ruralis*

principally in shorter leaves, emarginate at apex, with a fine and nearly smooth hair point; basal cells hyaline. Mac. Cat. 60.—On limestone rocks: Manitoba.

160. *Barbula lato-excisa* C. M. & Kindb.—Differs from *B. leptotricha* in the leaves being green, nearly flat at borders; outer basal cells faintly chlorophyllose, narrowly marginate. Mac. Cat. 60.—On the bases of trees: Vancouver Island.

161. *Barbula rotundo-emarginata* C. M. & Kindb.—Dioicous: plants loosely tufted, brown, about 1 cm. high: stems beset with emarald leaves or naked below; comal leaves close, patent when moist, scarcely curved, papillose, short obcordate, not sheathing; margins slightly or not recurved; cells subquadrate, inner basal greater and hyaline; costa red and thick, faintly rough at back, excurrent into a long hyaline or at base reddish more or less denticulate hair-point. Mac. Cat. 60.—*Sine loco*.

162. *Barbula aloides* Bruch & Schimp.—Dioicous: short, dull green: leaves erect spreading, longer, rigid, linear lanceolate, acuminate, acute, nerve incrassate in middle, convex at back, often reddish: capsule from curving of the seta, cernuous or subhorizontal, cylindraceous, subarcuate, rufo-fuscous on upper side, pale fuscous beneath; calyptra reaching a little beyond the lid; annulus of small cells, long persistent, lid rostrate, acute; teeth of peristome very slender, pale red, simply contorted, when dry arcuate incurved with points assurgent; spores smooth. *Tortula aloides*, Braithw.—Brit. Moss Fl. 1: 211. Newfoundland.

163. *Scouleria aquatica nigrescens* Kindb.—Plants coarse and rigid, in large dense black tufts: leaves oblong lanceolate, blunt and entire at rounded apex, vein ending below it, often radiculose at base; margins serrate, basal cells rectangular next vein, prosenchymatous near margin but extending upward only a short distance; some leaves simply hyaline at base with all cells rectangular. Bull. Torr. Bot. Club 22: 42. 1895.—Vancouver Island; Sicamous and Roger's Pass, B. C.

164. *Scouleria marginata* Britt.—Plants 3-4 cm. high, gregarious, in dense black tufts: stems wiry and naked at base, branching and densely foliate above: leaves crowded, curled and twisted when dry, only uppermost green, oblong lingulate, serrate above middle, or obscurely serrulate near base, teeth occasionally black and thickened; apex blunt, entire or toothed; vein thick, ending below it, smooth on back; basal cells green, rectangular, a narrow band near margin elongated, prosenchymatous, forming a dark dense border nearly to apex of leaf, superposed by rounded small cells: perichaetial leaves surrounding capsules, ovate lanceolate: capsules small, broader than long, cupuliform when old; lid persistent on columella, bordered with red; peristome none; mouth bordered; spores green with a minutely roughened coat. Bull. Torr. Bot. Club 22: 42. 1895.—Spokane Falls, Wash.; California.

165. *Grimmia apocarpa alpicola* H. & T.—More densely cespitose, lower, suberect: leaves shorter, broader, muticous; costa short: capsule larger, emergent, operculum long rostrate. Schimp. Syn. Musc. Eur. 243. 1876.—Greenland; Alaska.

166. *Grimmia conferta pruinosa* Braith.—More robust, in blackish tufts: leaves broader, upper gradually ending in long smoothish hairs: perichaetial bracts larger, distinctly papillose: teeth of peristome rufous-orange, more lanceolate, often reflexed against capsule. Braithw. Brit. Moss Flora 2: 7.—Cape Horn Mountains, Idaho.

167. *Grimmia chloroblasta* Kindb.—Differs from *G. conferta* principally in long hair pointed leaves: perichaetial leaves larger and greener: lid of capsule short, conic apiculate; teeth very cribose, nearly as in *Coscinodon pulvinatus*. Mac. Cat. 64.—On dry rocks: Spence's Bridge, B. C.

168. *Grimmia heterophylla* Kindb.—Differs from *G. conferta* in stem nearly simple: leaves patent when dry, upper caniculate; margins involute above: perichaetial leaves very much longer: the peristome reddish, not papillose: tufts about 2 cm. high, when dry dark green above. Mac. Cat. 64.—On rocks: Spence's Bridge, B. C.

169. *Grimmia atricha* C. M. & Kindb.—Differs considerably from *G. conferta* in tufts densely cohering: leaves small and when dry appressed, muticous, short, ordinarily ovate-oblong, not recurved at the margins; cells not incrassate: perichaetial leaves very much larger and broader than the others: capsule more wide-mouthed, teeth of peristome orange only below, yellow or hyaline above, entire, not rimose nor papillose; lid longer rostrate. Mac. Cat. 65.—On rocks: Sproat, B. C.

170. *Grimmia pachyneurula* C. M. & Kindb.—Tufts small, green: leaves small, when dry appressed, not twisted, when moist spreading, short ovate lanceolate, recurved on both sides, hairless and obtuse; cells uniform, subquadrate, not erose, the alar scarcely distinct; costa percurrent, thick, smooth at back: barren. Mac. Cat. 65.—On rocks: Revelstoke, B. C.

171. *Grimmia Phillibertiana* Britt.—Dioicous: plants pulvinate, in small dark green cushions: stems naked and decumbent below, branching and spreading above; leaves erect incumbent when dry, not secund, spreading when moist, lanceolate, carinate, with recurved margins and toothed hair points, generally deformed and bearing globose propagula, or retuse and bifid along midvein; cells above rounded, hexagonal, faintly sinuous and oblong at base, discolored, with slight enlargement at basal angles; costa heavy, rounded at back, sulcate above: perichaetial leaves broader, inner short, triangular and hyaline at base: pedicels one or two from same perichæstium, twisted, variously bent; capsule broadest at mouth, smooth when dry, pale with a red rim; teeth recurved, red, undivided, segments broad below, slender and papillose above; lid straight or oblique; calyptra mitrate; annulus delicate.—Bull. Torr. Bot. Club 18: 51. 1891.

172. *Grimmia elatior* B. & S.—Dioicous: robust, 4-6 cm.: stems erect, naked at base, slightly branching, in broad lax tufts of a yellowish green color, brown in interior: leaves erect, lanceolate, concave carinate, borders revolute, hair point long and nearly smooth; basal cells rectangular, marginal scarcely distinct, upper round quadrate, more or less papillose; costa strong: perichaetial leaves large, erect, laxly areolate: seta arcuate; capsule oval, striate, when dry sulcate; lid conic, straight; annulus large, compound (3); teeth broadly subulate, purple, densely articulate, irregularly cleft and perforated. Braithw. Brit. Moss Flora 2: 23.—Godhavn, Greenland. On rocks: Rocky Mountains.

173. *Grimmia arcuatifolia* Kindb.—Loosely tufted, tufts blackish, dark green above: stems 5 cm. long, denudate at base: lower leaves small, upper long and not crisped, hooked-curved when moist, ovate lanceolate, long acuminate and acute, reflexed at least at one border, with a short denticulate hair point; most basal cells pellucid, long, narrow, upper basal cells sinuous, marginal uniseriate, hyaline or not distinct; other cells chlorophyllose, quadrate, not sinuous; costa percurrent, canaliculate, pellucid in the middle. Mac. Cat. 69.—On dry rocks: Vancouver Island.

174. *Grimmia Hartmani* Sch.—Dioicous: tufts large, lax, dense-leaved, yellowish or olive green above, dark at base: stems 3-10 cm., procumbent, naked at base: leaves often more or less secund, erect spreading when moist, slightly crispate when dry, oblong lanceolate, upper ending in a slightly denticulate hair, carinate, revolute at border, sometimes only on one side; lower marginal cells quadrate, distinct, cells near costa rectangular, upper roundish-quadrate; young terminal leaves tipped with globose propagula, formed of quite a number of cells, or occasionally bifurcate: perichaetial leaves shorter, from a longish concave base, lanceolate, pointed, with laxer more transparent cells: capsule longish oval-oblong, smooth; seta erect or curved; annulus of three to four rows of small cells; calyptra mitriform, lobed; teeth lanceolate, entire or slightly perforated, orange red, smooth at the base. Husnot, Musc. Gall. 135.—Summits of Chilco Range, Idaho.

175. *Grimmia deplata* Kindb.—Tufts large and compact, brown or green above: stem elongate: leaves when moist arcuate, ovate lanceolate, recurved on both sides, generally hairless and obtuse, sometimes with a short hair point; alar cells hyaline, elongate, in 4-5 rows, the others more or less erose: perichaetial leaves from a sheathing base narrowed into a large canaliculate sublinear and obtuse acumen: capsule nearly smooth, when dry finally rugose, not distinctly costate; teeth red, deeply cleft below middle, when dry spreading, connivent when moist: Leak more or less oblique; pedicel arcuate when dry. Mac. Cat. 69.—Vancouver Isl., alt. 3,000 feet.

176. *Grimmia Arizonae* Ren. & Card.—Differs from *G. trichophylla*

and *G. California* in broader leaf base, upper $\frac{2}{3}$ bistratose, more dense, more obscure, hair point longer, striate and coarsely denticulate; from *G. Olneyi* in more robust habit, longer hair point and incrassate inferior leaf cells. *Revue Bryol.* 19: 85. 1892.—Arizona.

177. *Grimmia prolifera* C. M. & Kindb.—Tufts soft, coherent and very radiculose, when dry blackish below, proliferous with long green shoots: stem slender, naked below, 3-4 cm. long: leaves when dry incurved or crisped, upper often falcate when moist, narrow at the oblong appressed subvaginant base attenuate to a short acute acumen, mucous or rarely furnished with an apical hair-point-like cell, margins recurved at least at one side; lower basal cells narrow rectangular, others and upper subquadrate, all thin walled, faintly yellowish and pellucid; costa yellow brown, percurrent.—Differs from allied *G. contorta* principally in the leaves shorter, mucous, upper more distant than lower. *Mac. Cat.* 67.—British Columbia.

178. *Grimmia tortifolia* Kindb.—Nearly allied to *G. torquata*: differs principally in shorter leaves and leaf cells. *Mac. Cat.* 68.—On rocks: Revelstoke, B. C.

179. *Grimmia funalis* Sch.—Dioicous: densely caespitose, greenish above, brown below: stems 1-5 cm., branching, eradiculose: leaves when dry erect and spirally incumbent, when moist erect spreading, small, lanceolate, upper with hyaline points or extended into a smooth hair, margin recurved, costa narrow, vanishing at apex; basal cells rectangular, upper quadrate: perichaetial leaves broader at base, concave, piliferous; seta arcuate, capsule small, oval, slightly striate; lid conic, beak short, annulus large, compound (3-4); teeth purple, lacunose or torn at apex into two papillose legs. *Husnot, Musc. Gall.* 132.—Smith's Sound, Greenland.

180. *Grimmia Hendersoni* Ren. & Card.—Closely related to *G. decipiens*: seta longer, capsule subcylindric and narrower, lid longer rostrate, basal areolation looser. *Revue Bryol.* 19: 86. 1892.—Oregon.

181. *Grimmia pachyphylla* Leiberg.—Mats wide, dense, inflated: stems 5-12 cm. high, repeatedly dichotomous, subsimple, plants intermixed, erect or ascending from a decumbent base, radiculose: leaves imbricate when dry, spreading when moist, upper portion more or less recurved, oblong or broadly lanceolate, shortly decurrent, carinate above and reflexed on margins, strongly costate, nerve more or less channeled, above becoming laminoid, gradually narrowing into a sparingly toothed hair; cells long rectangular below, subquadrate in middle, small quadrate above, all sinuous; margin of 2-3 rows of quadrate cells; extreme base and angles of irregular oblong or subrhombic cells: dioicous: capsule oval or oblong, pendent on a twisted seta, red, narrowed at mouth, collum distinct, substrumose, when dry erect and irregularly 4-8 costate; teeth 2-cleft to below middle, sometimes merely lacunose along middle line, papillose above, ar-

triose, more dense,
enticulate; from *G.*
crassate inferior leaf

soft, coherent and
as with long green
when dry incurved
in the oblong ap-
acumen, mucous or
margin recurved at
others and upper
lucid; costa yellow
principally in the
wer. Mac. Cat. 67.

G. torquata: differs
Cat. 68.—On rocks:

caespitose, greenish
ose: leaves when dry
ng, small, lanceolate,
air, margin recurved,
upper quadrate:
seta arcuate, cap-
anulus large, com-
two papillose legs.

related to *G. deci-*
lid longer rostrate,
regon.

ense, inflated: stems
plants intermixed,
se: leaves imbricate
r less recurved, ob-
above and reflexed
eiled, above becom-
ched hair; cells long
rate above, all sinu-
base and angles of
ule oval or oblong,
llum distinct, sub-
eth 2-cleft to below
papillose above, ar-

ticulations few; annulus compound (2). Bull. Torr. Bot. Club 20: 113. 1893.—Granite, gneissoid and slate rocks: Idaho.

182. *Grimmia cinellidodonta* C. Müll.—Monocous, male flower terminal on a special branch: branches fasciculate: leaves subsquarrose-spreading, loose, strict when moist, rather long and narrow, regularly concave, from an oblong base gradually attenuate, apex rather obtuse, margin quite entire, flat or a little curved; costa thick, occupying whole point; cells minute, round, their walls smooth, shining: perichaetial leaves larger: capsules short pedicellate, immersed, hemispherical, macrostome, exannulate, often aggregated; operculum obliquely rostellate; peristome teeth broad, lanceolate, intense red, broadly trabeculate, apex perforate and irregularly cleft. Bot. Centralbl. 44: 388. 1890.—On wet rocks: Ellensburg, Wash.

183. *Grimmia crassinervis* C. Müll.—Monocous; habit of *G. conferta* but leaves wholly pointless, slightly papillose; costa rather thick, occupying almost all the acumen: perichaetial leaves long acuminate from a broad base, quite entire: capsule immersed, pyriform-cyathiform, macrostome; seta short; operculum very obliquely rostrate; calyptra long persistent, cucullate, smooth; teeth lanceolate, rather short, somewhat perforate at apex. Bot. Centralbl. 44: 389. 1890.—Victoria, Vancouver Is.

184. *Grimmia tenella* C. Müll.—Dioicous: tufts small, dirty green: stem slender, fasciculate branched below, branches parallel, somewhat flexuous slender: leaves erect spreading, scarcely crispate, subulate, narrowly oblong-acuminate; hair point rather short, hyaline, straight or slightly flexuous, slender, sharp, sharply denticulate; margin quite entire, scarcely involute; costa rather broad for the leaf, excurrent; cells all chlorophyllose, thick walled, a few at base quadrate, toward apex round: perichaetial leaves larger, broader: seta slender, short; capsule scarcely surpassing leaves, erect, small, oblong, truncate, leptodermons, pale with age, smooth; lid minute, obliquely rostrate; teeth small, slender; calyptra narrow, cucullate. Bot. Centralbl. 44: 388. 1890.—Cœur d'Alene, Idaho.

184a. *Grimmia Mannia* C. Müll.—Dioicous: tufts slender, compressed, intensely green; stems fastigiate branched, very small, densely foliate, loosely coherent: stem leaves densely imbricate, small, from a broad truncate base deeply carinate, open, obtusely short acuminate; margin plane, quite entire; costa slender, green, excurrent; cells at base small, green, empty, hexagonal, rather thick walled, above minute and obscure: perichaetial leaves very closely imbricate, much larger, more obtuse, cells larger, more involute, all rather fleshy: seta long exerted, slender, reddish, somewhat spirally twisted; capsule erect, globose-urn-shaped, thick walled, rufous; lid conic; peristome teeth short, red, more or less split and perforate. Flora 70: 233. 1887.—Napa Springs, Calif.

185. *Grimmia elongata* Klif.—Dioicous: tufts lax, pulvinate, dark below, olive green and shortly canescent at apex: stems slender, elongated,

dichotomous, sparingly branched, naked and decumbent at base: leaves divergent from an erect oblong base, rather rigid, elongate-lanceolate, lightly recurved at margin, subcomplicate-carinate, mucicous, wings asymmetric; costa stout, excurrent, uppermost leaves with short hyaline points; basal cells linear-rectangular, dilated towards margin, upper quadrate, sinuose, apical minute: perichaetial leaves oblong lanceolate, acuminate with short hair points: capsule exerted on a short straight seta, small, ovate, smooth; lid conic, obtuse; annulus of 2-3 rows of cells; calyptra cucullate; teeth yellow, broad, cleft or perforate only at the apex. Braithw. Brit. Moss Flora 2: 30.—Greenland.

186. *Grimmia sulcata* Sauter.—Stems erect, branching, in olive green tufts: lower leaves mucicous and obtuse, upper oblong lanceolate, with a short hyaline point, bistratose in upper part, longitudinally trisulcate especially above the middle; lower cells short rectangular: dioicous: perichaetial leaves shorter and less sheathing than in *G. alpestris*: capsule exerted, oblong sub-cylindric; lid convex, beak a little longer than in *G. alpestris*; annulus persistent, scarcely distinct; teeth of peristome entire, red; calyptra cucullate, covering $\frac{1}{4}$ - $\frac{1}{6}$ the capsule. Husnot, Musc. Gall. 129.—Franz Joseph Fjord, Greenland.

187. *Grimmia microtricha* C. M. & Kindb.—Plants in small blackish pulvinate tufts with greenish tops: leaves when moist sub-erect, short ovate lanceolate, margins recurved, plane only at the short base; cells short, alar narrow and hyaline: perichaetial leaves ovate obtuse: capsule oblong, smooth; teeth dark red; pedicel flexuous: dioicous, or monoicous on distinct branches.—Differs from *G. alpestris* principally in longer emergent capsules, recurved leaf margins, and short perichaetial leaves. Mac. Cat. 70.—On rocks: Revelstoke, B. C.

188. *Grimmia tenerrima* Ren. & Card.—In small, compact, gray tufts: leaves small, oblong-lanceolate, lower mucicous or with a short hyaline point, upper prolonged into a smoothish hair; borders generally reflexed in the upper part; costa canaliculate; basilar cells lax, quadrate, pellucid, thin walled, upper bistratose, subquadrate, with scarcely thickened walls: capsule exerted on a short pale pedicel, small, leptodermous, smooth, yellow or pale brown: lid convex apiculate; peristome orange red, teeth patulous when dry, papillose, perforated, more or less lacerate at the apex; calyptra cucullate: male flowers unknown. Bot. Gaz. 15: 40. 1890.—Moist bluff towards the snow line: Mt. Hood, Oregon.

189. *Grimmia mollis* B. and S.—Dioicous: tufts lax, soft, dark green: plants robust, stem bifurcating: middle leaves erect spreading when moist, very soft, oval-lanceolate or oblong, obtusely rounded, concave cochleariform, plane on the border, unistratose; costa narrow, vanishing below the apex; lower cells rectangular, upper roundish-quadrate: perichaetial leaves acuminate, with hyaline points: capsule short exerted, not passing be-

yond the summit of the perichaetial leaves, elliptic; annulus simple; teeth of the peristome linear lanceolate, lacunose. Husnot, Musc. Gall. 127.—Greenland.

190. *Grimmia sarcocalyx* Kindb.—Differs from the allied *G. leucophæa* principally in leaves with a faintly denticulate hair point: capsule short oval; lid obliquely beaked; pedicel doubly longer than the capsule; vaginule pale red, inflated and fleshy.—Leaves short, very broad at base as in *G. leucophæa*; cells nearly all quadrate. Mac. Cat. 66.—On rocks: Spence's Bridge, B. C.

191. *Grimmia sphaerica* Schimp.—Stems erect, with inflated branches, tufts compact, grayish green: leaves erect, lower small, muticous; upper gradually larger, oval-lanceolate, concave, acute, slightly revolute at the borders, bistratose, apex hyaline, large and dentate above in the perichaetial leaves; lower cells rectangular, upper quadrate or rounded with thickened walls: seta straight, very short; capsule sub-globose, symmetric, enlarged at mouth; lid convex, small, apiculate; annulus compound (2), persistent; peristome very rudimentary, scarcely reaching beyond the annulus, so that capsule appears gymnostomous; calyptra mitriform. Husnot, Musc. Gall. 124.—Canada.

192. *Rhacomitrium proteasum* Braun.—Allied to *R. aciculare*: differs in stems longer, less denudate, branched many times, inclined, erect above, in depressed tufts of a yellowish green color: leaves crowded, when dry imbricate, when moist recurved and erect spreading, rarely secund, linear lanceolate from a long base, with obtuse entire apex, concave; margins revolute beyond middle or only on one side, unistratose; costa distinct, vanishing below apex; cells round-quadrate or short rectangular near apex, in the middle rectangular, and below linear, papillose, sinuous: inner perichaetial leaves shorter, plicate, sheathing: capsule erect, long to about cylindrical; lid rostrate, beak straight; annulus compound, 2-3 rows; teeth of peristome split to base into two unequal papillose free or occasionally united legs; spores yellowish brown, punctate. Husnot, Musc. Gall. 139.—On rocks: Victoria, Vancouver Island; Greenland.

193. *Rhacomitrium Macounii* Kindb.—Plants fastigiate branching, innovations without lateral fasciculate branchlets: tufts loosely cespitose, naked at base, brown, with green tips: leaves loose, crispate when dry, patent or squarrose when moist, ovate-lanceolate, acute, muticous, smooth and entire, at one side slightly reflexed or erect, on other always erect; upper cells quadrate and obscure, scarcely or not erose, lower linear and sinuose, marginal cells uniseriate, quadrate-rectangular, hyaline, basal cells yellow; costa brown, percurrent: capsule oblong, dark brown, not striate; teeth orange, pertuse or cleft to below middle, smooth; pedicel straight or sub-erect and contorquate. Mac. Cat. 73.—In large masses on boulders: Rocky and Selkirk Mts. and Cold Range, B. C.

194. *Rhacomitrium alternatum* C. M. & Kingb.—Allied to *R. Macounii*: stem lower, more branched above: tufts loose, not naked at base, brown with green tips: leaves not crisped, when moist sub-erect or sub-patent, nearly straight, ovate lanceolate, acuminate-acute, often furnished with a short dentate hair point, smooth, reflexed at the base at least on one side; all cells erose, lower linear, upper short angular; costa greenish, stout, percurrent: perigonal leaves sub-ovate or short acuminate, acute or sub-obtuse: female plants not found. Mac. Cat. 73.—On boulders: Selkirk Mountains, B. C.

195. *Rhacomitrium robustifolium* Kindb.—Differs from *R. Macounii* in leaves less crisped, pellucid very squarrose when moist, more reflexed on borders, often furnished with a short hair point: capsule oval, striate or plicate when dry; teeth dark purple brown, deeper cleft, papillose; beak oblique, needle shaped, very much shorter than capsule; pedicel slightly curved. Mac. Cat. 73.—On rocks: Lake Griffin, B. C.; Vancouver Island.

196. *Rhacomitrium obtusum* (Lindb.) R. & C.—Dioicous: short, densely pulvinate: leaves quite hairless, ovate oblong, gradually acuminate, obtuse at point, nerve lost far below apex, margin narrowly revolute: capsule oblong, narrowed at mouth; lid acicular; peristome fugacious, irregular, teeth with two unequal legs. Mac. Cat. 74.—On rocks: Lake Superior.

197. *Rhacomitrium heterostichum alopecurum* Hüb.—Dioicous: yellowish green, scarcely hoary: stem slender, elongate, fasciculate branched: leaves lanceolate acuminate, acute with a short or obsolete hair point: capsule elliptic-oblong, narrow at mouth; lid conic rostrate; teeth short, pale, cleft to base. Mac. Cat. 74.—On dry rocks: Halifax, N. S.; Selkirk Mountains and McLeod's Lake, B. C.

198. *Rhacomitrium heterostichum occidentale* R. & C.—Stems often nearly simple: pedicel very short; capsule small, pale, not shining; peristome pale. Bot. Gaz. 15: 41. 1890.—On rocks: Lost Lake, Oregon.

199. *Rhacomitrium micropus* Kindb.—Habit of *R. heterostichum* or *R. canescens*: plants dirty green, branches nodose with numerous short branchlets: leaves very faintly papillose, ovate-lanceolate, long-acuminate, more or less revolute, hair point long and rough; cells yellow, upper short or elongate, alar distinctly quadrate: capsule small, oblong-cylindric; beak short, oblique; pedicel short. Mac. Cat. 77.—On rocks: Gold Range, B. C.; Hector, N. W. T.

200. *Rhacomitrium microcarpum* Palmeri Kindb.—Leaves long subulate, hairless, upper cells longer and more confluent, alar large and rectangular: capsule shorter pedicellate. Differs from related *R. Sudeticum* in deeply cleft peristome teeth and narrow leaf cells. Mac. Cat. 287.—St. Paul Island, Behring Sea.

CONSIN.

—Allied to *R. Ma-*
 not naked at base,
 sub-erect or sub-
 ut, often furnished
 the base at least on
 ular; costa greenish,
 acuminate, acute or
 —On boulders: Sel-

fers from *R. Ma-*
 e when moist, more
 point: capsule oval,
 deeper cleft, papillose;
 an capsule; pedicel
 n, B. C.; Vancouver

—Dioicous: short,
 gradually acuminate,
 rowly revolute: cap-
 some fugacious, irreg-
 —On rocks: Lake

Hüb.—Dioicous: yel-
 asciculate branched:
 late hair point: cap-
 e; teeth short, pale,
 fax, N. S.; Selkirk

& C.—Stems often
 not shining; peri-
 Lake, Oregon.

heterostichum or
 with numerous short
 te, long-acuminate,
 yellow, upper short
 long-cylindric; beak
 cks: Gold Range,

db.—Leaves long
 ent, alar large and
 related *R. Sudetti-*
 cells. Mac. Cat.

201. *Rhacomitrium speciosum* C. Müll.—Dioicous: tufts very broad and lax, plants intricate, robust, green: stems dichotomously branched, elongate: leaves crowded, when wet quickly and very distinctly squarrose-recurved, dimorphous; the lower (older) ovate from a broad base, many times lightly plicate, rather long decurrent, acuminate, obtusate, cucullate; upper (younger) hair pointed, the hair stout, rather long, hyaline, denticulate-serrate; margin quite entire, below (up to middle or beyond) broadly revolute; costa broad, deeply canaliculate, excurrent; cells at decurrent angles loosely parenchymatous, basilar longer: perichaetia! leaves involute, erose truncate at apex: capsule cylindric, erect, narrowed below mouth, smooth, plicate with age; seta short, smooth, twisted; lid conic, long rostrate; calyptra long subulate, apex slightly rough, persistent, lacinate at base with 16 lobes; teeth very long and narrow, dirty red, smooth, bifid almost to base; annulus none. Bot. Centralbl. 44: 388. 1890.—Victoria, Vancouver Is.

202. *Rhacomitrium languinosum subimberbe* Hartman.—Tufts extended, stems long, erect, a little flexuous, delicate, slightly nodulose, branches distant, short; leaf point shorter, sometimes almost disappearing. Fl. Miq. 46.—Miquelon Is.

203. *Rhacomitrium canescens muticum* Kindb.—Leaves without a hair-point; cells yellow; costa percurrent: barren. Mac. Cat. 77.—Gold Range, B. C.

204. *Rhacomitrium canescens Delamarei* Ren. & Card.—Tufts yellowish, stems long, nodulose, branches short, erect, leaves erect when dry, rigid, often broken at point, without a hair, almost smooth, costa percurrent: sterile, but very distinct from var. *lutescens* Lesq. & James. Fl. Miq. 46.—In extended mats on earth: Miquelon Island.

205. *Hedwigia ciliata subnuda* Kindb.—Leaves nearly hairless, the greater number broadly ovate, borders reflexed, cells larger, subquadrate. Mac. Cat. 78.—On rocks in woods: Ottawa; also near Wooler, Ont.

206. *Zygodon* H. & T.—Plants dichotomous, fastigiately branched, crowded and interwoven with radicles: leaves spatulate lanceolate, very chlorophyllose above, hyaline at base: capsule on an elongated pedicel, oval-oblong, with a soft swelling neck, less distinctly striate; peristome single or double, rarely none, the teeth resembling those of *Orthotrichum*; calyptra smooth, cucullate.

207. *Zygodon viridissimus* Brid.—Dioicous; tufts small, cushion like, bright green above, brownish at base: stems sparingly branched, fastigate, brown radiculose at base: leaves dense, when moist recurved squarrose, when dry erect-appressed, or slightly twisted, toward apex complicate, oblong or linear-lanceolate, acutely acuminate, deeply carinate above, expanded below, minutely papillose, margins plane; costa pellucid, ending abruptly below apex; cells at base thin, quadrate hexagonal, above

small hexagonal rotundate: capsule erect, oval-oblong, neck short, olivaceous, when dry pyriform, obscurely 8-costate; lid obliquely rostrate; calyptra fugacious, rostrate; peristome none. Braithw. Brit. Moss Flora 2: 61.—On trees: White Falls, between Norway House and Hudson Bay at York Factory.

208. *Zygodon conoidens* H. & T. Dioicous; tufts lax, light yellow green: stems nearly simple or sparingly branched, slender, beset with ferruginous radicles at base: leaves less dense, patent, curving upward, imbricated when dry, linear-lanceolate, acuminate, flat and slightly keeled, more papillose, nerve narrow, vanishing below apex; cells larger and more incrassate: capsule oval, tapering into a neck of equal length, narrower, striate towards mouth when dry and empty, leptodermous, pale fuscous; lid subulately beaked; peristome of 8 bigeminate pale yellowish teeth, reflexed when dry, truncate at apex, fugacious, endostome of 8 yellowish cilia, fugacious, often abortive. Braithw. Brit. Moss Flora 2: 62.—Eastern States and Virginia.

209. *Drummondia clavellata Canadensis* Kindb.—Leaves larger and longer: inflorescence monoicous. Mac. Cat. 81.—On trees: Pelee Island, Lake Erie.

210. *Ulota megalospora* Vent.—Primary stems creeping, tomentose; tufts dense, bright green: all leaves when dry cirrhate-crispate; those of primary stem small (1 mm.), narrowly long subulate from a broadly ovate concave base; apex of one row of cells; base cochleariform, cells narrow, thick-walled; upper cells roundish angular, 7-8 μ broad, walls thickened, each with a thick round papilla; margin irregularly crenulate by projecting cell walls; upper leaves of branches broader with a shorter subula: perichaetial leaves longer, from a longer base, short subulate: autoicous sporophyte 5 mm. high: capsule small, ovate, when dry sulcate and urceolate, constricted below mouth; striæ 8, extending to middle or beyond, of 4 rows of cells, neck long, defluent; stomata superficial; external teeth 8, pale, bigeminate, minutely papillose, above almost smooth, lineolate, split along divisural line $\frac{1}{2}$, otherwise entire; cilia 8, subulate, smooth, below of 2 rows of cells; annulus double; operculum apiculate from a conic base; calyptra sparsely hairy; spores very large, 55-61 μ diam., green, minutely papillose, mixed with small spores scarcely 15 μ diam. Bot. Centralb. 44: 389. 1890.—Cascade Mts., Wash.

211. *Ulota maritima* C. M. & Kindb.—Differs from *Ulota phyllantha* in plants smaller, darker, green or black, not yellow: leaves shorter and less circinate when dry, long attenuate to subulate apex, distinctly papillose, costa narrower, rough at back: capsule short oval with short collum, pedicel shorter, thicker, curved when moist; peristome reflexed, teeth trabeculate, cilia long, carinate, of two rows of cells, finely granulate and striolate in both. Mac. Cat. 84. Bull. Torr. Bot. Club 21:

72. 1894.—On rocks: British Columbia; Vancouver; Alaska Behring Sea; Miquelon Island.

212. *Ulota Hutchinsiae rufescens* Britt.—Plants more slender than the species, green or brown, not black: stems rufous tomentose: leaves less crowded, longer, acuminate; cells more dense and obscure, lower golden brown, rectangular, not linear or sinuous, marginal shorter, not hyaline: capsule pyriform, inflated, narrower at mouth; peristome double, teeth reflexed when dry, white, granulose, not trabeculate at apex; cilia 8, of two rows of cells, fugacious. Bull. Torr. Bot. Club 12: 69. 1894.—On trees in dense woods.

213. *Orthotrichum Shawii* Wils.—Monoicous: tufts lax, 1-2 cm. high, brownish radiculose at the base, green, olive-green or brownish above: leaves loosely imbricate when dry, when moist reflexed and spreading from an erect base, lanceolate from an ovate base, long and narrowly pointed, margins more or less revolute; costa vanishing below apex, lamina unistratose; cells papillose, thick walled, round-hexagonal, above larger and oval, near base elongated rectangular, at angles shorter and broader: perichaetial leaves erect, less keeled, inner often smaller: capsule ovate, immersed, neck half length of capsule, gradually narrowed into seta, when dry and empty almost urnshaped, light weak folds above; lid short, bordered with orange colored cells; annulus persistent, 2-3 rows of cells; peristome simple, teeth 16, when dry reflexed, linear lanceolate, pale, split in middle here and there along middle line, thickly papillose. Limpr. Laubm. 1: 90. Bry. Eu. Suppl. *Orthotrichum*, pl. 1.—California.

214. *Orthotrichum fastigiatum* Bruch.—Closely allied to *O. affine*, but generally smaller, tufts only 1 cm. high: stem reddish radiculose, fasciculately branched: leaves more rigid, when dry appressed, when moist reflexed and erect spreading, shorter and broader, long lanceolate, mostly short pointed, keeled, plicate at base, margin revolute; costa percurrent; cells on both sides with simple or furcate papillae, the upper thick-walled, roundish, in point larger, oval or oblong, at base linear (1:6), more transparent, at angles much shorter: perichaetial leaves larger and broader, the inner with plane margins: capsule immersed, somewhat thicker, long pyriform, 8 costate; neck furrowed and gradually narrowed to seta; annulus of one or two rows of cells: peristome double, light yellow, teeth eight, when dry reflexed, trabeculate at apex, vermicular striate; cilia 8, robust, shorter than teeth. Limpr. Laubm. 1: 82.—Lake Superior.

215. *Orthotrichum Sprucei* Mont.—Autoicous: in dark green, sparingly branched tufts: leaves erect, loosely imbricate when dry, erect spreading when moist, ovate-spatulate, rounded at point and muticous or with a small apiculus; margin revolute at base, then recurved to some distance below summit; cells at base quadrangular, elongated near nerve, passing gradually into large, rounded or angular, smooth or faintly papillose

ones; costa slender, vanishing below apex: perichæstrial leaves longer, narrower, sulcate, with a short nerve; sometimes with a long filiform point: capsule immersed, oval-pyriform with a long sulcate neck, broadly 8-striate; annulus compound (2); teeth 8, bigeminate, yellowish, reflexed when dry, densely and finely papillose; cilia 8, shorter than teeth, sometimes 16; lid conic, rostellate. Braithw. Brit. Moss Flora 2: 81. 1889.—On willows: Clark's Fork of Columbia River.

216. *Orthotrichum urnigerum* Myrin.—Monoicous: tufts lax, bright green to yellowish brown, reddish-brown radiculose at base: stems prostrate or in thick tufts, erect, 2-5 cm. long: leaves when dry loosely appressed, when moist becoming recurved and falcate spreading, lower lax upper larger and tufted, lanceolate from an ovate base, long pointed, keeled; margins strongly revolute as far as middle; costa vanishing below apex; cells uniform throughout in size, roundish quadrate or hexagonal, thick walled, papillose on both sides with simple or furcate papillæ, short rectangular at leaf base: perichæstrial leaves somewhat larger, erect, longitudinally furrowed at the base: capsule half immersed, thick oval, 8-striate; neck short, when dry suddenly contracted into seta; lid short, annulus persistent, compound; peristome double; teeth 16, papillose below, vermicular and longitudinally striate above; cilia 16. Limpr. Laubm. 2: 46.—Yellowstone National Park.

217. *Orthotrichum Roellii* Vent.—Tufts pulvinate cespitose, closely radiculose: stems erect, branching: leaves from ovate lanceolate to lanceolate acuminate, cells below incrassate, rotund, papillose with simple or furcate papillæ; cells above quadrangular, elongated, smooth; margin revolute nearly to apex: autoicous: capsule cylindrical ovate, scarcely exerted, when dry faintly sulcate to middle, striæ of 2 short rows of cells, other cells quadrangular; stomata superficial; collum short, defluent into seta; peristome teeth 16, simple, approximate in pairs, when dry erect or spreading, above longitudinally striate, striæ below inclined, rarely mixed with papillæ; scarcely a vestige of cilia found, but when present shorter than teeth; spores 16-18 μ , minutely papillose. Bot. Centralbl. 44: 360. 1890. *O. lonchothecium* C. M. & Kindb., Mac. Cat. 90.—Rocks: Ellensburg, Washington; trees: Krao Creek, Kootenai Lake, B. C.; Banff, Rocky Mountains.

218. *Orthotrichum Schlotthaueri* Vent.—Tufts dense, pulvinate, fuscous-green: stems erect, branched, 1.5-3 cm. high: leaves when dry closely appressed, when moist apex quickly recurved, then erect spreading, lanceolate from an oblong base, acute, 2-3 mm. long; upper cells rotund-angular, walls thick, papillæ thick, often furcate; margin reflexed: autoicous: capsule immersed, elongate-ovate and cylindrical, smooth when dry, not constricted under the mouth, not sulcate, more or less long pedicellate from defluent collum, all together 3-4 mm. high; stomata superficial; annulus

aves longer, nar-
ong filiform point:
k, broadly 8-stri-
h, reflexed when
h, sometimes 16;
889.—On willows:

tufts lax, bright
base: stems pros-
en dry loosely ap-
reading, lower lax
se, long pointed,
a vanishing below
ate or hexagonal,
ate papillae, short
arger, erect, longi-
thick oval, 8-stri-
; lid short, annu-
papillose below,
pr. Laubm. 2: 46.—

cespitose, closely
nceolate to lanceo-
ee with simple or
ooth; margin revo-
scarcely exerted,
ows of cells, other
defluent into seta;
dry erect or spread-
rarely mixed with
esent shorter than
bl. 44: 360. 1890.
cks: Ellensburg,
; Banff, Rocky

se, pulvinate, fus-
s when dry closely
spreading, lanceo-
ls rotund-angular,
d: autocous: cap-
nen dry, not con-
pedicellate from
perforial; annulus

double or triple; teeth of the peristome eight, yellowish, each divided to the base into two legs slightly split at the apex and marked with a median line, when dry erect or spreading, distinctly articulate, more or less densely papillose, especially below middle; cilia more or less perfect, smooth, fugacious, sometimes wanting. Bot. Centralb. 44: 390. 1890.—Garrison and Sun River Cañon, Montana.

219. *Orthotrichum euryphyllum* Vent.—Loosely caespitose, 2-3 cm. high, dark green, rufescent: stem erect, branching: upper leaves 4 mm. long, 1.5+ mm. broad, lower smaller, lanceolate from a broadly ovate base, apex roundish, entire, or obtuse and slightly toothed, margin revolute nearly to apex, so that apex is sometimes cucullate; costa ending far below apex; cells hexagonal, 12-13 μ diam., walls not thickened, reddish, papillae single, minute, or wanting: autoicous: capsule immersed, thick, broadly ovate, when dry constricted under the mouth, deeply sulcate to the middle, striæ 8, broad, of 4-6 rows of cells; column short, abruptly narrowed into the pedicel; annulus double or triple; peristome double, teeth 16, reddish, approximate in pairs, when dry radially spreading, each interruptedly cleft almost to the base in the middle line, papillae minute, and arranged in more or less regular lines; cilia 16, eight robust, equaling teeth, papillose below, the intermediate eight rudimentary or abortive.—Bot. Centralb. 44: 417. 1890. On stones: Ellensburg, Wash- ington.

220. *Orthotrichum nudum* Dicks.—Monoicous: tufts lax, soft, green or dirty green, 1-2 cm. high, brown radiculose at base: leaves broader and softer than in *O. cupulatum*, obtuse, longitudinally furrowed at the base; cells quite uniform: perichaetial leaves longer and broader and somewhat sheathing at base: capsule emergent or exerted, larger and thicker than in *O. cupulatum*, pyriform with a long neck abruptly contracted into seta, with alternately long and short striæ, lid red margined, beak short; annulus compound (2-3), persistent; peristome double, teeth 16, when dry erect, longitudinally striate or slightly papillose; cilia often rudimentary, 8 or 16, scarcely half as high as the teeth.—Limpr. Laubm. 2: 42. *O. cupulatum nudum*, Braithw. Brit. Moss. Flora 2: 78. On lime- stone rocks: Rockcliffe, Ottawa river.

221. *Orthotrichum strictum* Vent.—Sub-species of *O. Lyellii*. Leaves without gemmæ, rigid, lanceolate; capsule with short seta; otherwise as in *O. Lyellii*. Bot. Centralb. 44: 419. 1890.—Cascades, Enumclaw, Wash- ington.

222. *Orthotrichum bullatum* O. Müll.—Monoicous: tufts lax, yellow- ish green: stems slender, dichotomously branched, fastigate, 2 cm. high, flexuose: leaves not crowded, crispate or secund-twisted, when moist re- curved from an erect base, rather broadly oblong, bluntly acuminate, quite entire, almost smooth; margin strongly revolute; here and there

plicatulate below, concave; costa thickish, carinate, glabrous, yellowish at base, vanishing below apex; cells small, elliptical, not thickened, larger and yellowish or orange at base: perichaetial leaves larger, secund at apex; capsule immersed, inflated-oval, thin, pale yellow, strongly bullate, slightly plicate; operculum flat, erect rostrate; peristome simple, teeth 16, solitary, cupulate or erect, never reflexed, pale, narrowly lanceolate; calyptra slightly plicate, pale, shining, with few hairs.—Flora 70: 223. 1887. California.

223. *Orthotrichum speciosum* Roellii Vent.—Densely pulvinate, blackish green: leaves nearly smooth, papillae very minute, cells large as in *O. Killiasii*; margin revolute: capsule emergent, oblong-ovate, smooth; peristome normal. Bot. Centralbl. 44: 419. 1890. Chicago, Argyle, Ill.

224. *Orthotrichum elegans* Schwagr.—Tufts dark green, 1-2 cm. high, softer and smaller than in *O. speciosum*: leaves lanceolate acuminate, recurved at border; cells at base colorless, thin walled, quadrangular, elongated, passing gradually into rounded-hexagonal cells above, with one or two small simple papillae: monoicous: capsule small, thin walled, sub-cylindric, green and smooth before emptying, afterwards pale and slightly costate in upper part; collum distinct, more or less elongated; peristome double, 8 bigeminate teeth densely papillose, pale yellow, when dry reflexed against capsule but not revolute; cilia 8, filiform or linear, of rows of cells, papillose, convergent but not touching at points; operculum conic, apiculate, border pale red; hood conic-campanulate. Husnot, Musc. Gall. 169.—On ledges at Lake Pend d'Oreille, Idaho, to Kootenay River, B. C.

225. *Orthotrichum Killiasii* C. Müll.—Closely related to *O. speciosum* but generally smaller, tufts pulvinate or sometimes flat, rigid, dirty green: stem erect, branched, densely foliate: leaves appressed, when moist erect-spreading, narrow lanceolate, obtuse, younger leaves acute; margin revolute nearly to apex; cells thick walled, densely papillose with simple and furcate papillae, above roundish or oval, at the base yellowish red, rectangular, towards margin quadrate: inner perichaetial leaves larger, longitudinally furrowed at base: capsule immersed or emergent, long cylindric, with short 8-costate neck, when empty narrow cylindric, not furrowed and somewhat constricted at mouth; lid red margined, beak equal to diameter of base; annulus simple or compound (1-2); peristome double, teeth when dry erect, originally united in pairs but soon separated, densely papillose; cilia 8 (according to Schimper; to Venturi 16) robust, papillose, margins sinuous. Limpr. Laubm. 2: 92.—Disco Is., Greenland.

226. *Orthotrichum præmorsum* Vent.—Densely pulvinate, 2-3 cm. high, bright yellowish green, below reddish yellow: leaves long acuminate from an ovate base, often apiculate, when dry often curved into a cone, when moist becoming suddenly recurved, then erect spreading,

costa ending in apex or apiculus: margin narrowly revolute; cells ovate or rotund above, chlorophyllose, walls thickened below (especially near nerve), elongated, narrow walls irregularly thickened, papillæ in upper part of leaf salient, simple or furcate: inflorescence autoicous: capsule emergent, ovate, collum equaling it in length, defluent into seta, when dry slightly constricted under mouth and sulcate, striæ 8, indistinct; stomata immersed; annulus simple; peristome double, teeth pale yellow, 8, when dry reflexed against wall of capsule apex truncate and fimbriate, remainder entire, or rarely apex lacunose, papillæ minute, distinct; cilia irregular, 8, fugacious, sometimes fragmentary; spores 15-18 μ . Bot. Centrabl. 44: 418. 1890.—Yellowstone National Park, Wyoming.

227. *Orthotrichum rhabdophorum* Vent.—Densely pulvinate, dark green: stems erect, somewhat branched, 1.5-2 cm. long, below tomentose radiculose: leaves when dry imbricate, when moist becoming suddenly recurved, then erect spreading, lanceolate, or lanceolate from an ovate base, acuminate, margin strongly revolute nearly to apex; cells above rotundate, walls thickened, papillæ bi- or tri-furcate, salient; inflorescence autoicous: perichaetial leaves for most part reaching middle of capsule, sometimes capsule entirely exerted: capsule when dry and operculate ovate-cylindrical, with faint reddish striæ, when moist ovate-elongate, constricted below mouth, collum short, passing abruptly into seta, old capsules sulcate at least to middle; annulus double or triple; peristome simple or rarely double, teeth 16, when dry recurved, sparsely papillose; cilia if present more or less perfect, erect, spores 8-11-14 μ . Bot. Centrabl. 44: 418. 1890.—Cascades, Thorp, Washington.

228. *Orthotrichum arcticum* Schpr.—Tufts dense, more or less depressed, dark green or blackish, 1-2 cm. long; stems branched, rigid: leaves densely imbricate when dry, when moist becoming recurved and then erect-spreading, obovate or oval-lanceolate and more or less pointed, recurved at margin and revolute in lower half; costa vanishing below apex in lower leaves and almost at point in upper leaves: cells at base quadrangular, smooth and with unequally thickened walls, passing gradually above into relatively large rounded or subhexagonal cells with thick walls and provided with salient bi- or tri-furcate papillæ: perichaetial leaves larger at base, more or less acuminate and a little longer than others: inflorescence autoicous: capsule emergent and sometimes exerted, oval or oval oblong, when dry sub-urceolate and faintly furrowed; collum straight and abruptly contracted; annulus triple; peristome double, teeth 8, bigeminate, split to middle and more or less lacunose along the middle line, finely and densely papillose, reflexed when dry; cilia often fragmentary, not passing half height of teeth, of one or two rows of cells. Husnot, Musc. Gall. 172.—Greenland.

228a. *Orthotrichum Macounii* Aust.—Autoicous: robust, in broad

dense yellowish tufts fuscous below: stem 1 cm. long, sparingly branched: leaves when dry erect, when moist rather strict sub-patent, ovate lanceolate, very acute, sub-carinate, minutely papillose, apex entire, margin revolute; costa sub-percurrent; cells very minute, obscure, basal a little broader, scarcely more pellucid: capsule very narrow cylindrical, smooth, pale straw-yellow, long exerted, defluent when dry, into a long a strongly sulcate collum; peristome teeth 16, short, subulate, when dry erect-incurved, pale, hyaline, distinctly 8-10-articulate, minutely granulose papillose; cilia 8, half shorter than teeth, narrow; lid long apiculate; calyptra sparsely hairy. Bull. Torr. Bot. Club 6: 343. 1879.—Rocks: Cascades, B. C.; Washington; Idaho.

229. *Orthotrichum Blyttii* Schpr.—Tufts dense, wide, 1-3 cm. high, brownish or olive-colored: leaves erect spreading, imbricate when dry, recurved when moist, obovate-lanceolate, revolute at border almost to summit; cells at base quadrangular, smooth, passing gradually into rounded or sub-hexagonal cells above, with thickened walls, papillae simple or bi- or tri-furcate, sometimes salient but variable, older leaves not papillose: perichaetial leaves a little larger at base: capsule emergent, oval or oval-oblong, when dry 8 costate; collum straight and suddenly contracted at base, making fruit more or less pyriform; annulus persistent, compound (3-4); peristome double, teeth 8, split more or less along middle line, finely and densely papillose, reflexed when dry; cilia 8, more or less complete, sometimes very small or half height of teeth, of 1-2 series of cells. Husnot, Musc. Gall. 174.—Greenland.

230. *Orthotrichum pumilum Americanum* Vent.—Tufts small, lax, soft: leaves almost without papillae: capsule with 8 bands composed of 2 series of rectangular cells faintly marked; teeth 8, obtuse at point, split along middle line, finely papillose. Husnot, Musc. Gall. 190.—On trees, locality unknown.

231. *Orthotrichum Rogeri* Brid.—Tufts lax and irregular, 1-1.5 cm. long, green or dark green: leaves more or less flexuous when dry or loosely imbricate, when moist erect-spreading or only upper lanceolate half of leaf spreading, carinate, lanceolate from oblong base sometimes concave or even cochleariform, frequently rounded at point, obtuse, acuminate or short apiculate, entire on border or denticulate near apex; border more or less revolute; costa ceasing below apex; cells elongated quadrangular at base, smooth, walls slightly thickened, passing gradually into rounded cells with thickened walls above, papillae very small, simple, rarely more pronounced: inflorescence autoicous: capsule more or less emergent, costate when dry and empty, contracted below mouth before empty, oval oblong with collum slightly defluent into seta, sometimes longer than sporangium; annulus double, persistent; peristome double, teeth 8, bigeminate, reflexed when dry, yellowish orange or darker, more or less split or lacunose at points,

densely and finely papillose or slightly lineolate above; cilia 8, weakly papillose or lineolate, of 1 or 2 series of cells. Husnot, Musc. Gall. 186.—Idaho.

282. *Orthotrichum Hendersoni* Ren. & Card.—Pulvinate, yellow green; stems dichotomous, 1-2 cm. long; leaves patulous, flexuose when moist, slightly crispate when dry, from an oblong base linear lanceolate, acuminate, carinate, borders strongly revolute; costa vanishing below apex; cells thick walled, elongated, sub-rectangular below, in upper part: roundish or angular, papillose capsule subexserted on a short pedicel, oval oblong, suddenly constricted to pedicel, 8 striate, becoming cylindrical and contracted below mouth when old and empty; lid convex, apiculate; teeth 8, bigeminate, yellow, minutely granulose, not striolate lengthwise, reflexed when dry, split at apex; cilia 8, smooth; spores papillose. Bot. Gaz. 15: 42. 1890.—On bushes: Coast Mts., Oregon.

283. *Orthotrichum pulchellum productipes* Ren. & Card.—Much more robust than type, with larger leaves, a longer pedicel, and teeth of peristome larger and paler. Bot. Gaz. 15: 43. 1890.—On trees and shrubs: Portland, Oregon.

284. *Orthotrichum pulchellum leucodon* Vent.—Tufts caespitose, bright or pale green; peristome pale, becoming white.—Bot. Centralbl. 44: 419. 1890. Vancouver Is., Washington.

285. *Orthotrichum ulotaeforme* Ren. & Card.—Pulvinate, yellow green; stems dichotomous, 1-2 cm. long; leaves patulous, flexuose when moist, slightly crispate when dry, carinate; borders strongly revolute, sometimes sinuate at apex; costa vanishing below apex; cells thick walled, lower elongated, narrow, sub-sinuuous, upper roundish or sub-hexagonal, slightly papillose; capsule exserted on a long pedicel, oblong, 8 striate when dry, suddenly contracted to pedicel; lid depressed, rostrate; teeth 8, bigeminate, or 16 more or less connected in pairs, pale yellow, minutely granulose, striolate lengthwise, truncate and split at apex, reflexed when dry; cilia 16, long, nodulose, nearly smooth; spores papillose; inflorescence monoicous. Bot. Gaz. 15: 42. 1890.—On bushes with *O. Hendersoni*: Coast Mts., Oregon.

286. *Eucalypta subspathulata* C. M. & Kindb.—Monoicous; stem very short, about 0.5 cm. high; leaves bright green, spathulate or lingulate, twisting when dry, upper spreading when moist; inner basal cells short rectangular, smooth and hyaline, finally red-brown, outer ones much longer, narrower, and yellowish; costa faintly reddish below, yellow above, long excurrent; capsule cylindrical, short necked, smooth or finally furrowed when dry; peristome pale, partly incomplete and membranous, but distinct and high; calyptra small, papillose in narrower part, yellowish green, not covering whole capsule, not fringed; pedicel red. Mac. Cat. 83.—On rocks, Frazer River, B. C.; on earth: McLeod's Lake, B. C.

237. *Encalypta lelemitra* Kindb.—Nearly allied to *E. rhabdocarpa*, but leaves shorter, often subspathulate, costa vanishing at apex; peristome teeth nearly blunt; calyptra not papillose; spores larger. Mac. Cat. 94.—On rocks: Clearwater River, Athabasca.

238. *Encalypta cucullata* C. M. & Kindb.—Monoicous: leaves cucullate, perichaetial ones long acuminate acute, with a long hair point; costa percurrent, red at base; calyptra papillose all around. Mac. Cat. 96.—On earth: Columbia river, near Revelstoke, B. C.

239. *Encalypta lelocarpa* Kindb.—Monoicous: stem 3-4 cm. high, dichotomously branched, radiculose; leaves erect-patent, lingulate, faintly revolute nearly all around, without a hair point; lower decolorate brown, sub-acute, incurved; comal larger, green, obtuse, slightly twisted; basal cells hyaline, marginal very papillose; costa faintly papillose, not excurrent, in lower leaves brown, in comal green or reddish at base; perigonal leaves with a short thick tip; capsule straight, smooth, cylindrical with an apophysis; peristome simple, orange; pedicel red; calyptra papillose all around, not fringed. Mac. Cat. 95.—Crevices of rocks: summit of Mount Queest, B. C.

240. *Encalypta Alaskana* Kindb.—Differs from *E. longipes* Mitt. principally in capsule striate and not apophysate, peristome brown red, appressed to mouth when moist, costa nearly smooth; monoicous: lid of capsule obliquely rostrate. Mac. Cat. 260.—Mixed with a *Bryum* on earth: Ounalaska Island, Behring Sea.

241. *Encalypta apophysata* N. & H.—Stems 5-20 mm. high, erect, branching, in compact dark green tufts; leaves erect spreading when moist, crispate when dry, undulate, lanceolate elongate, apiculate by excurrent costa, revolute at base; seta rough at base, yellow above; capsule cylindrical, thin-walled, smooth; collum thick and distinct; lid conic, long beaked; teeth of peristome long, linear, entire or lacunose along middle line, articulations quite numerous, orange, papillose; hood irregularly lobed and lacinate at base; spores papillose. Husnot, Musc. Gall. 196.—Rocky Mountains.

242. *Merceya latifolia* Kindb.—Densely caespitose: plants 1-2 cm. high, brown-ferruginous below, green at tips, divided, at base radiculose; leaves quite smooth, spathulate-lingulate, obtusate or subacute, entire, slightly reflexed at base, plane above, broad-bordered with larger, orange-colored cells; basal cells narrow, upper small and round; costa percurrent or scarcely excurrent. Habit of *Barbula ruralis*. Mac. Cat. 97.—On upper slopes of Mount Finlayson near Goldstream, Vancouver Island, California.

243. *Tayloria acuminata* Hsch.—Monoicous: tufts more lax and generally shorter than in *T. splachnoides*; stem reddish tomentose, with numerous gemmae; leaves soft, loosely appressed or spreading, with reflexed point, when dry falcate, rhombic-lanceolate, long pointed; margins

of lower half of leaf reflexed and entire, above plane and toothed, apex not concave; costa longer and vanishing in point; cells somewhat smaller: seta 1-1.5 cm. high, capsule erect, smaller, oval or elliptic, when dry almost globose, suddenly contracted into the long thin neck; columella generally only slightly excurrent; lid suddenly obliquely beaked; peristome inserted deeply, separated by a layer of epidermis, teeth 16, hygrosopic, when moist involute, when dry appressed against the capsule wall, or somewhat irregular, warty-papillose on the outside, cross walls quite prominent. *Limpr. Laubm.* 2: 151.—On damp rocks: Dry Cañon, near Devils Lake, Rocky Mountains.

244. *Edipodium* Schw.—Plants small: leaves succulent, upper cells rounded quadrate, chlorophyllose: some flowers bisexual, others male: seta thick; capsule sub-spherical with a long collum; columella included; peristome absent; hood conic, cucullate; spores large.

244a. *Edipodium Griffithianum* Schw.—Stems 5-15 mm. high, erect, in soft, dark green tufts: leaves succulent, forming a rosette at summit of stem, obovate-epatulate, large and rounded at summit, entire, ciliate at base; costate to below apex; lower cells rectangular, hyaline; upper rounded-quadrate, chlorophyllose: seta pale green, thick, passing gradually into the very long collum; capsule sub-spherical, orange; lid convex-conic or shortly apiculate; peristome absent, hood fugacious, conic-cucullate; spores large, papillose. *Husnot, Musc. Gall.* 201.—Greenland.

245. *Physcomitrium turbinatum* (Michx.) Brid.—Plants light green, gregarious, 8-20 mm. high; autolcous: stems short and simple, or taller and branching: leaves oblanceolate or obovate from an oblong base, serrate above middle; costa vanishing below apex or occasionally excurrent into an acuminate apex; lower cells oblong, upper rhomboidal or hexagonal, marginal longer and narrower, often yellow and inflated at their upper ends: seta erect or twisted and occasionally arcuate: capsule erect, globose pyriform when fresh, becoming turbinate and contracted below mouth and spore sac when dry, dark brown and often urceolate when empty; lid convex or mamillate, occasionally apiculate when dry, blunt; mouth bordered by 8-12 rows of cells and a narrow row of orange colored cells, with a hyaline vesicular persistent annulus incurved after falling of lid; calyptra cucullate, oblique and split unequally, 5-8 lobed and beaked; spores rough.—*Bull. Torr. Bot. Club* 21: 199. 1894. A common but variable species in old fields, grassy open places in gardens, etc., from Florida to Ontario, west to the Rocky Mountains and California (?).

246. *Physcomitrium turbinatum Langloisii* (R. & C.) Britt.—Plants pale yellow; stems usually short and simple, slender, occasionally tall and branching to 2 cm.: leaves narrow, acuminate: seta filiform; capsule small, almost campanulate, not contracted below mouth when dry: neck tapering,

often contracted below spore sac when dry. Bull. Torr. Bot. Club 21: 200. 1894.—Low swampy ground and in gardens, probably only in the Southern states.

247. *Physcomitrium turbinatum* Floridanum (R. & C.).—Leaves longer, long acuminate, coarsely serrate: capsule strongly dilated at mouth, when empty cup shaped, varying. Bull. Torr. Bot. Club 21: 200. 1894.—Florida.

248. *Physcomitrium megalocarpum* Kindb.—The largest one of the genus, plants often 3-4 cm. high, light yellow or brown when old: stems short, simple: leaves spreading, flat and open when dry, not much twisted or shriveled, lanceolate from a lax oblong base; lower cells large, inflated at the angles, marginal longer and narrower in two rows, yellow, entire or serrulate; costa ending in acute or acuminate apex: seta erect or twisted and bent; capsule large, globose pyriform, nearly as broad as long, usually urceolate when dry, contracted at neck and below mouth when dry; lid conic, bluntly apiculate; mouth small, not flaring, bordered by a narrow orange-colored annulus with a second hyaline row and 8-12 rows of denser but slightly elongated cells; neck short, stomatose; spores rusty brown, spinose. Bull. Torr. Bot. Club 21: 200. 1894.—Pacific slope.

249. *Physcomitrium Kellermani* Britt.—Autoicous, antheridia terminal in basal buds: plants scattered or gregarious, dark brown when mature, small, seldom more than 3-5 mm. high: stems simple, with basal innovations: leaves few, rosulate, ovate acuminate; costa excurrent into subulate apex or ending below it; margins coarsely serrate; cells inflated, basal lax, scarcely elongated: seta short, scarcely exceeding perichæcial leaves, occasionally long and exserted; capsule short pyriform or broadly flaring, bright brown when old; neck short, tapering or swollen into an hypophysis, stomatose, rugose; mouth bordered with 4-7 rows of cells and a narrow, persistent annulus of darker cells with a hyaline incurved row almost invisible; lid small, conic rostrate; calyptra large, three lobed; spores large, rough, warty, but not spinose. Bull. Torr. Bot. Club 21: 204. 1894.—Kansas; Nebraska.

250. *Physcomitrium Coloradense* Britt.—Autoicous, antheridia in basal buds, few, large, without paraphyses; plants small, 3-4 mm. high, scattered or gregarious; stems short, simple or with basal buds; leaves few, radical, erect, concave, base short, auriculate; cells lax, upper cells shorter, marginal serrate or inflated above middle, with large irregular teeth, occasionally entire or serrulate only at apex; vein narrow, percurrent into a cuspidate apex or ending below it in lower leaves: seta short, immersed or partly exserted; capsule exserted, large for size of plants, nearly 2 mm. long, pyriform when fresh, becoming turbinate and contracted below mouth and spore sac when dry, bright orange or brown when mature; mouth bordered by 4-5 rows of narrow, elongated cells, and a

double annulas, the outer orange colored, inner vesicular, hyaline; lid large, conic, rostrate when dry, also bordered with orange cells; spores warty, not spinose. Bull. Torr. Bot. Club 21: 206. 1894.—Colorado; on muddy banks of the Missouri River, Great Falls, Montana.

251. *Physcomitrium Drummondii* Britt.—Plants gregarious or scattered, 3-10 mm. high; stems with short basal branches; leaves narrow, strictly erect, lanceolate-acuminate, serrate above middle, marginal cells longer and broader, inflated or yellow, basal cells lax; vein thick, ending below the apex or excurrent into a cuspidate point; seta stout, straw-colored or brown when old, short, erect, slightly twisted; capsules pyriform turbinate, not contracted below flaring mouth when dry, bordered by 6-8 rows of narrow elongated thick brown cells very distinct from cells with sinuous walls of rest of capsule; annulus narrow, orange-colored, persistent, with a second incurved hyaline row; lid conic-rostrate, bordered by orange, beak as long as spore sac, which is shallow and broad; neck contracted below spore sac, stomatose; spores large, warty. Bull. Torr. Bot. Club 21: 205. 1894. *Physcomitrium acuminatum* L. & J. Man. 198. 1884 in part.—Louisiana; Canada; Missouri; Oregon.

252. *Physcomitrium australe* Britt.—Autoicous, antheridia terminal and cotemporaneous with fruiting axis: plants gregarious, tall, slender, 3-4 cm. high; stems branching repeatedly and rooting at joints: lower leaves short, distant, with vein ending below the apex, upper crowded around base of seta, all narrow, oblong lanceolate, serrate above middle, marginal cells narrower and longer, teeth small, appressed; vein ending below acute apex: seta short, pale, twisted and curved; capsules often cernuous, subglobose becoming turbinate and flaring at mouth when dry and empty, small, often broader than long; neck tapering, contracted and stomatose; lid flat and apiculate when dry, conic when moist, bordered with brown; mouth bordered by 5-8 rows of large clear cells and a darker annulus, with a second row of hyaline cells incurved and almost invisible, surface cells irregular with thick walls; spores brown, rough, warty. Bull. Torr. Bot. Club 21: 201, 1894.—Apalachicola, Fla.

253. *Physcomitrium Californicum* Britt.—Autoicous, antheridia terminal, becoming lateral by innovations: plants gregarious, 10-15 mm. high; leaves few, basal, oblong-lanceolate, bordered by a double row of elongated cells, entire or occasionally serrulate above middle; vein thick, ending below acute apex; cells lax, oblong, marginal obliquely septate: seta slender, twisted, often brown; capsules small, globose when mature and empty, more or less cylindrical when young; lid conic, short, blunt; neck short, wrinkled, and contracted below spore sac when dry, stomatose; mouth bordered by a narrow orange-colored annulus and 5-11 rows of slightly denser scarcely differentiated cells; calyptra cucullate, lobed and

long beaked; spores brown, warty, not spinose. Bull. Torr. Bot. Club 21: 203. 1894.—On the ground: California.

254. *Physcomitrium acuminatum* (Schleich.) Br. & Sch.¹ Autoicous, antheridia terminal on basal branches: plants gregarious, pale green, slender, 10-15 mm. high: stems short, leaves almost radical, rosulate, oblong-lanceolate, acuminate; vein thick, ending below apex or excurrent into a cuspidate point; marginal cells elongated, in two rows, entire or subserulate at apex, lower cells elongated, often brown at angles: seta pale, slender, twisted, often bent; capsules small, pyriform, becoming turbinate when dry, with a broad flaring mouth, not contracted below it; neck tapering, often abruptly contracted when dry; mouth bordered by 4-7 rows of narrow cells, but slightly elongated or thickened, those of walls smaller than in *P. turbinatum* and rounded; annulus double, outer row orange colored, inner hyaline, vesicular, falling in fragments with lid or persistent; lid blunt, conic or apiculate, bordered with orange; spores small, yellow, spinose. Bull. Torr. Bot. Club 21: 203. 1894. L. & J. Man. 198 in part.—Not common: Central states west to Nebraska.

255. *Funaria calcaria occidentalis* Ren. & Card.—Differs from the type in the leaves more shortly and broadly acuminate, and the longer pedicel. Bot. Gaz. 15: 43. 1890.—Wet mud banks: Oregon City, Oregon.

256. *Bartramia breviseta* Lindb.—Tufts 1.5-3 cm. high, very thick, brownish green above, brownish tomentose below: stem dichasially branched, densely foliate: leaves obovate, half sheathing, appressed, pale yellow green, from a slightly shining base gradually narrowed into an erect-spreading narrow somewhat twisted brittle very sharp subulate point, margin plane, acumen toothed; cells of the sheathing part of leaf smooth, elongated rectangular, towards the margins narrower and colored, cells of the acumen much smaller, rectangular to quadrate, sharply mucillose in the corners; costa percurrent, filling the entire acumen: perichaetial leaves more sheathing, mostly exceeding the capsule in length: seta rigid, thick, brown; capsule large, erect, symmetric, almost globular, brownish, longitudinally furrowed, thin walled, small mouthed, when empty wide mouthed; lid small, convex, in the middle somewhat elevated; annulus none; peristome none or simple and rudimentary, formed of pale short obtuse irregular teeth. Limpr. Laubm. 2: 538.—Crevices of rocks: Ounalaska, Behring Sea.

257. *Bartramia glaucoviridis* C. M. & Kindb.—Differs from *B. pomiformis* in its glaucous green color, sheathing leaves not margined, suddenly short cuspidate, costa often excurrent: monoicous. Mac. Cat. 105.—On damp rocks: Columbia River, Revelstoke, B. C.

¹This description is given here because that in L. & J. Manual 198 is incomplete.

258. *Bartramia circinnatula* C. M. & Kindb.—Also allied to *B. pomiformis* but still more distinct than *B. glaucoviridis*: leaves green, loosely disposed, circinate, long sheathing, very long cuspidate: costa longer excurrent: probably dioicous. Mac. Cat. 105.—On rocks: Hastings, Burrard Inlet, B. C.

259. *Philonotis fontana cespitosa* Sch.—Stems usually simple, leaves more distant than those of *P. fontana*, more or less secund, oval, short acuminate, toothed, plane on border, not plicate: perigonal leaves broader than long, triangular, subulate, costa reaching apex. Husnot, Musc. Gall. 269.—Indiana; Illinois.

260. *Philonotis fontana microblasta* C. M. & Kindb.—Male flowers very small, brown; perigonal leaves all acute, strongly costate. Mac. Cat. 107.—On damp earth: Rogers Pass, Selkirk Mts., B. C.

261. *Philonotis fontana Columbia* Kindb.—Lower leaves narrow, costa long excurrent, perigonal leaves acute or subacute. Mac. Cat. 107.—On wet rocks: Revelstoke, B. C.

P. fontana brachyphylla Kindb., Prince Edward Is., and *P. fontana serrata* Kindb., Selkirk Mts. and Islands of Behring Sea, are *nomina nuda*. Mac. Cat. 107.

262. *Philonotis serrata* Mitt.—Dioicous: in stiffish yellow-green tufts with rufous tomentum, readily falling asunder, and with habit of *P. fontana*: leaves in spiral rows, imbricated when dry, erect or sub-falcate, dimorphous, those of male innovations ovate, bluntish, nerve vanishing, the rest deltoid ovate, acutely pointed: nerve thick, reaching apex or vanishing, all somewhat decurrent, concave, with two deep plaits at base on each side, margin revolute in lower third, bluntly toothed by single or double papillae; nerve very strong, rough at back, reaching to apex; cells above small and rectangular, with a papilla at lower and often at upper end, below laxer, oval, and longish with a central papilla on both sides: perichaetial leaves with nerve excurrent: capsule on a long straight seta, cernuous striate and furrowed; lid conic obtuse; peristome rufous, endostome orange, finely papillose. Braithw. Brit. Moss Flora 2: 212.—Washington.

263. *Philonotis glabriuscula* Kindb.—Tufts radiculose below, 4 cm. high: stem slender: leaves small, green, distant, spreading and straight when moist, indistinctly decurrent, short ovate-lanceolate, short acuminate, acute, slightly papillose, pellucid, plane at margins, not plicate, minutely serrulate principally above; cells oblong hexagonal, the lower narrower, less chlorophyllose; costa sub-percurrent: barren. Mac. Cat. 107.—By springs; Canaan Forks, N. B.

264. *Mielichhoferia cuspidifera* Kindb.—Differs from *M. nitida* in leaves broad-ovate, suddenly cuspidate, entire or slightly crenulate above, cells a little wider and costa often percurrent. Mac. Cat. 110.—On damp rocks: Hector, Rocky Mountains.

265. *Webera Cardoti* Ren.—Loosely cespitose, pale green or yellowish: stems simple, erect, slender, rigid and brittle: leaves small, erect, imbricate, oblong-lanceolate, decurrent, strongly revolute on borders, obtuse or sub-obtuse, rarely sub-acute, generally sinuate denticulate at apex; costa very broad, green, percurrent or vanishing very near apex, widening below and occupying one-third of base; cells lax, truncate or sub-attenuate, 2-3 times longer than broad: seta reddish, flexuous, often geniculate at base; capsule oblong-sub-pyriform, symmetric, erect, yellowish or brownish, tapering to an attenuate neck; lid conic; peristome small, pale yellow, pellucid, very minutely papillose; teeth triangular-lanceolate, long acuminate, lamellæ 15-20; endostome more or less perfect, with split segments and cilia, or reduced to a variously raised and lacerate membrane; annulus compound (2-3). Bot. Gaz. 14: 95. 1889.—On wet sandy rill-banks: Mt. Hood, Oregon.

266. *Webera polymorphoides* Kindb.—Tufts large, dense, dull green above, rufescent below, 4-5 cm. high: stem finally denudate at base, radiculose in middle: leaves crowded, not decurrent, ovate-oblong, acute or sub-obtuse, widely areolate, nearly entire and flat on borders, costa vanishing below apex; comal ones longer, sub-linear-lanceolate, short-acuminate, upper cells narrow, borders reflexed for greater part, costa thick, sub-percurrent: capsule oblong, short-necked, pendent; peristome pale yellow, segments with narrow basal membrane, cilia short, rudimentary; annulus narrow; lid mamillate; spores brown: dioicous. Mac. Cat. 111.—Hermit Mountain, Rogers Pass, Selkirk Mountains, B. C.

267. *Webera cruda minor* Ren. & Card.—Much smaller, capsule narrower, lid conic. Bot. Gaz. 15: 43. 1890.—Oregon.

268. *Webera longibracteata* (Broth.) R. & C.—Dioicous: loosely cespitose, light glaucous green, shining: stem about 2 cm. high, reddish, delicate, erect, flexuous, simple, somewhat brownish radiculose near base, laxly foliate: leaves spreading, long decurrent, nearly plane, linear-lanceolate, short acuminate, acute, margin slightly revolute from base to middle, or nearly plane, serrate from apex to middle, not bordered; costa pale vanishing below apex; all cells elongated, narrow, scarcely chlorophyllose, smooth: barren. *Pohlia longibracteata* Broth. Bot. Centralbl. 44: 419. 1890.—Argillaceous earth: Astoria, Oregon.

269. *Webera nutans subdenticulata* B. & S.—Stem simple: stem leaves narrower, longer pointed, plainly toothed; branch leaves ovate-lanceolate and loosely imbricate: capsule pendent, shortened, color uniform. Limpr. Laubm. 2: 251.—Miquelon Island.

270. *Webera nutans macrospora* Kindb.—Leaves denticulate to the middle; costa excurrent; spores large. Mac. Cat. 113.—Summit of Gold Range, B. C.

271. *Webera canaliculata* C. M. & Kindb.—Allied to *W. nutans*:

median and comal leaves longer attenuate, denticulate sometimes below middle, narrowly areolate, costa thick, canaliculate and excurrent, comal revolute at borders, only lowest shorter and short-decurrent; peristome pale; lid low and flat. Mac. Cat. 113.—On rocks: Vesuvius Bay, Salt Spring Island, Gulf of Georgia, B. C.

272. *Webera microcaulon* C. M. & Kindb.—Resembling a small form of *W. polymorpha* in the very short stem and acute leaves agglomerate in small buds: differs principally in dioicous inflorescence and very large spores, when unripe about 0.03 mm. Capsules (not ripe) short obovate, annulate; neck short; lid low mamillate; pedicel straight, arcuate at apex: comal leaves scarcely revolute at borders, inner perichaetial much shorter: tufts very compact; leaves green or finally blackish. Mac. Cat. 114.—Digges Island, Hudson Strait.

273. *Webera subcucullata* C. M. & Kindb.—Habit of *Mielichhoferia nitida*: intermediate between *W. cucullata* and *W. pycnoderrens*: resembles the last in small compact tufts, small (unripe) short-necked capsule, and mamillate lid; but stems subjulacous, leaves dull green, densely crowded, not decurrent; lower leaves short, sub-obtuse, nearly as in *W. cucullata* but leaf cells narrower. Mac. Cat. 113.—Crevices of rocks: Mount Queest, Gold Range, B. C.

274. *Webera Ludwigii* Sch.—Dioicous: soft, laxly caespitose, red or blackish at the base, dark green above, scarcely glossy when dry: stem and branches purple, erect, slender, sparingly radicles: lower leaves remote, broadly ovate, obtuse, entire, more crowded upward, erect spreading, long decurrent, not carinate; comal leaves densely crowded, oblong-lanceolate, serrulate at apex; margin narrowly recurved, nerve purple, vanishing below apex, thick at base; cells rather lax, thin, rhombo-hexagonal above, more rectangular at base: capsule on a flexuose reddish seta suddenly bent below capsule, sub-pendulous, oval-pyriform, brownish, annulate, slightly constricted below mouth; lid conical, obtuse, apiculate; peristome pale yellow, teeth linear-lanceolate, basal membrane of endostome reaching middle of teeth, processes gaping at keel, cilia 2-3. Brit. Moss Flora 2: 154 as Pohlia.—Oregon; Cascade Mts. and Gold Range, B. C.; Greenland.

275. *Webera gracilis* De Not.—Dioicous: much more slender than *W. commutata*, in loose, yellow green, glossy tufts, blackish at base, with many rigid filiform shoots, short in fertile plants, elongated and often with red axillary gemmae in sterile: leaves erect, appressed when dry, rigid, ovate and ovate-lanceolate, short-pointed, without chlorophyll; margin plane, faintly serrate at apex; nerve thinner, lost at or below point: capsule turgidly ovate, cernuous, small, reddish brown; lid orange, hemispherical, apiculate. Braithw. Brit. Moss Flora 2: 154.—Oregon.

276. *Webera micro-denticulata* C. M. & Kindb.—Tufts dense, glossy

n or yellowish:
l, erect, imbric-
ders, obtuse or
at apex; costa
widening below
attenuate, 2-3
ulate at base;
brownish, tap-
yellow, pellu-
ng acuminate,
segments and
prane; annulus
rill-banks: Mt.

nse, dull green
at base, radic-
g, acute or sub-
costa vanishing
acuminate, up-
ck, sub-percur-
ale yellow, seg-
mentary; annulus
t. 111.—Hermit
er, capsule nar-

ous: loosely ces-
h, reddish, delt-
lose near base,
, linear-lanceo-
base to middle,
red; costa pale
chlorophyllose,
atrabl. 44: 419.

ole: stem leaves
-lanceolate and
form. Limpr.

iculate to the
mmit of Gold

W. nutans:

green, about 3 cm. high: leaves small, loose when dry, open erect, decurrent, short, ovate-oblong, nearly entire, more widely areolate with a red and not percurrent costa; comal ones longer, lanceolate, acute or sub-obtuse, narrow areolate, revolute at borders nearly all around, faintly denticulate above, costa pale and sub-percurrent: capsules (not ripe), small obovate, short-necked; lid low, mamillate: dioicous. Mac. Cat. 114.—Close to perpetual snow on Gold Range, B. C.

277. *Webera pycno-decurrrens* C. M. & Kindb.—Tufts dense, glossy, bright green, 1-1.5 cm. high: leaves small, acute; lower ovate-oblong, crowded but short decurrent; comal very much longer sublanceolate, revolute at borders for greater part, denticulate above; inner perichastial leaves very much shorter, looser areolate; costa not excurrent: capsule obovate, red-brown, distinctly short-necked, annulate, orange-margined at the mouth; cilia sometimes appendiculate, inner membrane broad, teeth finally dark yellow; lid convex, orange-margined, mamillate; pedicel geniculate at middle: dioicous. Mac. Cat. 114.—On earth near perpetual snow on the Gold Range, B. C.

278. *Webera camptotrachela* Ren. & Card.—Stems erect, slender, simple or with few branches: leaves little crowded, erect, narrowly oblong lanceolate; acuminate, acute; borders plane or slightly revolute below, distinctly denticulate in upper part; costa strong, percurrent; cells elongated, sub-hexagonal or rhomboidal, 6-10 times longer than broad: external perichastial leaves more elongated, long narrowed-acuminate, more or less revolute on borders, serrulate with costa generally excurrent, 2 or 3 inner bracts smaller and shorter: seta reddish, flexuous, often geniculate at base: capsule small, subhorizontal or cernuous, oblong sub-pyriform, tawny-brown, with a long attenuated curved collum; lid convex, apiculate; annulus double; teeth yellowish, densely trabeculate; segments of endostome generally imperfect; cilia variable in length. Bot. Gaz. 18: 199. 1888.—California.

279. *Webera Columbia* Kindb.—Differs from *W. pulchella* principally in leaves more denticulate at least in the middle, costa red, annulus detached, revoluble. Leaves small, slightly reflexed, not glossy: capsule small, segments with 2 cilia. Mac. Cat. 115.—Moist banks: British Columbia and North West Territory.

280. *Webera albicans urceolata* Ren. & Card.—Capsule very short. Revue Bryol. 20: 1. 1893.—Oregon.

281. *Webera micro-apiculata* C. M. & Kindb.—Tufts small, dense and shining, bright green above, decolorate below, about 2 cm. high: leaves small, narrow areolate, densely imbricate and appressed when dry, when moistened sub-erect, not decurrent, ovate-lanceolate with a needle-shaped often incurved point, lower nearly entire, comal a little longer,

faintly and distantly sinuolate-denticulate above, borders not revolute, costa not excurrent: barren. Mac. Cat. 115.—Damp rocks: Revelstoke, B. C.; Hector, Rocky Mountains.

282. *Bryum Froudel* Kindb.—Habit of *Webera nutans*. Agrees with *Bryum inclinatum* in synoicous inflorescence, symmetric capsule, etc.; differs in leaves long acuminate, cells long and narrow, upper sub-linear, costa very long excurrent, peristomial segments quite free from teeth, spores smaller, cilia wanting. Mac. Cat. 120.—St. Paul Island, Behring Sea.

283. *Bryum sub-purpurascens* Kindb.—Agrees with *B. purpurascens* in red tufts, elongate shoots, costa scarcely or faintly excurrent, capsule long-necked, constricted below mouth, teeth orange colored, segments free, cilia smooth, spores small, lid large and mamillate, pedicel red: differs in inflorescence dioicous, all leaves red margined, costa long excurrent, and principally in curved sub-clavate capsule, resembling that of *B. meeseoides*. Mac. Cat. 119.—On wet earth: Port Moody, B. C.

284. *Bryum angustirete* Kindb.—Differs from *B. pendulum* in leaves narrow, ovate-lanceolate, reflexed all around; upper cells very narrow, sub-linear, basal cells reddish; costa red: capsule sub-cylindric; pedicel arcuate above; teeth paler; spores small: synoicous. Mac. Cat. 119.—Damp earth: Rocky Mountains; Revelstoke, B. C.

285. *Bryum Roellii* Philib.—Polygamous: yellowish green: leaves aggregated on upper part of stem, ovate, or elongate-lanceolate, acuminate, almost perfectly entire; cells small; costa long excurrent into a rigid scarcely denticulate point; margin slightly thickened, rather distinct, not colored, in lower part sub-plane, above broadly reflexed: seta 2-4 cm. long; capsule oblong, about 3.5 mm. long, becoming pale; lid conic, often darker colored; annulus broad, pale; peristome from a red base very pale, internal closely adherent to outer throughout its whole length, segments irregular, laterally affixed to teeth or obsolete, cilia none; dorsal lamina of teeth very slender, scarcely visible; ventral lamina pale, divided and excavate, with the vertical dissepiments of the internal membrane adnate in 3-4 rows of cells; spores 20-25 μ . Revue Bryol. 17: 56. 1890.—Cascades, Washington.

286. *Bryum brachyneuron* Kindb.—Agrees with *B. pendulum* in inflorescence synoicous, peristome orange, spores large: differs in leaves decurrent, short ovate, costa broad, abbreviate, not excurrent, shoots bearing globose gemmæ, peristomial teeth very much broader: stem red, very short: pedicel 1 cm. long or shorter, often scarcely emerging from tufts: costa of lowest leaves red, percurrent only in leaves of shoots and perichatral ones: capsules ventricose, short-necked, constricted below mouth. Mac. Cat. 120.—St. Paul Island, Behring Sea.

287. *Bryum Archangelicum* Schimp.—Synoicous, also with male and

more rarely with female flowers; tufts low, thick, pale green, reddish radicle within; leaves not decurrent; lower ovate-lanceolate, comal leaves loosely imbricate, lanceolate from an ovate base, concave, narrowly margined, generally somewhat revolute, more rarely plane; costa strong, excurrent into a long yellow faintly toothed hair; cells above rhombic, below rectangular, at the insertion red, and quadrate or rectangular: seta arcuate above; capsule nodding or pendent, obovate, not constricted under mouth; peristome yellowish, pale above, narrowly bordered, inner peristome same height, free or slightly adherent, cilia none or rudimentary.—*Limpr. Laubm.* 2: 308.— On earth: Gaspé Co., Quebec; on damp rocks: Hector, Rocky Mountains; Sabine Island, Greenland.

288. *Bryum mamillatum* Lindb.—Autoicous: tufts very thick and low: lower leaves small and distant, comal leaves crowded, not decurrent, long lanceolate, pointed, margin with a thick yellow border; costa very thick, excurrent into a short faintly toothed point; cells thin walled, rhombic above: capsule pendent, symmetric, globose-pyriform, neck short, rarely somewhat curved, longitudinally furrowed when dry; teeth of peristome orange, insertion red, yellow above, bordered; inner peristome free, yellow, basal membrane one-third length of teeth, cilia short, three.—*Limpr. Laubm.* 2: 328.— Greenland.

289. *Bryum Labradorense* Philib.—Plants branching, in compact tufts, radicle: leaves pale green, reddish at the base, oval-lanceolate, with a large sheathing base, decurrent, acuminate with a short point formed by costa, which is flexuose and somewhat toothed; otherwise margin entire; lower leaves plane and nearly emarginate; upper leaves faintly marginate with two rows of elongated cells; border slightly reflexed at base; cells distinct, compact, rhomboidal: polygamous: capsule oval, narrowed slightly at the base to form a short collum; lid convex with a scarcely salient beak; teeth of peristome very short, regularly attenuate, nearly triangular; inner peristome adherent and imperfect, segments linear, lacunose between articulations; cilia none; annulus very large; spores very large. *Rev. Bryol.* 14: 55. 1887.— Labrador.

290. *Bryum stenotrichum* C. Müll.—Synoicous: tufts slender, low, yellowish: fertile stems short, surrounded by a few slender very short rosulate stipitate innovations: perichaetial leaves erect-imbricate, spreading when moist, small, concave, from a broad-ovate purpurascens base longish acuminate; costa yellowish, thick, percurrent into an elongate slender scarcely dentate sharp point; margin strongly revolute from base to plane point, quite entire, broadly yellowish bordered; lamina confluent with subula; cells regular, small, yellowish, scarcely granulose: stem leaves smaller, shorter aristate: seta slender, pale red, 1 inch long, cernuous arcuate above; capsule small, from a slender neck narrowly oblong, ochraceous brown; lid small, conic, short pointed; annulus broad, revolvable;

peristome small, outer teeth short, densely trabeculate, inner ones slender, short, sulcate, split to the short shallow sulcus, short cuspidate, cilia very short, single, rudimentary. Flora 70: 219. 1887.—Alaska.

291. *Bryum Edwardsianum* C. & M. Kindb.—Nearly allied to *B. Warnum*, agreeing in peculiar peristome and very large spores (about .05 mm.), but differing principally in leaves longer acuminate, entire, revolute at borders; costa long excurrent; capsule narrower and lid lower; flagelliform branches absent: monolecous. Mac. Cat. 120.—On damp sandy soil: Prince Edward Island.

292. *Bryum Knowltoni* Barnes.—Plants densely caespitose, interwoven with red and brown rhizoids: stems copiously branched by innovations, reddish: leaves closely imbricated in bud like tufts at top of innovations, not twisted when dry, youngest bright green, older dirty yellow, carinate, concave, ovate to obovate-lanceolate, lower shorter, upper narrower, all abruptly and shortly acuminate; costa shortly excurrent, or dissolving in or ceasing below apex; margin entire, or rarely slightly denticulate here and there, slightly revolute or plane, border usually indistinct; cells rectangular and hyaline below, rhomboidal and densely chlorophyllous above: polygamous: capsule red brown or paler, rugose, pendent, oblong-pyriform; operculum small, strongly convex, apiculate, long persistent; annulus triple, revoluble; teeth linear lanceolate, strongly barred within, smooth above; segments of endostome free, strongly nodose, split between along keel, cilia two, rudimentary. Bot. Gaz. 14. 44. 1889.—Crevices in rocks: Funk Island, Newfoundland.

293. *Bryum fallax* Milde.—Stems 5-10 mm., branching: tufts yellowish: leaves erect spreading, oval, decurrent, concave carinate, very shortly mucronate by excurrent costa, entire, marginate, revolute at border, upper cells hexagonal: dioicous: capsule pendent, symmetric or arcuate, oblong pyriform, contracted at mouth; collum as long as sporangium; lid conic, apiculate; annulus large; teeth yellow, segments of endostome split along keel, cilia rudimentary. Husnot, Musc. Gall. 238.—St. Matthew Island, Behring Sea.

294. *Bryum caesium* Blytt.—Dioicous: tufts thick, olive color and reddish green, when old almost copper colored, reddish radiculose within: leaves spreading, rigid, when dry falcate incurved and somewhat twisted, narrowed at base and decurrent, lower leaves small and distant, ovate or obovate, sharp pointed, upper leaves oblong lanceolate or broad lanceolate, long pointed, red marginate; margin revolute, entire; costa thick, red, generally excurrent; cells thick-walled, pitted, above rhombic-hexagonal, the basal rectangular: perichatium leaves lanceolate, much smaller, rigid: capsule pendent or nodding, club pyriform, somewhat curved, reddish brown; neck one-half sporangium, when dry furrowed; lid small, yellow, convex; annulus compound (2); teeth narrow,

linear lanceolate, yellow, orange at insertion, yellowish above, broad hyaline margined, finely punctate; endostome adherent, yellow, papillose, segments narrow, free, split, cilia 2-3, broad, short. *Limpr. Laubm. 2: 332.*—Greenland: Smith Sound, Clavering and Sabine Island.

295. *Bryum mamilligerum* Kindb.—Subspecies of *B. intermedium*, differing in leaves distinctly margined, nearly flat on borders: capsule oblique, distinctly constricted below mouth, often horizontally patent; lid mamillate, not apiculate; spores larger: stem very short. *Mac. Cat. 122.*—On damp rocks: Devils Lake, Rocky Mountains.

296. *Bryum cirrhatum megalosporum* Kindb.—Differs principally in large spores, about .03 mm. *Mac. Cat. 122.* On wet soil on rocks: Burrard Inlet, B. C.

297. *Bryum cuspidatum* Sch.—Synoicous: tufts low, rarely 2 cm. high, dense, radiculose within: leaves decurrent, narrower than in *B. binum* and longer pointed; lower leaves small, oval, short pointed, upper larger, oval-lanceolate, pointed by excurrent costa, uppermost crowded in a coma, and over twice as large, with a very long acumen; margin recurved, yellow marginate, entire; costa excurrent as a toothed acumen; cells thin walled, faintly pitted, below rectangular, red, at decurrent angles somewhat elongated: capsule inclined to almost pendent, obovate, when deperculate constricted below the mouth, lid short conic; peristome and endostome of equal length, teeth gradually narrowed, papillose, broadly bordered; endostome free, pale yellow, segments half length of teeth, split along keel; cilia three, appendiculate. *Limpr. Laubm. 2: 343.*—Illinois; Wisconsin; Idaho; Montana; Oregon; Washington; Vancouver Island.

298. *Bryum binum angustifolium* Kindb.—Leaves narrow, loosely disposed: pedicel shortly emerging above innovations: stem 4 cm. high. *Mac. Cat. 123.*—On wet rocks: Cape Vincent, Ont.

299. *Bryum binum atrotheca* Ren. & Card.—Capsule black red: leaves scarcely denticulate or quite entire at the point. *Bot. Gaz. 19: 238. 1894.*—Newfoundland.

300. *Bryum leucolomatum* C. M. & Kindb.—Nearly allied to *B. binum*; differs principally in stem more robust and elevate, above 8 cm. high: leaves pale bordered, inner perichæcial ones cuspidate; capsules not ripe, lid apiculate: synoicous. *Mac. Cat. 123.*—In a marsh: Revelstoke, B. C.

301. *Bryum Hendersoni* Ren. & Card.—In robust, yellowish green tufts: stems robust, purple, tomentose, erect, dichotomous, 2-4 cm. long, lower leaves distant, smaller, then becoming gradually larger, upper crowded, erect spreading when moist, loosely appressed when dry, concave, cucullate at apex, broadly obovate-lanceolate, or oblong sub-spatulate, short acuminate and reflexed apiculate by excurrent costa; generally dentic-

ulate above on the back by the prominence of cell-apices; margin narrowly revolute but flat toward point, strongly serrate above; cells reddish and rectangular at base, oblong-hexagonal in middle, ovate-hexagonal or rhomboidal in upper part, the marginal elongated, linear flexuose, forming a more or less distinct border; capsule inclined or pendulous, narrowly cylindrical, incurved, constricted below mouth and tapering to a long attenuate neck; lid convex or sub-conic, apiculate; teeth yellow, densely trabeculate, segments split, cilia 1-3, appendiculate; annulus of 3-4 rows of cells, very broad; dioicous. Bot. Gaz. 19: 44. 1894.—Moist sunny bluffs: Portland, Oregon; California.

302. *Bryum microstegium* Sch.—Synoicous: aspect of *B. sub-rotundum*: tufts very low, thick, green: lower leaves small, distant, oval-lanceolate, costa ending below the point; comal leaves numerous, crowded, almost imbricate, when moist erect spreading, outer oval-lanceolate; inner larger, oblong lanceolate, long acuminate, costa excurrent, marginate, border plane, entire, sometimes slightly recurved; cells thin walled, above narrowly rhombic, at base violet purple, rectangular: seta twisted, curved above; capsule nodding, with narrower and shorter neck, oblong-pyriform, mouth symmetric, small; when dry neck furrowed and capsule rugose; lid conic, apiculate; teeth faintly bordered, weakly papillose, abruptly narrowed above; endostome almost hyaline, segments split along the keel cilia three, appendiculate. Limpr. Laubm. 2: 348.—Greenland; Labrador.

303. *Bryum pallescens laxifolium* Kindb.—Leaves loose, not glossy, long and narrow; upper cells narrow: spores small. Mac. Cat. 124.—In damp woods: Kananaskis Pass, Rocky Mts.

304. *Bryum pallescens longifolium* Kindb.—Leaves dense, glossy, very long attenuate; upper cells narrow: spores small. *l. c.*—Wet gravelly soil: Morley, Rocky Mountains.

305. *Bryum nitidulum* Lindb.—Synoicous: tufts low and dense, above yellowish green, below reddish, within densely brown radiculose: stem scarcely branched, leaves gradually larger above, broad oval, acute, narrowly marginate, margin revolute to recurved point, entire; costa thick, nearly percurrent; cells small, thick walled, pitted, upper irregular rhombic, basal rectangular, reddish: capsule pendent, small, oval-pyriform, gradually narrowed into neck, large-mouthed, pale yellow, shining; lid convex, with long point; annulus triple; peristome large, teeth pale yellow, lanceolate-acuminate; endostome free, basal membrane one-half height of teeth, segments split along keel, cilia 2-3, appendiculate. Limpr. Laubm. 2: 347.—Greenland.

306. *Bryum teres* Lindb.—Autoicous: tufts low, yellowish green, shining: leaves erect, not decurrent, broad oval, obtuse, the upper with small points, concave, not marginate, entire; lower leaves with plane margins,

comal leaves revolute to middle, inner to apex; costa purple at base, vanishing below point; cells faintly pitted, above rhombic-hexagonal, basal red, rectangular hexagonal; seta curved above; capsule pendent, symmetric, pyriform, when dry contracted below the mouth; lid small, convex; annulus triple, separating in fragments; teeth lanceolate, abruptly subulate above the middle, narrowly bordered, papillose, endostome free, basal membrane $\frac{1}{2}$ height of teeth, segments widely gaping along keel, cilia 2-3, long appendiculate. *Limpr. Laubm.* 2: 367.— Franz Joseph Fjord and Sabine Island.

307. *Bryum microglobum* C. M. & Kindb.—Tufts compact, green, radiculose below, small (about 1 cm. high); branches very short, bearing small buds; leaves narrow margined, sub-entire; stem leaves ovate oblong, short acuminate, faintly revolute at borders, branch leaves ovate acute and not revolute; upper cells short except narrow margined ones; costa more or less short excurrent; capsule small, inclined, pale globose-pyriform, small mouthed; peristome perfect, teeth pale; cilia short, appendiculate; lid convex with conical mamilla; pedicel arcuate at apex; diolcous. *Mac. Cat.* 129.— On earth: London, Ont.

308. *Bryum micro-erythrocarpum* C. M. & Kindb.—Nearly allied to *B. erythrocarpum*; differs in stem shorter; leaves distinctly yellow-margined, laxer areolate; costa sometimes excurrent; capsule more ventricose, constricted below the mouth; lid longer apiculate. *Mac. Cat.* 124.— In wet gravel: Vancouver Island.

309. *Bryum Blindii* B. & S.—Diolcous; gregarious or in small dense tufts, brownish or light green, somewhat shining; stem fasciculately branched, radiculose below; lower leaves broad oval, short-pointed; upper stem and branch leaves ovate, short-pointed; comal and perichæstrial leaves oblong-lanceolate; all leaves imbricate, concave, not margined, entire, margin plane, only upper comal and perichæstrial leaves slightly revolute; costa strong, percurrent or excurrent; cells yellow-walled, considerably thickened, above rhomboid and rhomboidal, at base red, thin walled, rectangular and rectangular hexagonal; seta generally bent at base, hooked above; capsule pendent, symmetric, obovate or globose-pyriform; neck thick; mouth small, when empty scarcely changed; lid small, strongly convex, with small mamilla; annulus triple, separating spirally; peristome inserted below mouth, teeth yellow, with pale incurved point when dry, bordered, papillose; endostome yellow, papillose, basal membrane $\frac{1}{2}$ height of teeth, segments lanceolate, suddenly subulate, somewhat split, cilia three, nodose or appendiculate. *Limpr. Laubm.* 2: 419.— On damp earth Rocky Mountains; on débris: Selkirk Mountains, B. C.

310. *Bryum alpiniforme* Kindb.—Allied to *B. alpinum* in habit. leaves chlorophyllose, basal cells quadrate, costa red; differs in leaves smaller, marginate, loosely disposed, more distinctly decurrent, border re-

flexed nearly all around, cells wider, upper hexagonal oval, costa often more excurrent: barren. Mac. Cat. 271.—On rocks: islands in Lake Nepigon, Ont.

311. *Bryum hæmatocarpum* C. M. & Kindb.—Tufts very tomentose below green innovations: leaves loosely appressed, when dry slightly corrugate but not twisted, crowded, not decurrent, subovate, acute, pellucid, narrow marginate, revolute at the entire borders all around, those of the innovations less distinctly revolute and margined to above the middle; cells pale yellow, upper short-rhomboidal; costa yellow, percurrent or in uppermost leaves slightly excurrent: capsule large, sub-oblong-cylindric, blood-red, twice as long as pale collum, pendent or patent, not appressed to pedicel; peristome dark yellow or orange below, pale above; segments free, very much shorter than membrane, cilia faintly appendiculate or rather nodose; annulus broad; lid nearly flat and apiculate: dioicous. Mac. Cat. 125.—On damp rocks: British Columbia.

312. *Bryum percurrentinerve* Kindb.—Tufts compact, dull green above, decolorate below: leaves when dry appressed and corrugate, when moistened sub-patent, crowded and slightly decurrent, ovate-obtusate, only uppermost ovate oblong and sub-acute, narrow margined, revolute at the entire borders for the greatest part; cells pellucid, upper wide, sub-rhomboidal; costa red, percurrent, in uppermost leaves pale. Allied to *B. Muhlenbeckii*, but differing principally in color, thinner, broader and shorter leaves, at the apex patent or sometimes reflexed, not cucullate. Mac. Cat. 216. Bull. Torr. Bot. Club 17: 274.—On dripping rocks: waterfall near Kamloops, B. C.

313. *Bryum capitellatum* C. M. & Kindb.—Tufts small, compact, yellowish green above, reddish rufescent below: stems sub-julaceous, clavate acute, radiculose at the base; innovations short: leaves gradually larger upwards, not decurrent, very concave, not margined nor revolute at the entire borders; cells wide, pellucid, upper sub-rhomboidal, lower sub-rectangular; lowest leaves very small, short-elliptic and blunt, loosely disposed, others crowded, median sub-oval and blunt, uppermost ovate-oblong, sub-acute; costa finally red, percurrent in upper leaves, abbreviate in lower: barren. Mac. Cat. 127.—Borders of ditches: Vancouver Island.

314. *Bryum rubicundulum* C. M. & Kindb.—Tufts compact, olive green, very radiculose to innovations: leaves appressed when dry, concave, obtuse, lower oblong, upper ovate, forming small buds, chlorophyllose, not yellow or hyaline, rarely reddish; borders revolute, narrow-marginate nearly all around; upper cells wide; costa red, sub-percurrent: perichætal leaves narrow, sub-acute, narrowly areolate: capsule inclined, obovate-oblong, short-necked, constricted below the mouth when dry, purplish brown; peristome pale yellow; inner membrane low, cilia short, nodulose or appendiculate: dioicous. Mac. Cat. 129.—Summit of Hermit Mountain, B. C.

315. Bryum Vancouverense Kindb.—Differs from *B. caespiticium* in leaves longer acuminate, at base red, reflexed only below middle; cells small, upper very narrow; costa very thick, excurrent in most leaves; capsule not pendent, not constricted below the mouth; lid deplanate; spores larger; pedicel very long; stem indistinct with gemmiform innovations. Mac. Cat. 129.—On wet slopes: Mt. Finlayson, Vancouver Island.

316. Bryum synoico-caespiticium C. M. & Kindb.—Tufts dense, about 1 cm. high, radiculose only at base, green above, brown decolorate below; leaves erect, not decurrent, sub-ovate, acuminate, not margined, recurved at borders, slightly denticulate above; costa long excurrent, very rigid, thick, often denticulate, light brown; capsule turgid, oval, very short-necked, not or slightly constricted below mouth, pendent; peristome perfect; lid low, umbonate; spores about .015 mm.; pedicel geniculate below middle, arcuate at apex, thick, red; synoicous. Mac. Cat. 128.—On earth: old channels of the Illecillewaet River, Revelstoke, B. C.

317. Bryum heteroneuron C. M. & Kindb.—Sub-species of *Bryum capillare*: differs in leaves finally purplish-colored, less contorted when dry, distinctly serrate above, finally red-margined; costa dark purplish, either long excurrent with point thick and very rigid or vanishing below apex; dioicous: agrees with *B. Donianum* Grev. in clavate branches and smaller spores, but leaf borders not incrassate, capsule not so large. Mac. Cat. 130.—On earth: Ottawa, Ont.; Hastings, B. C.; on rocks: Vancouver Island.

318. Bryum heteroneuron brevispldatum Kindb.—Leaves short pointed. Mac. Cat. 131.—Vancouver Island.

319. Bryum elegans Nees.—Dioicous: tufts dense and soft, green above, reddish brown within, densely brown radiculose; lower stem and branch leaves obovate, concave, margin plane, border narrow; comal leaves crowded, imbricate, obovate from a narrow reddish base, carinate concave, margin plane, recurved only at base, toothed above; costa quite strong, excurrent as a smooth somewhat recurved awn; cells very lax, thickened, mostly rhombic hexagonal, at base rectangular and generally reddish; seta curved above; capsule horizontal or nodding, symmetric, long oblong to almost cylindric, with a neck equally long, when dry constricted below mouth; lid very convex, scarcely apiculate; annulus broad; teeth of peristome narrowly margined; basal membrane of the inner peristome $\frac{1}{2}$ height of teeth; segments narrow, cleft or only split; cilia delicate, appendiculate. Limpr. Laubm. 2: 388.—Miquelon Island.

320. Bryum Floridanum Ren. & Card.—Closely allied to *B. Donianum* but much more delicate, leaves shorter, border narrow, scarcely incrassate, margins plane. Revue Bryol. 20: 4. 1893.—Florida.

321. Bryum acutifusculum C. Müll.—Dioicous; tufts slender, low, rather dense, tomentose; fertile stem short, simple or with few branches: leaves

aggregated in a small sub-rosulate tuft, erect-imbricate, spreading when moist, small, oblong-acuminate, regularly concave; costa carinate, yellowish, flexuous, excurrent into a short sharp point; margin strongly revolute, bordered, quite entire; cells small, very regularly rhomboidal, empty; capsule on a short (1.2 cm.) slender flexuous fuscous seta, minute, pyriform-oval, narrow, fuscous, leptodermous; neck elongate (comparatively), very slender, arcuate; lid minute, short conic and acutish from a convex base; annulus broad, revoluble; teeth of peristome very narrow, yellowish, median line none, margin scarcely cristate, prolonged into a sub-hyaline filiform point; segments of endostome a little shorter, very slender, little sulcate, neither gaping nor perforate, cilia single, equaling segments, remotely somewhat appendiculate. *Flora* 70: 220. 1887.—On hornblende rocks: Chilcoot, Alaska.

322. *Bryum Sawyeri* Ren. & Card.—Loosely or densely cespitose, green or brownish; stems short, radiculose below, branching by several innovations, more or less copiously provided in upper part with brown thick simple articulate very caducous filaments arising from axils of leaves; leaves regularly distant, or upper tufted, open when moist, erect imbricate, often slightly twisted when dry, rather concave, oblong sub-spatulate, narrowed at base, shortly acuminate, plane on borders, obsolete denticulate in upper part; costa percurrent or shortly excurrent; cells large, hexagonal, thin-walled, marginal cells elongated; seta reddish, capsule brown or rufescent, pendent, finally oblique or subhorizontal, defluent into a long attenuate neck, constricted below mouth after fall of convex apiculate lid; teeth of peristome densely trabeculate; segments split, cilia long, appendiculate; inflorescence seems to be dioicous, male flowers unknown. *Bot. Gaz.* 14: 95. 1889.—On sandy ground at base of trees or on decayed logs: Enterprise and Beauclerc, Florida; Louisiana.

323. *Bryum extenuatum* Ren. & Card.—In loose yellowish tufts: stems depressed, radiculose, branching below perichatium by elongated innovations, erect, slender, flexuous, generally attenuate and flagelliform; stem leaves distant, equal, erect spreading when moist, imbricate when dry, concave, from a long decurrent base ovate-lanceolate, shortly acuminate-cuspidate, entire or sub-sinuate toward point; innovation leaves much smaller and narrower; margins revolute from base to above middle, costa excurrent into a very short point or vanishing just below apex; cells rhomboidal or hexagonal, rectangular at base, longer and narrower on margin but not forming a distinct border; seta flexuous, reddish below, yellowish above; capsule pendulous, narrowly cylindrical, constricted below mouth when dry, tapering to a long attenuate neck; lid convex, acutely apiculate; teeth pale yellow; internal membrane very broad, segments

split, cilia 2-3, long appendiculate; annulus of 2-3 rows: dioicous. Bot. Gaz. 15: 57. 1890.—Wet sunny bluffs: Portland, Oregon.

324. *Bryum erubescens* Kindb.—Stems short, scarcely 1 cm. high, loosely tufted: leaves of innovations green, small; stem leaves few, reddish-brown, not decurrent, ovate-lanceolate, acuminate, red margined, not revolute at entire borders; areolation pellucid, decolorate, upper cells narrow rhomboidal or hexagonal; costa red, more or less excurrent; inner perichætal leaves smaller, immarginate, slightly sinuate above, with a percurrent costa: capsule small, regular, pendent, narrow-oblong, constricted below mouth; cilia appendiculate; lid convex, mamillate; spores extraordinarily large, about 0.03 mm.: dioicous. Mac. Cat. 118.—On roots of trees: Lake Louise, Rocky Mountains.

325. *Bryum obtusifolium* Kindb.—Closely related to *B. Neodamense*: tufts broad and tumid, 4-7 cm. high, dirty green tinged with red: stems soft: leaves lax, long decurrent, oval, obtuse, very concave, not limbate, slightly reflexed to cucullate apex, entire; cells very lax, slightly thickened, faintly pitted, nearly all rhombic-hexagonal: sterile. Limpr. Laubm. 2: 420. 1892.—Godhavn and other localities in Greenland.

326. *Bryum erythrophyllum* Kindb.—Closely allied to the arctic *B. obtusifolium*: barren stems laxly cohering, blood red, about 1 cm. high: leaves small, reddish, loosely disposed, flaccid, not decurrent, lower leaves ovate, sub-obtuse, upper leaves sometimes more attenuate, furnished with a rigid point to the excurrent costa, entire, sub-immarginate, not reflexed at margins; cells lax, nearly uniform; costa red, percurrent or slightly excurrent, in a short recurved point. Mac. Cat. 131.—In boggy places: Truro, N. S.; Revelstoke, B. C.

327. *Bryum erythrophyllodes* Kindb.—Nearly allied to *B. erythrophyllum*: agreeing in leaves loosely disposed, rose-red, not decurrent: differs in leaves quite immarginate, narrower, ovate-lanceolate, long acuminate or acute; upper cells narrower; costa in upper leaves somewhat long excurrent, rigid: barren. Mac. Cat. 272.—Cypress Hills, Alberta.

328. *Bryum anæctangiaceum* C. M. & Kindb.—Tufts compact, radiclese, small, scarcely 2 cm. high; leaves small, rose colored, loosely disposed, not decurrent, spirally contorted when dry, ovate-acute, pellucid, narrow marginate, entire, faintly revolute, laxly areolate; costa thick, reddish, percurrent or short excurrent: probably monoicous, but male flowers not well developed. Mac. Cat. 130.—On rocks: Hector, Rocky Mountains, B. C.

329. *Bryum crassirameum* Ren. & Card.—In robust wide compact yellowish-green tufts: stems robust, stout, branched: leaves erect-spreading when moist, imbricate when dry, ovate-lanceolate, shortly acuminate, not decurrent, very entire or sub-sinuate at apex; margins revolute from base to near apex: costa percurrent or vanishing just below point; cells rectangular at base, hexagonal above, narrower on margins but not forming a distinct

border: seta reddish, flexuous; capsule pendulous, badious or ferruginous, cylindrical, constricted below mouth when dry, tapering to an attenuate neck; lid conic or sub-convex, apiculate; teeth yellow, densely trabeculate; segments split, cilia 2-3, appendiculate; annulus triple: dioicous.—Bot. Gaz. 15: 57. 1890. Wet running bluffs: Oregon City, Oregon.

330. *Bryum pseudotriquetrum gracilescens* Sch.—Branches elongated, delicate; leaves more distant. Husnot, *Musc. Gall.* 249.—Wisconsin; Wyoming; Washington.

331. *Bryum pseudotriquetrum hyalodontium* C. M. & Kindb.—Differs principally in capsule pale brown and peristome pale, hyaline. *Mac. Cat.* 133.—On wet rocks: Hector, Rocky Mountains, B. C.

332. *Bryum denticulatum* Kindb.—Differs from *B. pseudotriquetrum* in leaves distantly denticulate to middle, lanceolate acuminate, reflexed at base: barren. *Mac. Cat.* 133.—In bogs and springy places: Cypress Hills, Alberta; Selkirk Mountains, B. C.

333. *Bryum hydrophilum* Kindb.—Closely allied to *B. pseudotriquetrum* but leaves narrower, ovate-lanceolate, reflexed at base or not at all: costa yellow; tufts loose, radiculose only below: flowers and capsules not found. *Mac. Cat.* 133.—In springy places: Vancouver Island.

334. *Bryum Duvallii lato-decurrrens* C. M. & Kindb.—Differs in the upper leaves, looser disposed, longer decurrent; uppermost more distinctly apiculate, with a sometimes excurrent costa; longer short and broad: stem more radiculose. *Mac. Cat.* 132.—Wet earth: Selkirk Mts. and Revelstoke, B. C.

335. *Bryum Duvallii Gaspeanum* Kindb.—Differs in leaves narrow lanceolate, short decurrent, greater part margined, sometimes faintly denticulate above; costa percurrent or short excurrent. *Mac. Cat.* 272.—On wet rocks: Gaspé Co., Que.

336. *Bryum meesioides* Kindb.—Dioicous: densely cespitose: leaves yellowish, brown-margined, not decurrent, ovate-oblong or short lanceolate, short acuminate, faintly denticulate at apex, slightly reflexed on borders at base; cells rhomboidal oblong; costa red, scarcely excurrent, abbreviated in lower leaves: capsule narrowly pyriform, defluent to a longer collum, immature, arcuate, pale, wide-mouthed, constricted below mouth; lid convex, pale yellow, mamillate; teeth pale yellow, connivent; annulus double; cilia perfect, long and appendiculate; basilar membrane very low or indistinct. *Mac. Cat.* 133.—Wet rocks: Gaspé Coast, Que.; Banf, Rocky Mountains; Selkirk Mountains, B. C.; Vancouver Island.

337. *Bryum Ontariense* Kindb.—Intermediate between *B. roseum* and *B. Beyrichii*: comal leaves very numerous, lingulate, abruptly and shortly acuminate, revolute to $\frac{3}{4}$ or $\frac{3}{4}$, yellow-margined above with great confluent teeth; costa stout, excurrent: capsule pale, with a distinct curved collum half as long, teeth papillose and hyaline above; lid convex,

short apiculate, not oblique. Mac. Cat. 135. *B. roseum* L. & J. Man. 240, not Schreb. In old logs and sometimes on limestone rocks in maple woods: Ontario; Quebec; New Brunswick; Ohio.

338. *Bryum lucidum* Britton.—Plants slender, scattered, not gregarious, light glossy green: stems from radiculose stolons, simple and naked below: leaves rosulate, not twisted when dry, broadly elliptical above, with parallel margins at base, blunt, with costa vanishing below apex, or with a serrate cuspidate point; vein heavy, frequently red at base; margins of long proserchymatous cells forming small appressed teeth, entire below; cells of the lamina parenchymatous, elongated-hexagonal: dioicous: setae single, stramineous, lustrous and sulcate; capsule reflexed, horizontal or erect, constricted below mouth, neck short: teeth with a faint median line, segments of endostome split, cilia 3-4, not appendiculate, faintly papillose, very irregular, variously divided and elongated; lid apiculate. Bull. Torr. Bot. Club 18: 53. *Bryum simplex* Kindb., Mac. Cat. 135. *Mntum Roellii* Broth., Bot. Centralbl. 44: 420. 1890. Idaho; Montana; Washington; Gold Range, B. C.

339. *Bryum bullatum* C. Müll.—Dioicous: very small, slender: perichætium with few very short very slender julaceous branches: stem leaves minute, very densely imbricate, oval, very shortly acuminate, carinate-concave; margin plane, quite entire; costa yellow, slender, vanishing below apex; cells small; pellucid, elongate, narrowish: perichætial leaves larger, longer, from a lanceolate narrow base ligulate attenuate, obtusish; costa much longer, flexuous, carinate, reddish; cells much longer, looser, yellowish: capsule nutant on a slender reddish seta scarcely 1 in. long, very small, oval barrel-shaped, when young bullate-tuberculose; collum none; annulus broad; lid shortly convex-conic, ochraceous; peristome teeth small, slender; margin scarcely cristate; segments very narrow, cilia nodose (?). Flora 70: 221. 1837.—Takhin Valley, Alaska.

340. *Bryum hematophyllum* Kindb.—Stems short, loosely tufted or solitary: leaves small, red, densely crowded, not decurrent, ovate or ovate-oblong, acute, indistinctly margined, reflexed at entire borders; cells red, nearly uniform and loose; costa very broad, not (or very shortly) excurrent: perichætial leaves oblong lanceolate: capsule (not ripe) regular, pendent, obovate, lid convex, short apiculate, pedicel thir, spores small: dioicous. Mac. Cat. 118.—On wet rocks: Rocky Mountains.

341. *Bryum oligochloron* C. M. & Kindb.—This species is still doubtful, being found only in a barren state.—It is peculiar in upper leaf-cells very narrow; leaves ovate-lanceolate, long acuminate, nearly hyaline and costa long excurrent. Mac. Cat. 129.—On rocks: British Columbia.

342. *Bryum microcephalum* C. M. & Kindb.—This is a doubtful species, not having been found with capsules. It somewhat resembles *B. caespiticium*; stem proliferous with small male buds, leaves contorted,

costa percurrent or short excurrent, inflorescence dioicous. Mac. Cat. 134.—
On rocks: British Columbia.

343. *Bryum pygmaeo-alpinum* C. M. & Kindb.—Tufts small, dense: stem sparingly radiculose to the buds: leaves small, olive-green, finally reddish, flaccid, not glossy, laxly areolate, hyaline; uppermost close, others loosely disposed, not decurrent, very much smaller; all subovate or oval obtuse, uppermost sometimes subacute; borders not margined nor revolute; upper cells short rhomboidal; costa broad, purplish, percurrent or in uppermost leaves slightly excurrent: barren. Mac. Cat. 126.—On wet rocks: Vancouver Island.

344. *Bryum sanguilentum* Ren. & Card.—Quite similar to *B. torquescens*, but more delicate: leaves when dry not twisted, scarcely bordered: capsule narrower: flowers dioicous. Differs at first sight from small forms of *B. capillare* in the narrower blood-red capsule, and in the leaves scarcely or indistinctly bordered. Musc. Am. Sept. 20: 31. 1893. *B. capillare* var., Lesq. & James Man. 236.—California.

345. *Mnium cuspidatum tenellum* Kindb.—Differs in all parts being smaller. Mac. Cat. 136.—On earth in woods: Prince Edward Island.

346. *Mnium macroclliare* C. M. & Kindb.—Bisexual: loosely tufted: stem densely foliate, sometimes stoloniferous: leaves green or bright green, sub-distichous, not decurrent, pale-margined, simply or doubly dentate-ciliate, sub-ovate; perichætal smaller and narrower; costa always excurrent: capsules single or two, oblong, inclined or pendent; lid pale, conic, red-margined, short-rostrate; teeth yellow; pedicel purplish below, yellowish above. Mac. Cat. 137.—On rocks: Revelstoke, B. C.; on earth in damp woods: Ontario.

347. *Mnium insigne intermedium* Kindb.—Agrees with *M. insigne* in leaves long decurrent, long dentate, cells hexagonal-rotundate: differs in syncious inflorescence. Mac. Cat. 139.—On earth: Prince Edward Island; wet rocks: British Columbia; in damp woods: Vancouver Island.

348. *Mnium serratum Macounii* Kindb.—Inflorescence paroicous: leaves red-margined; costa excurrent: lid obliquely rostrate. Mac. Cat. 139.—In dry woods: Prince Edward Island.

349. *Mnium decurrens* C. M. & Kindb.—Dioicous: loosely tufted: stem very loosely foliate, naked below, elongate: leaves green or finally reddish, sub-distichous, long decurrent, often red at margins and wings, doubly dentate, lanceolate: inner perichætal ones sublinear, red cuspidate; cells smaller than in *M. umbratile*; costa sub-percurrent, often red, in inner perichætal leaves excurrent: capsule single, oval, inclined; lid pale, conic, not margined, short apiculate, not rostrate, teeth dark yellow; pedicel reddish. Resembles *M. orthorrhynchum* in size of capsule; leaf cells not much larger: differs from *M. umbratile* in narrow leaves, less twisted when dry. Mac. Cat. 140.—On stones: British Columbia.

350. *Mnium inclinatum* Lindb.—Dioicous: close to *M. orthorrhynchum*: plants 2-3 cm. high, slender: leaves larger above, uppermost oblong, obtuse, narrowly thick margined, double-toothed; costa toothed on back, vanishing in point, cells small, not pitted: seta 1.5 cm. long, curved above; capsule hypnoid, erect or inclined, ovate; neck long, gradually narrowed into seta; peristome yellow; membrane of endostome $\frac{1}{2}$ length of teeth, reddish-brown, segments split; cilia delicate, scarcely nodulose; lid either straight or obliquely beaked. *Limpr. Laubm.* 2: 456.—On rocks and trunks of trees (Drummond); Ottawa, Ontario; British Columbia.

351. *Mnium pseudo-lycopodioides* C. M. & Kindb.—*M. lycopodioides* L. & J. *Manual* 246; not of Schwägrichen. From the closely allied *M. orthorrhynchum* it is separated by excurrent costa of upper leaves, and narrower capsules. *Mac. Cat.* 140.—Newfoundland; New Brunswick; Quebec; Adirondack and White Mts.; Rocky Mountains; British Columbia.

352. *Mnium Niagaræ* Kindb.—Intermediate between *M. spinulosum* and *M. serratum*, but readily distinguished by wide mouthed capsule: loosely tufted without creeping stolons, fertile stem simple: leaves dark green, crisped when dry, indistinctly decurrent, doubly dentate at red-margined borders, lower more loosely disposed, oblong, upper lanceolate; cells rotundate-hexagonal, often pellucid; costa red brown, excurrent at least in upper leaves; perichætal leaves narrow, sublinear, dentate, innermost very short: capsule solitary, pendent, narrow, often appressed to pedicel, sub-clavate-cylindric, constricted below wide mouth, teeth yellow; lid obliquely rostrate; pedicel pale red: inflorescence synoicous. *Mac. Cat.* 141.—On earth in woods: Ontario.

353. *Mnium Blyttii* B. & S.—Dioicous: tufts dense, 6-8 cm. high, bright green with red, densely red radiculose: leaves more crowded above, lower oval, acute, entire, upper oval oblong, acute, uppermost spatulate, acute, all narrowed at base, decurrent, red marginate, toothed to middle, with a double or single row of short, blunt teeth; cells not in divergent rows, irregularly roundish-angulate, faintly thickened, not pitted: capsule horizontal or nodding, long oblong, slightly curved; lid strongly convex, blunt; annulus double or triple; teeth of peristome yellowish-green, blunt, papillose, bordered; endostome nearly orange, basal membrane $\frac{1}{2}$ length of teeth, segments broad, split, suddenly subulate. *Limpr. Laubm.* 2: 460.—Bases of stumps and on earth: Rocky Mountains and British Columbia.

354. *Mnium hymenophyllum* B. & S.—Densely cespitose, tufts from glaucescent to bright green, reddish below, when old blackish brown, interwoven with red rhizoids: plants 2-6 in. high (occasionally higher), erect, branched: leaves rather crowded, erect-spreading, long; decurrent, broadly ovate-acuminate, rather obtuse border narrow, entire or with prominent

cells here and there; costa vanishing below apex; cells roundish-hexagonal: dioicous; male flowers discoid, antheridia few, paraphyses strongly thickened at apex; female flowers and fruit unknown. Bry. Eu. Mnium Suppl. 5. pl. 400. Hartm. Skand. Flora 2: 51. [Ed. 10.].—Greenland.

355. *Aulacomnium palustre congestum* Boul.—Tufts thick, radiculose: branches short: upper leaves slightly secund, sharp pointed. Limpr. Laubm. 2: 530.—Isl. Miquelon.

356. *Aulacomnium palustre laxifolium* Kindb.—Leaves laxly disposed. Mac. Cat. 145.—British Columbia.

357. *Timmia Megapolitana Bavarica* Brid.—Leaves uniform in size; leaf sheath not papillose on back.—*F. Bavarica* Hessel. Limpr. Laubm. 2: 581.

358. *Timmia Norvegica* Zett.—Dioicous: tufts lax, yellowish-green, below brown and laxly radiculose: leaves when old easily breaking off at insertion, above longer and narrower, from a short faintly sheathing base gradually linear-lanceolate, short pointed, erect-spreading, when dry falcate or spirally incurved twisted and almost crispate, from apex to middle short toothed, faintly toothed for remainder; costa strong, red, percurrent or vanishing; cells coarsely papillose in lower dorsal half, upper cells roundish quadrate or hexagonal, below gradually rectangular, in sheathing part elongated rectangular, distinctly pitted: capsule horizontal, oblong oval, short-necked; cilia of endostome delicate, not appendiculate.—Limpr. Laubm. 2: 580. On muddy banks: Hector and discharge of Devils Lake, Rocky Mountains; Greenland.

359. *Timmia Austriaca brevifolia* Ren. & Card.—Stems shorter; leaves more crowded, shorter and erect-imbricate, scarcely flexuous when dry. Bot. Gaz. 19: 238.—Springdale, Colorado.

360. *Atrichum undulatum altecristatum* Ren. & Card.—Lamellae of the leaves much higher than in typical form; capsule narrower, erect. Bot. Gaz. 15: 58.—Kansas; Pennsylvania.

361. *Atrichum lelophyllum* Kindb.—Plants erect, simple, naked below, densely foliate above: leaves smooth, lower very small, upper lanceolate, acute-cuspidate, not margined, base scarcely broader; borders sharply serrate with large teeth, basal only entire; basilar cells narrow, others round-hexagonal, all pellucid; costa percurrent; lamellae 6, disappearing below apex, entire or slightly crenulate; perichaetial leaves long cuspidate, costa excurrent: capsule single, sub-cylindrical, broad, slightly curved, constricted at mouth; lid obliquely rostrate from conic base. Mac. Cat. 148.—In damp woods: Vancouver Island; Selkirk Mountains, B. C.

362. *Atrichum rosulatum* C. M. & Kindb.—Dioicous: stem indistinct or very short: leaves rosulate, undulate, not margined, dentate at middle, sometimes beset with transverse rows of scales, lower sub-oblong, upper

short-lanceolate, sub-obtuse; costa scarcely percurrent, nearly smooth; barren. Mac. Cat. 148.—On bank of Columbia River, Revelstoke, B. C.

363. *Oligotrichum hercynicum* (Ehrh.) DC.—Stems 1-3 cm. high, erect, rigid; tufts large, lax, glaucous green or brownish; leaves erect-spreading when moist, imbricate and incurved when dry, oval and half sheathing at base, then lanceolate-linear, acute, concave, incurved on border and dentate above middle; costa with 10-15 lamellae on upper surface, contorted-undulate, and with 2-4 short dentate lamellae in the upper part on dorsal side; capsule ordinarily oblique, sometimes erect, oblong-cylindric, slightly contracted below mouth, at first yellowish, becoming brown, smooth, irregularly plicate when empty; lid convex-conic, beak short; teeth of peristome unequal, variable in number. Husnot, Musc. Gall. 277.—Greenland; St. Lawrence Island; Behring Sea.

364. *Oligotrichum hercynicum latifolium* C. M. & Kindb.—Leaves shorter, entire; capsule distinctly plicate. Mac. Cat. 149.—Selkirk Mts.

365. *Pogonatum erythrodontium* Kindb.—Stem erect, simple, or furnished with a short innovation above, naked below; leaves green, spreading, flexuous and not incurved when dry, plane and patent when moist, linear, lanceolate, acuminate, red-denticulate all around, not sheathing, undulate, hispid or margined; cells round-hexagonal, chlorophyllose (also the marginal); lamellae distinct, not confluent, numerous, 20-30; costa narrow, percurrent; capsule single, short, ovate cylindrical, erect, papillose, wide mouthed; teeth red brown, narrowly hyaline margined all around, narrowly lingulate; dioicous. Mac. Cat. 150.—Vancouver Island; British Columbia; Alaska.

366. *Pogonatum alpinum microdontium* Kindb.—Leaves nearly entire, or indistinctly denticulate.—Mac. Cat. 152.—St. Paul Is., Behring Sea.

367. *Pogonatum Macounii* Kindb.—Dioicous; laxly caespitose, green, brownish when old; stem rooting only at base, dichotomously branched; leaves very long; when dry spreading or patulose, flexuose and convolute; when humid sub-plane, linear-lanceolate; cuspidate from short sheathing dirty white base, densely and sharply serrate, spinulose and reddish at back towards the apex; lamellae numerous, about 60, entire, each of round hexangular cells; costa long excurrent; lower cells or leaf and base elongate and narrow, upper oblong, cell walls often oblique and irregular; perichaetial leaves shorter and more acute; seta robust, pale, finally orange; capsule obliquely inclined, cylindrical-oblong, without neck, narrowed below mouth, not angulose; lid large, conic-subulate, orange at base; teeth of peristome 32, short. Mac. Cat. 153.—On earth and rocks: British Columbia; Vancouver Island; Alaska.

368. *Polytrichum conorhynchum* Kindb.—Differs from *P. formosum* in leaves canaliculate, capsule shorter and broader, not attenuate at base:

nearly smooth:
 elstoke, B. C.
 1-3 cm. high,
 leaves erect,
 oval and half
 e, incurved on
 on upper sur-
 lae in the upper
 s erect, oblong-
 wish, becoming
 vex-conic, beak
 Husnot, Musc.

Kindb.—Leaves
 Selkirk Mts.
 , simple, or fur-
 green, spread-
 nt when moist,
 not sheathing,
 trophyllous (also
 s, 20-30; costa
 erect, papillose,
 ned all around,
 Island; British

aves nearly en-
 s., Behring Sea.
 espitose, green,
 ously branched:
 and convolute;
 short sheathing
 and reddish at
 e, each of round
 and base elon-
 l irregular; per-
 finally orange;
 , narrowed be-
 t base; teeth of
 British Colum-

P. formosum
 nuate at base:

differs from *P. gracile* also in capsule acute angulate, lid conic, rostrate, leaves long sheathing. Mac. Cat. 154.—In boggy ground; Selkirk Mts.

369. *Polytrichum sexangulare* Florke.—Stem 2-6 cm. long, generally simple, erect, or decumbent, naked at base; tufts large, brownish green, not tomentose: leaves rigid, erect incurved, sometimes incurved, large oval and sheathing at base: then abruptly contracted into a lanceolate-canaliculate point, sub-obtuse, very entire; lamellæ about 30; perichaetial leaves longer sheathing, mucicous: capsule erect or inclined, angular, six-sided; lid long beaked; teeth unequal; calyptra not reaching base of capsule. Husnot, Musc. Gall. 279.—Mt. Hood, Oregon.

370. *Polytrichum Ohioense* Ren. & Card.—Stem erect, simple or bipartite, a little tomentose below: leaves spreading when moist, erect flexuous when dry, from a sheathing base linear-acuminate, cuspidate, serrate; lamellæ about 50; perichaetial leaves longer with a hyaline base; pedicel 4-8 cm. long, reddish below, pale above; capsule erect, finally horizontal, tetragonal or pentagonal, rarely hexagonal, acute angled, rather narrowed towards the base, with a very small or indistinct hypophysis. Bot. Gaz. 13: 199. 1898.—Northern, eastern and southern United States; British Columbia; Canada.

371. *Polytrichum Canadense* Kindb.—Stem low, about 6-8 cm. high: pedicel not much longer; capsule blackish, much shorter than in the type.—Prince Edward Island; Quebec, Ontario; Lake Nepigon; Selkirk Mountains, B. C.

372. *Polytrichum commune* minus Weiss.—Plants shorter and more slender: leaves dense, shorter and more erect; perichaetial bracts less distinct. Capsule much smaller and shorter, less acutely quadrangular, lid with short straight beak; calyptra pale golden brown. Braithw. Brit. Moss Flora 1: 59.—New Jersey.

373. *Polytrichum polare* C. Müll.—Dioicous: low, dendroid, everywhere dark green, below with closely appressed small scales, above fastigiate branched; branches long for height of plants, rather slender, again divided, flexuous, equal: stem leaves closely appressed, when moist erect spreading, rather strict, from a large broadly-sheathing base obovate; shortly lanceolate; costa broad, multilamellose, opaque, occupying whole lamina, excurrent into a short blackish point; margin strongly serrate with robust teeth (rarely none) lobed at base and very sharp at apex; cells elongate, very narrow, close, flexuous, yellow, at margin very slender forming a pale border, above sinuate-narrowed; scales of undivided lower stem quite entire or slightly dentate at base, very shortly lanceolate; perichaetial leaves like; capsule on short thick red seta (scarcely 1 in.), inclined, small, quadrangular, bistrucosae at base; teeth very narrow, whitish. Bot. Zeit. 17: 205. 1869.—Greenland.

874. *Polytrichum juniperinum* Waghornel Kindb.—Leaves dentate in upper part, not in lower part only; marginal cells papillose. Mac. Cat. 155.—Labrador.

875. *Buxbaumia indusiata* Brid.—Dioicous: very similar to *B. aphylla*: seta 5-10 mm. high, thicker, yellowish red with smaller warts; capsule erect or sub-erect, when old less inclined, ovate-oblong, both surfaces same color lightly shining, pale olive-green, when empty yellowish brown; dorsal surface faintly swollen, the external cuticle at length ruptured along the dorsal side and revolute; neck shorter and thicker; lid shorter, convex-conic; annulus narrower: separating as single cells; outer peristome free, formed of three or four rows of teeth; teeth numerous, irregular, small on outside; larger within, linear, triangular in cross section, perforate here and there in middle line, pale red, papillose, when moist erect; inner peristome as in *B. aphylla*. Lampr. Laubm. 2: 640.—On decaying coniferous logs: Washington; Idaho; Catskill Mts., N. Y.

876. *Buxbaumia Piperi* Best.—Dioicous: stemless; leaves reduced to yellowish ovate-oblong or palmate crenate-laciniate bracts; cells oblong hexagonal: bracts of fertile plants producing long rhizoids enveloping the fleshy vaginule in a felt like mass: seta about 1 cm., arcuate or flexuose, warty, obliquely inserted; capsule inclined, ovate oblong, unsymmetric, greenish, becoming pale yellow, section broadly elliptical, upper surface not deeply impressed nor strongly margined; neck distinct; cuticle thin, not glossy or but slightly so, rolling back in segments after loosening of broadly conical obtuse operculum; endostome of linear papillose hyaline fluted segments, lightly cohering by their thickened margins forming a truncated cone; peristome of a single layer, with rudiments of a second deeply inserted, teeth linear, reddish or dirty white, papillose, articulated, revolute, lightly connate; pseud-annulus usually of 3 layers. Bull. Torr. Bot. Club 20: 116. 1893.—On rotten wood or on ground covered with woody debris: Washington; Idaho.

877. *Fontinalis antipyretica Oregonensis* Ren. & Card.—Stems very slender, soft, pinnate, yellowish and shining above, not naked below; stem leaves distant, open, lanceolate; branches cuspidate; branch leaves erect, imbricate at the top of branches; cells very long and narrow. Bot. Gaz. 14: 96. 1889.—In swamps, on roots of trees: Oregon.

878. *Fontinalis antipyretica rigens* Ren. & Card.—Plants rigid, shining, of a yellowish or copper green recalling the aspect of *Hypnum cordifolium*: stems rigid, naked at base, branches spreading divaricate: leaves distant, spreading or erect-spreading, imbricate only at ends of branches, lanceolate, obtuse or sub-obtuse; cells narrow, elongated; frutification unknown. Monog. des Fontin. in Mem. Soc. Sci. Nat. Cherbourg 23: 55. 1892.—Washington; Vancouver Island.

379. *Fontinalis Neo-Mexicana* Columbia Card.—Plants quite soft, lax, generally shining, bright or yellowish green: stems often red at base: leaves distant, spreading, imbricate at end of stem and branches, sometimes slightly flexuose when dry. Monog. des Fontin., l. c., 61.—British Columbia; Washington; Idaho; California.



380. *Fontinalis maritima* C. Müll.—Plants slightly rigid, yellowish green: stems elongated, naked at base; branches numerous, sometimes fasciculate, rigid, delicate, elongated, flexuose, erect, triangular cuspidate: leaves small, crowded, imbricate for the whole length of stem and branches, oval-lanceolate, acuminate, acute, entire, carinate-conduplicate, nearly straight; cells as in *F. Neo-Mexicana*, very narrow and elongated, walls rigid and slightly thickened; cells of the angles numerous, sub-hexagonal, yellowish or sub-hyaline: frutification unknown. Monog. des Fontin., l. c., 61.—Growing with *Polysiphonia* and other marine algæ, Neah Bay, Washington.

381. *Fontinalis Kindbergii* Ren. & Card.—Plants robust, soft, brownish, or sometimes yellowish or red, copper colored and shining above: stem generally more than a foot long, flexuose, naked and black at base, divided into many more or less pinnate branches; branches spreading or erect, a little flexuose, plumose, sometimes cuspidate: leaves dimorphous; stem leaves erect-spreading or loosely imbricate, more or less carinate or only plicate, broad oval-lanceolate, long narrow acuminate, cuspidate, entire or sub-denticulate at summit, concave, upper large, lower much smaller and shorter acuminate; branch leaves tristichous, divaricate, narrowly lanceolate, long acuminate, concave, inflexed on margins, canaliculate above, rounded or sub-carinate at back; cells long linear, at angles enlarged, sub-square, yellowish or ferruginous; perichæstrial bracts convolute, suborbicular, entire at truncate-rounded apex: capsule immersed, oblong, lid conic; teeth narrowly linear, slightly papillose, often connected in pairs at apex, with 25-35 lamellæ, not perforated at dorsal line; lattice cone of endostome perfect, papillose, transverse bars appendiculate: dioicous. Bot. Gaz. 15: 58. 1890.—In streams, lakes, and ponds: Vancouver Island; Oregon; Washington; Idaho.




382. *Fontinalis Kindbergii* Howellii Ren. & Card.—Stems rigid, subdendroid: regular pinnate, spreading, often recurved and plumose, leaves more rigid, dimorphism much more pronounced. *F. Howellii*, Bot. Gaz. 13: 200. 1888.—Oregon and Washington.

383. *Fontinalis chrysophylla* Card.—Plants rigid, shining, golden yellow above, brown at base: stems delicate, naked below, irregularly pinnate; branches distant, spreading, plumose, sub-attenuate: leaves scattered, rigid, spreading or erect spreading, slightly dimorphous; stem leaves narrow lanceolate, concave, carinate-conduplicate, narrow acuminate, obtuse or sub-obtuse, entire or slightly denticulate at summit; branch leaves and





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

1.8
2.0
2.2
2.5
2.8
3.2
3.6
4.0

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

1.0
1.1
1.2
1.4
1.6
1.8
2.0

© 1981

those of upper part of innovations narrower, long and narrowly acuminate, almost subulate; acute, not carinate, canaliculate, sometimes subtubulose above, entire or sub-denticulate at apex; cells compact, narrow, very long, attenuated, walls rigid, thickened; cells of angles dilated, large, oblong sub-hexagonal or sub-rectangular, brown or yellowish, forming distinct auricles; perichætal leaves broadly oval, obtuse, finally lacerate at summit: capsule immersed, oblong sub-cylindric; lid and peristome unknown. Monog. des Fontin., l. c., 67.—Olympic Mountains, Washington.

384. Fontinalis Delmari Ren & Card.—Dull green, naked below, much divided; branches irregularly pinnate, branchlets erect-spreading, attenuate; leaves crowded, erect, loosely imbricate, oblong lanceolate, obtusely acuminate, concave, not keeled, slightly incurved on the borders; cells linear-elongated, sometimes rather flexuous; alar cells few, small sub-hexagonal; perichætal leaves often lacerate at rounded apex: capsule immersed, oblong sub-cylindric; lid conic acuminate; teeth of peristome narrowly linear acuminate, lamellæ 14-20, divisural line distinct at base only and not perforated; cilia united at apex only, lower transverse bars imperfect, papillose, not appendiculate. Bot. Gaz. 14: 96. 1889.—Growing on stones in streams: Island of Miquelon.

385. Fontinalis mollis C. Müll.—Plants soft, floating, of a beautiful green: stems long, naked at base, much divided, irregularly pinnate, branches robust, unequal, short-cuspidate or sub-obtuse, erect spreading: leaves loosely imbricate, or erect spreading, soft, plicate when dry, very concave, inflexed on borders, broad oval, obtuse, entire; cells at angles sub-hexagonal, yellowish, greenish or concolorous and scarcely distinct; others linear rhombic, sub-flexuose, very chlorophyllose, walls delicate; upper perichætal leaves oval sub-orbicular, truncate and finally lacerate at summit: capsule immersed, oval; lid conic, obtuse; peristome purple, teeth linear-acuminate, faintly papillose, lamellæ 23-32, entire or slightly perforated near base in middle line, generally cohering in pairs at summit; transverse bars of endostome imperfect, strongly muricate. Monog. des Fontin., l. c., 90.—Washington.

386. Fontinalis Novæ-Angliæ Howel Card.—Leaves more acuminate, sometimes sub-acute; capsule half immersed. Monog. des Fontin., l. c., 93.—On rocks in streams: Ft. Edwards, N. Y.

387. Fontinalis Novæ-Angliæ Eatonl Card.—More shining, copper color: stems softer, less regularly pinnate, branches erect: leaves crowded, more erect, a little firmer and more elongated. Monog. des Fontin., l. c., 94.—On rocks in streams: New Hampshire.

388. Fontinalis Cardoti Ren.—Plants soft, dirty green or yellowish: stems flexuous, naked at base, much divided, pinnate branches often fasciculate, erect-spreading, robust, obtuse or short attenuate: leaves quite crowded, erect-imbricate, very concave, inflexed at margin, broadly oval-

lanceolate, obtuse, denticulate at summit, rarely almost entire; alar cells dilated, oblong, yellowish or sub-hyaline, forming distinct auricles; others narrow, linear, sub-flexuose, attenuate, walls rigid and thickened; upper perichaetial leaves sub-orbicular, broadly rounded or sub-apiculate at summit, not lacerate; capsule immersed, narrow, cylindrical; lid conic, elevated, acuminate; peristome purple, teeth narrowly linear acuminate, strongly papillose, lamellæ 13-16, middle line scarcely apparent; cross bars of endostome imperfect, cilia very muricate, united only at summit. Monog. des Fontin., l. c., 95.—Virginia.

389. *Fontinalis involuta* Ren. & Card.—Plants slightly rigid or soft, dirty green or yellowish; stems finally naked at base, much divided, pinnate; branches spreading or erect-spreading, distichous, obtuse or cuspidate, generally compressed; leaves quite crowded, rigid, erect-spreading, oval oblong, oblong-lanceolate or sub-linear, very concave, strongly inflexed at border, canaliculate, generally cucullate at apex, terminated by a broad short obtuse or sub-acute slightly denticulate acumen; cells of angles dilated, oblong or sub-hexagonal, brown or yellowish, forming distinct auricles; others linear, narrow, walls delicate or slightly thickened; fructification unknown. Monog. des Fontin., l. c., 96.—Louisiana; Florida; New Jersey.

390. *Fontinalis nitida* Kindb. & Arn.—Plants soft, lax, delicate, generally somewhat shining above, yellowish green, dark below, becoming black when old; stems delicate, flexuose, more or less naked below, branching; branches long, erect-spreading, more or less cuspidate; leaves scattered, erect-spreading, plane or slightly concave, oval lanceolate or oblong lanceolate, gradually narrowed, acuminate, acute or sub-obtuse, entire or sinuate at the apex; cells of angles dilated, oblong, sub-rectangular or sub-hexagonal, yellowish, forming small auricles; the remainder, elongated linear-rhombic with delicate walls; upper perichaetial leaves oval sub-orbicular, rounded or sub-apiculate, finally a little lacerate at summit; capsule immersed, small, oval, rounded at base; lid short conic; peristome reddish orange, teeth narrowly linear, often cohering in pairs, papillose, generally not perforate along middle line; cross bars of endostome perfect, strongly muricate, almost spinulose. Monog. des Fontin., l. c., 103.—On rocks and trunks of trees at the edge of water: British Columbia.

391. *Fontinalis tenella* Card.—Plants soft, very delicate, pale yellowish green; stems delicate, more or less naked at base, divided in elongated branches, sub-simple; leaves erect-spreading, soft, plane, narrowly lanceolate, very long and gradually acuminate, acute, entire or sinuate at summit; cells of angles dilated, oblong or sub-rectangular, yellowish or greenish, forming small auricles; others narrow, linear, walls delicate; upper perichaetial leaves broadly oval or sub-orbicular, rounded, generally small apiculate and finally lacerate; capsule immersed, small, oval or oblong,

rounded at base; lid short conic; peristome orange red, teeth narrowly linear, often cohering in pairs, strongly papillose, lamellæ 15-20, entire along middle line; cross bars imperfect, strongly muricate. Monog. des Fontin., l. c., 105.—On submerged trunks of trees and grauitic rocks: Lake Pend d' Oreille, Idaho.

392. *Fontinalis Durlæi* Sch.—Plants soft, lax, pale olive or yellowish green, often blackish at base: stems delicate, more or less flexuose, entire, foliate or naked below; branches distant, but quite numerous, unequal, short or elongated, spreading or erect, attenuate, tapering or cuspidate: leaves more or less scattered, spreading or erect-spreading, imbricate at extremity of branches, soft or slightly rigid, slightly concave, often almost plane, sometimes a little plicate longitudinally; stem leaves broad, oval-lanceolate, or oval-oblong; branch leaves oblong lanceolate, acute, sub-acute or sub-obtuse, slightly denticulate at the summit; cells of angles short, sub-hexagonal or sub-rotundate, brown or yellow; others elongated, narrow, attenuate, linear-rhombic, walls delicate and soft, sinuous, upper cells much shorter; upper perichaetial leaves oval sub-orbicular, rounded at apex, short and obtusely apiculate, finally lacerate: capsule immersed, oval or short oblong, rounded at base, not contracted below mouth when dry; lid conic: peristome beautiful purple or orange-red, teeth linear, sometimes sinuous on margin, papillose, frequently cohering in pairs, entire or perforate along middle line; cross bars of endostome perfect. Monog. des Fontin., l. c., 111.—California.

393. *Fontinalis flaccida* Ren. & Card.—Plants very soft, yellowish: stems slender, naked below; branches subpinnately ramulose; branchlets spreading, slender, distant: leaves soft, distant, open, but convolute-imbricate at top of branches, elongated, narrowly lanceolate, plane or nearly so, obtuse or truncate, slightly denticulate at apex; cells thin-walled, median very long, upper much shorter; those of angles large, lax, sub-rectangular, or sub-hexagonal, hyaline or brownish, forming very distinct auricles; perichaetial leaves broadly oval, truncate and finally lacerate at summit: capsule immersed, sub-cylindric, rounded at base, not contracted below mouth when dry; lid conic, elevated; peristome purple, teeth often cohering in pairs, narrowly linear-acuminate, weakly papillose, entire along middle line; cross bars of endostome imperfect, muricate. Bot. Gaz. 13: 201. 1888.—Growing in stagnant or running water: Louisiana; Alabama; White Mountains.

394. *Fontinalis microdonta* Ren.—Plants very slender and delicate, soft, entirely yellow or dark green at the base: stems delicate, filiform, reddish, naked at base, irregularly pinnate, branches distant, erect or erect-spreading, attenuate or cuspidate: leaves scattered, erect-spreading, soft, narrowly linear lanceolate, very long acuminate, acute, slightly denticulate, sinuate or entire at summit; stem leaves almost plane, branch leaves

sub-canalicate; cells of angles dilated, oblong, yellowish or sub-hyaline; others linear, narrow, attenuate, subflexuous, walls delicate; perichæatial leaves broad oval oblong, slightly narrowed above, entire or finally more or less lacerate, and sometimes plicate at summit: capsule immersed, narrow, cylindrical; lid conic, elevated; peristome purple, teeth narrowly linear acuminate, sometimes cohering in pairs at summit, papillose, entire or perforate along middle line; endostome very rudimentary, with complete transverse bars, cilia mucicate, simply appendiculate. Monog. des Fontin., l. c., 120.—New Jersey.

895. *Fontinalis dichelymoides* Lindb.—Plants somewhat shining, yellowish above, dark brown at base, having the external appearance of *Dichelyma* or certain submerged forms of *Hypnum fluitans* and *H. Kneiffii*: stems slender, flexuous, not or slightly naked at base, more or less regularly pinnate; branches distant, erect, spreading, more or less elongated, sub-attenuate and often slightly curved at summit: leaves scattered, erect spreading or sub-secund, narrowly lanceolate, incurved at border, canalicate, long acuminate, acute, entire; cells of angles dilated, oblong, sub-rectangular or sub-hexagonal, hyaline or yellow, forming quite distinct auricles; others linear, flexuous, attenuate, very narrow: fructification unknown. Monog. des Fontin., l. c., 122.—Growing on submerged rocks. Lake Vermillion, Minn.

896. *Fontinalis filiformis tenuifolia* Card.—Still more delicate than type, plumose, pale green: leaves very distant, more elongated, very soft when moist, rigid when dry: sterile. Monog. des Fontin., l. c., 123.—Louisiana.

897. *Fontinalis Langloisii* Card.—Plants slender, delicate, slightly rigid, dirty green or dark at base, yellowish above: stems delicate, flexuous, naked and black below, pinnate and sub-bipinnate; branches distant, distinct, very delicate, plumose, cuspidate: leaves very distant, sub-rigid when dry, erect-spreading or sub-imbriate, narrowly lanceolate, tubulose or sub-tubulose, generally cucullate, sub-obtuse or obtuse, rarely acute, nearly entire or slightly denticulate at summit; cells at angles a little dilated, oblong, greenish or sub-hyaline; others linear-rhombic, with delicate walls: fructification unknown. Monog. des Fontin., l. c., 126.—Louisiana.

898. *Dichelyma capillaceum elongatum* Kindb.—Stems more elongate, branches more distant, leaf base longer than excurrent part of costa. Mac. Cat. 160.—North of Lake Superior.

899. *Neckera Menziesii amblyclada* Kindb.—Stems densely pinnate; branches obtuse, rarely attenuate and flagelliferous: leaves shorter, sub-oblong. Agrees with the common form in the paraphylliferous stem. Barren. Mac. Cat. 162.—Rocky Mountains, B. C.

400. *Neckera Menziesii limnobioides* Ren. & Card. — Habit of a *Limnobium*, soft and dilated, cespitose, dark-rufescent: leaves smooth, or scarcely undulate, short, entire or obsolete denticulate above, obtuse or apiculate, areolation lax, costate to middle or beyond. — Bot. Centralbl. 44: 422. 1890.

401. *Neckera peterantha* C. M. & Kindb. — Subspecies of *N. oligocarpa*: secondary stems nearly simple, about 1 dm. long, rigid and more robust: leaves larger and more crowded, one branch of the costa prolonged sometimes to middle: antheridia and archegonia very numerous: capsule emergent. Mac. Cat. 162. — On rocks: Rocky Mountains, B. C.

402. *Neckera Douglasii Macounii* Kindb. — Leaves less attenuate: capsule turgid oval, completely exerted on long pedicel. Mac. Cat. 163. — Hanging in long festoons from limbs of trees in shady woods: British Columbia and Vancouver Island.

403. *Homalia Macounii* C. M. & Kindb. — Very nearly allied to *H. trichomanoides*: leaves longer, rather lingulate, lowest basal cells yellowish; perichaetial leaves more suddenly narrowed to very short acumen: segments of peristome cleft between articulations. Mac. Cat. 163. — *H. trichomanoides* and *H. obtusata* Lesq. & James, Manual 285. — Canada; Newfoundland; British Columbia and Vancouver.

404. *Pterigynandrum filiforme heteropterum* Sch. — Plants more robust, darker green; branches shorter and larger: leaves secund, oval, spatulate, rounded and shortly apiculate at summit. Husnot, Musc. Gall. 311. — Washington.

405. *Pterigynandrum papillosulum* C. M. & Kindb. — Differs from *P. filiforme* in branch leaves acuminate and acute, denticulate nearly all round, less papillose; branches blunt and turgid as in the variety *heteropterum* of this species. Mac. Cat. 165. — On rocks: British Columbia.

406. *Antitrichia Californica ambigua* Ren. & Card. — Branches not julaceous: leaves not so closely imbricate, generally subsecund and narrower; cells longer: pedicel often flexuous. Resembles in habit *A. curtipendula*, but differs in cylindrical, narrow capsule, perichaetial leaves longer acuminate, and shorter cells. Bot. Gaz. 15: 59. 1890. — Portland, Oregon.

406a. *Antitrichia tenella* Kindb. — Tufts loose, green: secondary stems irregularly divided; branchlets numerous, the greater number very short: leaves small, sub-patent or loosely appressed when dry, ovate-acute or short-acuminate, dentate at apex, not striate, reflexed to apex: greater number of cells short, inner and middle narrow; costa with 1-2 accessory branches at base; disappearing below apex; perigonal leaves elongate: dioicous: female plants unknown. Mac. Cat. 165. — On rocks: Nanimo River, Vancouver Is.

407. *Thelia compacta* Kindb.—Stems closely creeping; tufts green, very dense and thick; branches erect, terete, obtuse, unilateral: paraphyllia none: leaves cochleariform, rotundate-obtuse, short apiculate, very scabrous at back, with simple, incurved, papilliferous cilia: borders spinulose dentate, or fimbriate ciliate; cilia long, curved up and dentate; costa obsolete or very short; perichæstial leaves oblong, lanceolate, narrowly acuminate, fimbriate, capsule pale brown, ovate-cylindrical; teeth subulate, short and broad, sometimes horizontally divaricate when moist, distantly articulate, dusky, upper article cleft: basilar membrane short, scarcely $\frac{1}{4}$ length of teeth without segments; operculum conic-obtuse. Mac. Cat. 166.—Abundant on stems of young maples: central Ontario; New Brunswick.

408. *Leskea sub-obtusifolia* C. M. & Kindb.—Plants loosely tufted, yellowish green or fuscous: stem sparingly radiculose, irregularly divided, beset with paraphyllia; branches curved at apex: leaves distant, decurrent, entire, distinctly papillose at back, loosely appressed when dry, spreading when moist, margins recurved at base; stem leaves broadly ovate, obtuse or sub-acute; branch leaves oblong, obtuse; cells round, alar quadrate; costa sub-percurrent: perichæstial leaves large, short, ovate-lanceolate, short acuminate, costate: capsule oblong, inclined, sub-curved; lid short, mamillate; monoicous. Mac. Cat. 169.—On trees subject to inundation: British Columbia.

409. *Leskea cyrtophylla* Kindb.—Tufts dense, dark green or brown, not shining: stems irregularly branching, filiform, with few rhizoids: leaves nearly uniform, very small, appressed when dry, open-erect when moist, concave, from a broad rotundate ovate base short acuminate, entire, faintly papillose, borders reflexed below; cells rotundate, sub-hyaline; costa indistinct or none: dioicous: capsules not found. Mac. Cat. 169.—On rocks on islands in Lake Nepigon.

410. *Myriniola* Schimp.—Leaves oval or ovate lanceolate, entire, smooth, shortly costate; cells rhomboidal except those of basal angles which are quadrate: monoicous: annulus none; peristome double, calyptra cucullate, descending to middle of capsule.

411. *Myriniola* (?) Dieckii Ren. & Card.—Cespitose, dirty green: stems depressed, soft, elongated, irregularly branching, branches ascending, curved, julaceous, attenuate: leaves imbricate, concave, ovate lanceolate, sub-acute or obtuse, margin plane throughout or revolute at base, entire or sinuate; costa broad, green, sometimes sub-bifurcate, disappearing far below apex; cells rhomboidal-hexagonal, smooth, wall straight, alar quadrate, numerous, all strongly chlorophyllose; perichæstial leaves acuminate, apex obsoletely denticulate: pedicel red; capsule erect, sub-cylindrical: dioicous. Bot. Centralb. 44: 421. 1890.—Growing on tree trunks: Oregon.

412. *Anomodon attenuatus brevifolius* R. & C.—Leaves shorter, wider above, shorter apiculate, sometimes obtuse, apex entire or denticulate;

cells more distinct; costa less translucent. *Hedwigia* 32: 245. 1893.—Indiana; Illinois; Wisconsin.

413. *Anomodon heteroleus* Kindb.—Plants densely tufted, green, finally fuscous or blackish: stem creeping, subpinnate, much branching and furnished with numerous, small, flagelliform branchlets, densely beset with very small oblong obtuse and nerveless leaves, paraphyllia broad; stem leaves sub-distant, decurrent, appressed when dry, open-erect when moist, from a broadly ovate base, suddenly narrowed to a long subulate or sub-linear acumen, entire, faintly papillose; margins revolute at base; branch leaves more attenuate; cells round oval, marginal basal ones quadrate; costa vanishing below acumen; dioicous: fruit not found. *Mac. Cat.* 172.—On flat limestone rocks and roots of trees: Ontario and Rocky Mountains.

414. *Lescurea* Schimp.—Primary stem obscurely creeping, secondary fertile stems ascending, fasciculately radiculose, perichaetial branch not radican: leaves erect-spreading, costate, smooth, sulcate; cells narrowly oval or oblong rhombic, at angles narrowly quadrate: calyptra long, narrow, sub-tubulose, early deciduous; annulus narrow; teeth confluent to a basilar membrane, firm, narrowly lanceolate, vermicular verrucose, orange; segments from a narrow membrane, equaling or shorter than teeth, irregularly appendiculate, whitish yellow; spores minute.—*Synop. Musc. Europ.* 620. 1876.

415. *Lescurea imperfecta* C. M. & Kindb.—Tufts loose, green, not shining: stem pinnate, radican; paraphyllia few: stem-leaves smooth, decurrent, often bistriate, from a short ovate base suddenly narrowed into a long subulate or filiform often curved acumen, when dry loosely appressed with a patent acumen, distant and patent-open when moist; basal margins recurved; branch leaves long attenuate; inner cells near the costa oblong sub-linear, margined sub-quadrate, the others oval-oblong; costa vanishing in base of acumen; perichaetial leaves nerveless; cells sub-linear: capsule small, oblong, straight; lid conic, sub-obtuse; peristome double, teeth incurved; cilia short or none, basal membrane indistinct; seta smooth, fine, flexuous: dioicous. *Mac. Cat.* 170.—On earth and bark at bases of trees: Revelstoke, B. C.

416. *Platygyrium repens orthocladus* Kindb.—Branches elongate, not curved: all basal leaf cells orange; segments linear, not completely free at base, smooth or denticulate at one side, not shorter than teeth. *Mac. Cat.* 172.—On old logs: Ottawa.

417. *Pylaisæa pseudo-platygyrium* Kindb.—Tufts intricate: stem irregularly divided, or sub-pinnate; branches thick: leaves crowded, upper glossy green, others finally brown, from ovate-oblong base, long acuminate, acumen sub-filiform, often curved, distinctly denticulate: borders recurved to acumen; cells narrow and confluent except the quad-

rate alar ones; costa double, distinct; perichaetial leaves serrulate, abruptly narrowed to long, hair-point-like acumen: capsules straight, sub-erect, cylindrical, wide mouthed; segments free, longer than teeth; lid conic, short apiculate: monoicous. Mac. Cat. 173.—On decayed tree trunks: Lake Nepigon.

418. *Pylaisæa Selwynii* Kindb.—Differs from *P. intricata* in denser, darker green tufts: leaves broader, short-acuminate, reflexed to acumen at one border or both; the short alar and marginal cells more numerous: capsule short oval; segments adhering to two-thirds of teeth. Mac. Cat. 174.—Very abundant on old cedar fences: Ottawa.

419. *Pylaisæa filari-acuminata* C. M. & Kindb.—Agrees with *P. velutina* in leaves filiform acuminate, but acumen distinctly denticulate, often twisted; alar cells more numerous: capsule thicker, oblong; peristomial teeth nearly free from segments; lid not found. Mac. Cat. 174.—On logs subject to inundation: Revelstoke, B. C.

420. *Homalothecium sericeum* Sch.—Stem pinnate, creeping; branch leaves narrow, ovate lanceolate, short-decurrent, long subulate or filiform acuminate, plicate, faintly denticulate nearly all around or sub-entire; margin scarcely reflexed; cells narrow, alar quadrate; costa long, vanishing in base of acumen; perichaetial leaves scarcely plicate, attenuate to a filiform point: capsule erect, sub-cylindric, straight or slightly curved; teeth pale; segments short, basilar membrane high, to one-third the segments; cilia none; annulus broad: dioicous. Mac. Cat. 176.—On bark of trees: Vancouver Island; on rocks: Rocky Mountains.

420a. *Homalothecium sericeoides* C. M. & Kindb.—Differ from *H. sericeum* in branch leaves more densely crowded, not decurrent, narrower, heteromorphous; some long acuminate, sub-entire, less reflexed at margins; others short-acuminate, strongly reflexed at margins to serrulate acumen: capsules and seta unknown. Mac. Cat. 175.—Crevices of granite rocks: Revelstoke, B. C.

421. *Homalothecium Nevadense subulatum* Ren. & Card.—Leaves long acuminate subulate, generally less plicate. Hedwigia 32: 253. 1893. *H. sericeoides* C. M. & Kindb., Mac. Cat. 175.—Washington; Idaho; British Columbia.

422. *Homalothecium corticolium* Kindb.—Tufts dense, glossy: stems pinnate, creeping; branches densely crowded, curved: stem leaves ovate, abruptly narrowed to recurved or straight acumen; branch leaves ovate-oblong, acute or short-acuminate, straight; all leaves more or less denticulate and reflexed all around; alar cells quadrate, not numerous, marginal also quadrate, others oblong-rhomboidal; costa stout, vanishing above middle; perichaetial leaves entire, long-acuminate: capsule cylindrical-oblong, larger, slightly curved; teeth yellow; segments with high basilar

membrane; lid short apiculate: monoicous. Mac. Cat. 274.—On rocks: Vancouver Island.

423. *Cylindrothecium Macounii* (C. M. & Kindb.) Ren. & Card.—Tufts loose: stem sparingly divided, translucent; branches much compressed, elongate, not attenuate, shining green above: leaves patent, concave, short, ovate-lanceolate, attenuate to short subulate point, basal angles rounded; margins scarcely recurved below, faintly denticulate all around; cells faintly chlorophyllose, long sub-linear, lowest basal dilated, oblong, or the alar often sub-quadrata; costa none or very short and double; perichætal leaves small, convolute or connivent, longer acuminate, more distinctly denticulate at apex: dioicous. *Entodon Macounii* C. M. & Kindb., Mac. Cat. 177.—On earth: Ontario.

424. *Cylindrothecium aciculare* (C. M. & Kindb.) Ren. & Card.—Tufts compact, brown-yellow or variegated with green: stem much divided, very radiculose; branches very short, turgid, not attenuate: leaves imbricate, with difficulty loosed from stem, scarcely open when moist, finally golden yellow, from ovate oblong base suddenly narrowed to fine aciculiform or subulate point, denticulate nearly all around; cells not chlorophyllose, linear lanceolate or fusiform, alar not well defined; costa generally wanting: barren. *Entodon acicularis* C. M. & Kindb., Mac. Cat. 176.—Ottawa.

425. *Cylindrothecium expallens* (C. M. & Kindb.) Ren. & Card.—Tufts loose, pale yellow: stems sparingly divided, radiculose at the base; branches elongate, much compressed, not attenuate: leaves sub-distichous, patent, short ovate-lanceolate, acute, concave, nearly entire, denticulate only at apex, distinctly auriculate, not recurved at margins; cells not chlorophyllose, long sub-linear, inner basal dilated sub-oblong; auricles excavate, well defined, with large oval or roundish finally golden yellow cells; costa none or double, sometimes reaching to middle: barren. *Entodon expallens* C. M. & Kindb., Mac. Cat. 177.—In boggy soil in woods: Rocky Mountains.

426. *Climacium dendroides Oregonense* R. & C.—Leaves narrower at base: less serrate at apex, sometimes sub-entire. Bot. Gaz. 15: 59. 1890.—On ground and old logs: Oregon.

427. *Climacium Americanum Kindbergii* Ren. & Card.—A remarkable form, distinct by shorter, more distant, loosely intricate leaves; cells nearly equal, short, ovate, scarcely 1-2 times longer than broad. Bot. Gaz. 15: 59. 1890.—Louisiana; Massachusetts.

428. *Orthothecium intricatum* Hartm.—Stems 2-5 cm. long, delicate, decumbent below, becoming erect, branching somewhat: tufts compact, soft, olive or yellowish green: leaves erect, spreading, subsecund, narrowly lanceolate, long-acuminate, entire, not plicate: capsule erect, oval or

oblong, contracted below mouth; lid conic; membrane of endostome very short; segments a little longer than the teeth; cilia none: very rarely fruiting. Husnot, Musc. Gall. 317.—Rocky Mountains; Greenland.

429. *Pseudoleskea atrovirens brachyolados* Sch.—Plants larger, leaves larger, more shortly acuminate, not secund; cells of middle part shorter and differing little from others: capsule shorter. Husnot, Musc. Gall. 306.—Wyoming.

430. *Pseudoleskea atrovirens filamentosa* Boulay.—Branches elongated, very slender, hooked at the tips: tufts rather loose, pure yellow at the surface: leaves secund, oval-oblong, narrowed into a long linear sharp acumen, scarcely plicate, 1×0.3 mm.; median cells linear, 1:4-8, translucent, lateral cells quadrate, opaque, rather narrow, enlarged at the base: capsule oblong, arcuate. Musc. de la France 1: 162. 1884.—Washington; Oregon; Montana.

431. *Pseudoleskea atrovirens atricha* Kindb.—Tufts very dense, soft-fuscescent, with green tips: stem very lax, nearly without paraphyllia and rhizoids. Mac. Cat. 180.—On rocks: Griffin Lake, B. C.

432. *Pseudoleskea rigescens denudata* Kindb.—Stem more rigid, nearly simple, curved only at apex, naked below and radiant: barren. Mac. Cat. 181.—Selkirk Mountains, B. C.

433. *Pseudoleskea falcisnps* C. M. & Kindb.—Plants densely tufted: stems much branching: leaves very papillose, denticulate above, short, ovate-lanceolate, attenuate to an acute or filiform often curved point, long decurrent; margins recurved to or above middle, not in upper part; cells rotundate, quadrate at angles; costa vanishing far below acumen: dioicous. Mac. Cat. 182.—On rocks: Sicamous, Revelstoke and Quesnel, B. C.

434. *Pseudoleskea stenophylla* Ren. & Card.—Dioicous (monoicous?), male flowers small, numerous, gemmiform: tufts very intricate, yellowish: stem slender, tough, prostrate, strongly radiculose, irregularly pinnate; branches slender, attenuate, ascending: leaves erect-spreading, narrowly lanceolate, gradually long acuminate, entire or obsoletely denticulate above; margin revolute below; costa strong, vanishing in the green acumen; cells thickened, alar quadrate numerous, median sub-linear truncate, upper rhomboidal ovate, apical large and obtusely papillose; paraphyllia numerous, triangular lanceolate or subulate; inner perichætal leaves sheathing, oblong, rather suddenly acuminate, entire, costa very smooth, vanishing in acumen: seta red, smooth, above twisted to right, 6-8 mm. long; capsule erect, inclined or pendulous with age, ovate-oblong, red-brown, slightly constricted below mouth; peristome teeth brown, linear-lanceolate, lamellose within; segments from a short basilar membrane narrowly linear-subulate, entire, equaling teeth. Bot. Centralbl. 44: 421. 1890.—Cascade Mts., Easton, Washington.

435. *Pseudoleskea tectorum* Sch.—Stems 1-5 cm. long, decumbent,

delicate; branches numerous, short: tufts depressed, very dense, dark or reddish green: leaves erect-spreading when moist, imbricate when dry, broadly oval lanceolate, acuminate, entire; costa delicate, often unequally bifid, vanishing toward middle; margin of 3-5 rows of rounded transversely elongated cells in lower part, the remainder oblong; inner perichætal leaves sheathing, lanceolate, abruptly narrowed into a narrow point, longitudinally plicate: dioicous: capsule erect, cylindric, ferruginous; operculum convex-conic, rostrate; teeth of peristome, short, yellow; endostome with narrow segments, a little shorter than teeth, constricted at articulations giving them a moniliform aspect, basilar membrane and cilia absent. Husnot, Musc. Gall. 302.—Greenland.

436. *Pseudoleskea malacoclada* C. M. & Kindb.—Plants densely tufted: stems not rigid, much branching, sparingly radicate, without paraphyllia; branches green or finally fuscous: leaves crowded, entire, not decurrent, not distinctly papillose; stem leaves from a concave broadly ovate or rotundate base suddenly narrowed to subulate or filiform recurved acumen; branch leaves more attenuate to subulate and patent point; basal margins recurved; cells rotundate, alar quadrate; costa very short and double or none; perichætal leaves loosely connivent with a patent or curved acumen: dioicous. Mac. Cat. 182.—On dry rocks: British Columbia.

437. *Heterocladium heteropterum* Sch.—Stems 2-6 cm. high, filiform, decumbent, delicate, giving off irregular ascending branches: tufts dark-green: leaves spreading or sub-secund when moist, loosely imbricate when dry, papillose on both sides; stem leaves oval, acuminate, denticulate all around; costa faint, bifurcate, short; cells rounded-quadrate, oblong-linear in the middle; branch leaves smaller: capsule horizontal, oblong, neck distinct; lid conic, rostrate: rarely fruiting.—Husnot, Musc. Gall. 307. On damp rocks: British Columbia; Vancouver Island; Greenland.

438. *Heterocladium Vancouveriense* Kindb.—Plants minute, densely and irregularly pinnate, sparingly radiculose, rarely flagelliferous, dark green, not glossy; branches short, very slender: leaves ovate-deltoid, denticulate above middle, pellucid, faintly papillose; stem leaves decurrent, reflexed at base, acute; branch-leaves smaller, looser, more concave, oval and obtuse; cells hexagonal oval or rhombic, alar and marginal quadrate-rhombic; alar and marginal quadrate-rhombic; costa slender, short, scarcely reaching to middle; paraphyllia few or none; perichætal leaves longer acuminate, cells narrower: capsule small, oblong, sub-erect; peristome perfect, teeth yellowish, cilia 2, elongate: dioicous. Mac. Cat. 183. On rocks: Vancouver Island.

439. *Heterocladium frullanopsis* C. M. & Kindb.—Stem irregularly divided, brown red, not radiculose; branches short, obtuse, when dry incurved: paraphyllia none; leaves uniform, dark-green, not shining, when dry loosely or not at all appressed, sub-squarrose when moist, distant and

decurrent, cochleariform, roundish oval, obtusate, incurved at the apex, minutely denticulate above to middle, very papillose; upper cells rhombic, lower oblong, alar quadrate, inner basal reddish; costa short, more or less distinct, sometimes stout and simple: dioicous. Mac. Cat. 183.—On trees: New Brunswick.

440. *Heterocladium aberrans* Ren. & Card.—Intricate-cespitose, pale or yellowish-green: stems flexuous, creeping, radiculose, more or less regularly pinnate; branchlets ascending, flexuous: stem leaves squarrose, auriculate, from a cordate-ovate base, long-acuminate, generally subulate; borders plane, sinuate-crenulate all around; costa forked, with one division longer and vanishing about middle; areolation loose, pellucid, of soft thick-walled cells, elongated, linear, truncate or obtuse, 4-10 times longer than broad toward costa, others irregular ovate, roundish or sub-hexagonal, sometimes slightly papillose; branch leaves shorter, acute or obtuse; perichaetial leaves acuminate to a reflexed denticulate point, thin-nerved: seta purple, smooth; capsule horizontal, ovate, curved; lid unknown; teeth yellow, acuminate, densely trabeculate; segments narrowly split, cilia shorter, nodulose. Bot. Gaz. 15: 59. 1890.—On logs: Idaho.

441. *Thuidium setum lonchoneuron* Kindb.—Stems irregularly divided, sparingly radiculose; branchlets thick: stem leaves more broadly revolute at borders below acumen, less papillose: costa stout, percurrent or excurrent; branch leaves quite acute: capsules not found. Mac. Cat. 194.—On old logs: Leamington, Ont.

442. *Thuidium microphyllum lignicolum* (Kindb.) Best.—Monoicous: tufts yellowish or bright green: stems simply pinnate with few rhizoids and short scarcely ramose paraphyllia; branches close, distichous, attenuate, flexuous or slightly recurved: stem leaves from broad cordate base attenuate to a long often curved point, faintly striate, reflexed on borders; branch leaves shorter, acuminate; all denticulate from middle upward, and papillose at back or on both sides; cells obscure, rounded; costa vanishing in or below apex: capsule cylindrical, arcuate, light brown; teeth pale; cilia long, perfect; annulus double; lid conical, short apiculate. *T. lignicola* Kindb. Mac. Cat. 185.—On rotten logs: Ontario and British Columbia.

443. *Thuidium paludosum elodioides* (Ren. & Card.) Best.—Throughout similar in habit to *Elodium paludosum*; leaves short acuminate, cauline leaves fimbriate at base; cells shorter, elliptical or oval, papillose. Differs from *Thuidium Blandovii* by more slender habit: stems more remotely and less regularly branched: stem leaves narrower; cells looser; paraphyllia shorter; perichaetial leaves narrow, entire, very long subulate: monoicous: sterile. *Thuidium elodioides* R. & C., Hedw. 32: 251. 1893.—Hobart, Indiana; New Bremen, Ohio.

443a. *Thuidium Philiberti* Limpr.—Plants medium sized, yellow to dark green, in intricate mats; stem 4-8 cm. long, creeping, pinnately

branched; branches pinnate or bipinnate; paraphyllia multiform: stem leaves triangular-cordate-acuminate, usually with a hyaline filiform point; margins revolute or recurved, at least below; cells oblong-quadrate to oblong-rhomboidal: dioicous; perichaetial bracts loose, flexuous-spreading or reflexed, serrate, rarely with a few short cilia on the innermost, acumen about 3 times as long as the short scarcely costate body: capsule oblong-cylindric, curved horizontal; annulus narrow, indistinct, tardily or imperfectly deciduous; operculum conic-rostrate, curved; spores 12-16 μ , almost smooth, mature in Oct.—In swampy places on ground or base of small trees: New Jersey; Pennsylvania.

444. *Tripterocladium rupestre* Kindb.—Plants intricate, caespitose, pale brownish-green, not glossy: stems irregularly divided; branchlets short, filiform, not creeping: leaves small, densely crowded, when dry appressed, open-erect when moist, scarious, pellucid, not chlorophyllose nor papillose, sub-ovate, acute or long acuminate, upper denticulate above, borders recurved below, sometimes to middle or above; cells in lower leaves sub-linear above, looser at base, in upper leaves looser, oblong linear, in all sub-quadrate at borders and base; costa none. Mac. Cat. 187.—On the face of a cliff: Yale, B. C.

445. *Camptothecium lutescens occidentale* R. & C.—Robust, branches strongly sericeous, capsule narrower, longer. Hedwigia 82: 254. 1893.—Washington.

446. *Camptothecium dolosum* Ren. & Card.—Sub-species of *C. aeneum*: branch leaves shorter, broader; nerve sometimes dilated at apex and denticulate; inner perichaetial leaves with apex suddenly truncate, lacerate or deeply incised dentate. Hedwigia 82: 255. 1893.—Washington.

447. *Camptothecium aureum* Sch.—Plants smaller and more delicate than in *C. lutescens*; branches more crowded and shorter; tufts golden yellow: leaves shorter and less deeply plicate; cells of basal angles rounded-quadrate, forming distinct auricles; paraphyllia quite numerous, ordinarily oval lanceolate; perichaetial leaves entire: pedicel scabrous; capsule oblong, contracted below mouth; lid conic, obtuse, teeth of peristome orange. Husnot, Musc. Gall. 319.—Vancouver Island.

448. *Camptothecium Amesiae* Ren. & Card.—Widely caespitose, bright yellowish green: stems prostrate, creeping, radiculose, pinnately ramulose; branchlets crowded, short, equal, erect, a little curved: stem leaves broadly triangular, narrowly long acuminate, carinate, plicate, generally plane on one side and revolute on the other, slightly serrulate at apex; costa vanishing in acumen: cells linear, attenuate, upper shorter, alar numerous, quadrate or subrectangular; inner perichaetial leaves ecostate, narrowly lanceolate, long acuminate, entire: seta short, purple, rough, a little twisted to left; capsule narrow, long cylindric, sub-erect, or sub-horizontal and slightly arcuate; lid unknown; teeth orange, long

acuminate-subulate, strong, densely trabeculate; segments split their whole length; cilia long. Bot. Gaz. 17: 202. 1888.—Mixed with *Hypnum pinnatifidum*, California.

449. *Camptothecium Nuttallii tenue* Kindb.—Branches longer, filiform: leaves smaller only at base, hooked denticulate at apex. Mac. Cat. 189.—Perpendicular rocks: British Columbia.

450. *Brachythecium laetum fallax* R. & C.—Branches julaceous, elongated: leaves narrower, longer acuminate, more or less revolute; alar cells soft, hardly incrassate. Hedwigia 32: 257. 1893.—Calumet River, Ind.

451. *Brachythecium laetum Roellii* Ren. & Card.—Stems depressed, pinnate; branches julaceous, short, obtuse: leaves densely imbricate, broader, concave, shorter acuminate: alar cells as in preceding variety. l. c.—Calumet River, near Hobart, Ind.

452. *Brachythecium laetum pseudo-acuminatum* Ren. & Card.—Denticulate, habit *B. acuminatum*: leaves deeply plicate, basal angles excavate; alar cells less numerous but distinct. l. c.—Calumet River, near Hobart, Ind.

453. *Brachythecium digastrum* C. M. & Kindb.—Tufts laxly cohering, olive-green, not shining: stem rigid, sub-pinnate or irregularly branching, radiculose below; branches sub-julaceous, obtusate: stem leaves when dry loosely appressed or sub-imbricate, crowded, patent or sub-secund when moist, decurrent, not auricled, plicate, biventrose, ovate, short acuminate with flexuous acumen, or when dry serpentine-corrugate; borders more or less recurved but not reflexed, sub-entire or faintly denticulate above; lower basal cells wide and sub-rhombic, alar rather quadrate-rectangular and not very distinct, upper confluent small very chlorophyllose, inner median sub-linear, others oblong-lanceolate; costa thick and sub-flexuous, long, vanishing near acumen; branch leaves ovate oblong, more distinctly revolute at borders, denticulate at acumen, narrower areolate: capsule asymmetric, sub-cylindric, curved; lid long conic; seta smooth; teeth of peristome conic connivent when moist, cilia nodulose, not appendiculate: monoicous. Mac. Cat. 190.—Ottawa, Ont.; New Brunswick.

454. *Brachythecium Fitzgeraldi* C. Müll.—Dioicous: tufts low, pulvinate, broad, yellow, loosely interwoven: stem with branches short, more or less parallel, slender, round-julaceous; branchlets very short, rather spreading, single: stem leaves closely appressed, when moist scarcely spreading with cordate base semicircularly impressed, rather broad ovate, short acuminate; more or less ventricose-concave on both sides of the narrow vanishing deeply canaliculate green costa; margin nearly plane, everywhere slightly denticulate; cells very narrow, long, pale yellow; alar cells many, small, hexagonal: fruit unknown. Flora 70: 224. 1887.—Florida.

455. *Brachythecium acuminatum sub-albicans* R. & C.—Facies of *B. albicans*: more robust, pale yellow: branches silky, julaceous: areolation denser, cells narrower. Bot. Gaz. 15: 60. 1890.—Louisiana; Florida.

456. *Brachythecium cyrtophyllum* Kindb.—Sub-species of *B. acuminatum*: habit of a small form of *B. albicans*: plants caespitose, green, faintly glossy: stems irregularly divided, not creeping; branchlets filiform, sub-obtuse: leaves small, close, loosely appressed when dry, open-erect when moist, ovate-acute or short-acuminate, not sulcate nor decurrent, serrulate at least above middle; areolation loose, upper cells narrowly rhomboidal, inner sub-linear, alar sub-quadrate somewhat numerous and chlorophyllose; costa stout, reaching to $\frac{3}{4}$; perichaetial leaves ecostate: dioicous. Mac. Cat. 191.—On elm logs in thick woods: Ontario.

457. *Brachythecium Roellii* Ren. & Card.—Dirty or yellowish green: stems soft, depressed, scarcely radiculose, subpinnate; branches elongate, flexuous: leaves ovate-lanceolate, decurrent, quite long and narrowly acuminate, faintly plicate; margin sinuate or denticulate, plane in middle, revolute at base and at acumen; costa reaching beyond middle to $\frac{3}{4}$ length; cells narrow, elongate, linear, alar cells few sub-quadrate: evidently dioicous. Hedwigia 32: 263. 1893.—Vancouver.

458. *Brachythecium glareosum* Sch.—Related to *B. salebrosum*: stems decumbent, often very long, 15-20 cm., forming large yellow tufts, softer, shining: leaves more erect, longer acuminate; cells narrower, those of angles oblong: dioicous: capsule oblique or sub-horizontal; lid conic, longer. Husnot, Musc. Gall. 324.—On earth at roots of trees: Revelstoke, B. C.

459. *Brachythecium albicans occidentale* Ren. & Card.—Stems depressed, laxly foliate: leaves subsecund, less long acuminate, sometimes very distinctly denticulate. Hedwigia 32: 258. 1893.—Washington; Montana.

460. *Brachythecium harpidioides* C. M. & Kindb.—Tufts compact, soft, radiculose below, whitish or bright green, not shining: stems intricate, irregularly branching or pinnate: leaves spreading, somewhat loosely disposed, decurrent, more or less arcuate, not auricled, plicate, nearly flat, ovate, subulate acuminate; borders faintly denticulate, recurved often all around; cells distinctly chlorophyllose, alar large sub-quadrate, others lanceolate: dioicous: capsules not found. Mac. Cat. 194.—On old logs in woods: Revelstoke, B. C.; New Brunswick.

461. *Brachythecium salebrosum Waghornii* R. & C.—Monoicous: tufts very dense: stems erect, turgid, not radiculose: leaves more crowded, imbricate: lid mamillate. Differs from *B. mamilligerum* Kindb. in leaves imbricate, not patulous when dry: stems not radiculose. Bot. Gaz. 19: 238. 1894. Battle Harbor, Labrador.

462. *Brachythecium mamilligerum* Kindb.—Sub-species of *B. salebrosum*: monoicous: tufts very dense, radiculose: stem sub-pinnate: leaves very patent when dry, striate, long, narrowly ovate-lanceolate often filiform pointed; borders not or narrowly recurved, faintly serrate; alar cells few; costa generally vanishing in middle, sometimes longer; perichæatial leaves long aristate: capsule sub-oval, curved; segments narrowly rimose, not open; cilia radiose, not appendiculate; annulus none; lid mamillate. Mac. Cat. 192.—On old wet logs: Sicamous, B. C.

463. *Brachythecium salebrosum turgidum* Hartm.—Habit of *B. glareosum*: branches long, sericeous-yellow, leaves appressed, entire. Hartm. Skand. Fl. 2: 16. 1871.—In peat bogs: Stephen, Rocky Mts.; Greenland.

464. *Brachythecium pseudo-albicans* Kindb.—Differs from *B. albicans* in leaves looser, sub-distichous, shorter acuminate, faintly striate, denticulate all around; alar cells greater, all basal cells finally reddish: barren. Mac. Cat. 194.—On earth in woods: Vancouver Island.

465. *Brachythecium spurio-acuminatum* C. M. & Kindb.—Differs from *B. acuminatum* in tufts lax, loosely adhering to substratum: leaves denticulate, recurved at borders nearly all around; alar cells still more numerous and very chlorophyllose; perichæatial leaves subulate-acuminate, not filiform pointed: inflorescence monoicous. Mac. Cat. 191.—On logs in woods: Ontario.

466. *Brachythecium erythrorrhizon* Sch.—Monoicous: intricate cespitose: stems creeping, divided, strongly radiculose; branches ascending incurved; branchlets short: leaves crowded, laxly imbricate, younger secund, broad ovate-lanceolate, long acuminate, unequally broad sulcate; perichæatial leaves loosely imbricate: capsule cernuous short-ovate, incurved. Schimp. Syn. 2: 646.—Washington.

467. *Brachythecium sub-erythrorrhizon* Ren. & Card.—Monoicous: intricate-cespitose, yellowish green, facies of *B. velutinum*: stems creeping, radiculose, sparingly branching, branches procumbent: leaves sub-homomallous, oblong-lanceolate, long narrowly acuminate, bi-triplicate, sharply serrate all around; borders plane or partly revolute; costa vanishing above middle; sometimes forked and shorter; cells rather loose, pellucid, rhomboideal linear, alar quadrate sub-obscure; perichæatial bracts lanceolate, long acuminate-subulate, acumen serrate: seta smooth, reddish; capsule sub-erect, turgid ovate, not or scarcely constricted below mouth when dry; lid unknown; teeth of peristome yellowish, triangular-lanceolate; segments narrow, split along divisural line; cilia 1 or 2, long, filiform. Bot. Gaz. 19: 238, 1894.—Colorado.

468. *Brachythecium pseudo-collinum* Kindb.—Agrees with *B. collinum* in pinnate and creeping stem: decurrent leaves denticulate all around: short sub-oval capsule and smooth pedicel: differs in stems julaceous: leaves

larger and longer, ovate-lanceolate, more loosely disposed, spreading or patent, gradually tapering into short, half-twisted acumen; cells chlorophyllose, alar ones larger; costa longer, reaching above middle: monoicous. Mac. Cat. 196.—Queen's Co., New Brunswick.

469. *Brachythecium laevisetum* Kindb.—Habit of *B. populeum*: plants caespitose: green, glossy: stem irregularly ramulose: leaves close, sub-erect, open, ovate-lanceolate, acuminate and acute, sulcate; borders serrulate from middle upwards; basal cells dilated; costa long, reaching at least middle of leaf; perichætical leaves nerveless: capsule small, incurved, oblong; outer teeth hyaline-margined, light brown; cilia short, not appendiculate; lid highly convex, not apiculate; pedicel very smooth: monoicous. Mac. Cat. 193.—On rotten logs: Gold Range, B. C.

470. *Brachythecium Idahoense* Ren. & Card.—Imbricate-caespitose, bright green: stems depressed, creeping, irregularly pinnate; branches ascending, sub-incurved; leaves crowded, sub-secund, from an ovate base lanceolate, long acuminate, plicate, costate to above middle, borders denticulate all around or sub-entire, plane or more or less revolute; cells linear, attenuated, those of angles sub-quadrate, numerous; perichætical leaves rather suddenly acuminate, obsoletely costate or sub-ecostate: seta purple, smooth; capsule horizontal, ovate, gibbous, curved; lid obtusely conic; teeth lanceolate acuminate, densely trabeculate; segments broadly split; cilia long, nodulose: monoicous. Bot. Gaz. 15: 60. 1890.—On logs: Idaho.

471. *Brachythecium Neve-Angliæ Delamarei* R. & C.—Stems shorter, almost simple: leaves more distinctly imbricate, abruptly contracted into a short point. Fl. Miq. 50.—Island of Miquelon.

472. *Brachythecium latifolium* (Lindb.) R. & C.—Near *B. rivulare*: plants dioicous: very much smaller, straight, acute, acuminate at apex; irregularly and remotely sub-pinnate to sub-simple; branches short, divaricate, acute, simple: stem leaves pellucid, spreading, very long and broadly decurrent, rhomboid-triangular, gradually long acuminate, concave, not plicate; margin recurved below broadest part, very often auriculate; costa slender, vanishing at middle or a little above; cells nearly three times shorter, large, rhomboid-prosenchymatous, not vermicular: fruit not seen. *Hypnum latifolium* Lindb., Musci Scand. 35.—Miquelon Is.

473. *Brachythecium pseudo-Starkel* Ren. & Card.—Dioicous: loosely caespitose, green: stem erect or ascending, flexuous, laxly pinnate, branches elongate, attenuate: leaves not close, patulous, ovate-lanceolate, plicate acuminate, acumen long, sometimes tortuous; margin generally serrate; costa extending into the acumen; cells linear-rhomboidal, elongate, attenuate, alar lax soft quadrate hyaline: not fruiting. Bot. Cent. 44: 423. 1890.—Washington.

474. *Brachythecium rivulare obtusulum* Kindb.—Stem irregularly divided; branches simple and elongate: leaves glossy, ovate, blunt or

short acute, striate, decurrent, indistinctly denticulate above or from middle; cells dilatate, principally the lower and the uppermost, alar and basilar finally orange-reddish, alar rarely greater, costa short and simple. Mac. Cat. 201.—New Brunswick: Ontario; Revelstoke, B. C.

475. *Brachythecium platycladum* C. M. & Kindb.—Tufts densely cohering, bright green, shining; stem irregularly branching; branches short obtuse, complanate: leaves loosely imbricate or patent, nearly flat, long decurrent, distinctly auriculate, faintly striate, broad, ovate, suddenly and generally short acuminate; borders not recurved, faintly sinuate or subentire below middle, more distinctly denticulate above; cells pale, upper narrow, lower near base dilated, alar large and well-defined; costa short, reaching little above middle: capsule sub-oval, faintly curved; teeth dark yellow, entire at borders; cilia not appendiculate; lid unknown: dioicous. Mac. Cat. 185.—On stones: Ottawa, Ont.

476. *Brachythecium spurio-rutabulum* C. M. & Kindb.—Differs from *B. rutabulum* in dioicous inflorescence: leaves distinctly plicate, longer cuspidate: seta rough, short. Tufts dense: stems pinnate, and creeping leaves shining, when dry very spreading, loosely disposed, long decurrent borders recurved below the middle, faintly denticulate. Mac. Cat. 197 and Sicamous.—On bases of trees: Burrard Inlet, B. C.

477. *Brachythecium nanopes* C. M. Kindb.—Allied to *B. populeum* in habit, peristome, monoicous inflorescence, pedicel faintly rough above, long and subpercurrent costa of leaves: differs in stems not creeping, nearly without rhizoids, branches unilateral, leaves smaller and narrower at base, stem leaves long filiform apiculate, sub-entire and not or indistinctly recurved at borders, lower decurrent: capsule smaller, pedicel shorter, peristome pale orange, cilia long and indistinctly appendiculate. Mac. Cat. 201. On earth: Revelstoke, B. C.

478. *Brachythecium trachypodium* Sch.—Stems 3-5 cm. high, decumbent, furnished with branches and branchlets: tufts golden yellow, or greenish in part: leaves erect, loosely imbricate, oval or long lanceolate, long acuminate, denticulate, costate to middle, faintly plicate; cells of basal angles quadrate, others linear: some perichaetial leaves abruptly, others gradually acuminate: monoicous: seta robust, very papillose; capsule almost erect, or oblique, oval or oblong, contracted below mouth; lid convex-conic, obtuse; peristome of *B. velutinum*; calyptra reaching base of capsule. Husnot, Musc. Gall. 328.—Greenland.

479. *Brachythecium reflexum Pacificum* Ren. & Card.—More robust, stem leaves larger, ovate-lanceolate, hardly triangular, somewhat acuminate, margin sub-revolute at base. Hedwigia 82: 262. 1893.—Mt. Hood, Oregon.

480. *Brachythecium reflexum Demetrii* Ren. & Card.—Habit stronger,

branches thicker, erect, leaves broader, softer. Bot. Gaz. 19: 239, 1894. Squaw Is., Labrador.

481. *Brachythecium glaciale* Sch.—Stems 2-5 cm., decumbent, much divided, branchlets subglaucescent: green or dark yellow: leaves erect-imbricate, decurrent; stem leaves broadly oval, abruptly or shortly acuminate, denticulate throughout, costate $\frac{3}{4}$ length, plicate; branch leaves narrower, longer acuminate; cells of the basal angles quadrate or rectangular, the middle linear of variable length; perichaetial leaves erect-imbricate: monoicous: seta papillose; capsule almost erect, oblique or horizontal, oval or oblong; cilia nodulose, not appendiculate. Husnot, Musc. Gall. 323.—Greenland.

482. *Brachythecium curtum* Lindb.—From *H. Starkei* differs in its leaves broad and short; margin plane and short serrate; costa more slender, much shorter, smooth on back; cells broader. Differs from *B. cecidipodium* in perichaetial bracts shorter acuminate, more serrate: seta rather thick, slightly scabrous; capsule nodding more than horizontal; lid perfectly conic. Musci Scand. 35. 1879.—On earth in woods: New Brunswick; Prince Edward Isl.; Ottawa.

483. *Brachythecium gemmascens* C. M. & Kindb.—Tufts very dense, finally green, rufescent: stem irregularly branching, furnished with numerous male buds: leaves narrow, ovate-lanceolate with a twisted point, faintly denticulate all around, crowded, not decurrent nor auricled; cells pale, nearly all narrow, only lowest one or two basal rows dilated, alar not larger than inner; costa prolonged above middle, vanishing below acumen: capsule small, round-oval, oblique; teeth dark-orange; segments shorter than very high basal membrane, cilia not appendiculate; lid conic, short pointed; seta minutely verrucose, about 1 cm. long; monoicous. Mac. Cat. 195.—On wet logs: Columbia River, above Revelstoke, B. C.

484. *Brachythecium leucoglaucum* C. M. & Kindb.—Tufts loose with but few rhizoids, whitish or sub-glaucous-green, faintly shining: stem sub-pinnate or irregularly branching: leaves from ovate base suddenly tapering into a somewhat long filiform often half twisted point, sharply serrate above, faintly denticulate below, striate, decurrent, borders reflexed below; alar cells small, numerous, others narrow; costa long, sometimes sub-percurrent; perichaetial leaves long, filiform-acuminate, arcuate-squarrose: capsule curved, oblong-cylindric; lid sub-obtuse when moist; pedicel faintly rough; peristome conic-connivent, teeth serrulate, pale above; segments open in middle; papillose above, cilia not appendiculate; monoicous. Mac. Cat. 197.—On loose earth: New Brunswick.

485. *Brachythecium rutabulum Canadense* Ren. & Card.—Of more delicate habit; leaves narrower, deeply plicate, long acuminate. Revue Bryol. 20: 19. 1893.—Canada; Miquelon Is.; Washington.

486. *Brachythecium rutabuliforme* Kindb.—Agrees with *B. rutabulum* in very rough pedicel, monoicous inflorescence and form of leaves: differs principally in rigid stem, distinctly appendiculate cilia of peristome, and short pedicel: leaves sub-ovate, short-acuminate, nearly entire, faintly denticulate. Mac. Cat. 198.—On stones in brook: British Columbia.

487. *Brachythecium Columbicorutabulum* Kindb.—Tufts dense, faintly shining, finally brownish-green: stems elongate, pinnate creeping: stem leaves patent, from broadly ovate base long cuspidate, with filiform generally prolonged point, decurrent, very plicate, nearly entire, borders shortly reflexed below; cells not chlorophyllose, alar larger and well distinct; perichætal leaves sub-erect-patent with a long filiform arcuate point, faintly and distinctly denticulate: capsule oblong-cylindric, curved; peristome teeth not serrulate, segments very open in middle, cilia faintly nodulose, not appendiculate; pedicel very rough; monoicous. Mac. Cat. 198.—On wet and rotten logs in woods: Columbia River, B. C.

488. *Brachythecium lamprochryseum* C. M. & Kindb.—Tufts large, laxly cohering to substratum, with few rhizoids: golden yellow, shining or finally decolorate: stem elongate, often pinnate; branches generally short or sometimes more elongate and faintly curved above, sub-acute: leaves open, more or less loosely disposed, long decurrent, distinctly auriculate, very plicate, from triangular ovate base short acuminate, filiform or subulate cuspidate, often curved at apex, nearly flat, only auricles faintly revolute; borders faintly denticulate all around; most cells very long and narrow, lowest basal ones dilated and short, also the alar, all very sparingly chlorophyllose; costa broader at base, faint, reaching to middle, but in the smaller narrower and more loosely disposed leaves of some branchlets longer, reaching to acumen: perichætal leaves ecostate, when dry squarrose, inner sheathing with a short subulate acumen and a long filiform point: capsule short, sub-ovoid, thicker near base, slightly contracted below mouth, arcuate; lid unknown; teeth finally brown at least at base; cilia not appendiculate; pedicel very rough: monoicous. Mac. Cat. 199.—On stones: Vancouver Island.

489. *Brachythecium mirabundum* C. M. & Kindb.—Tufts large, very laxly cohering, nearly without rhizoids, silky or yellowish green, faintly shining: stem elongate, irregularly divided or prolonged into sciuroid-curved obtuse branches: leaves loosely imbricate, crowded, when dry sub-rugose, when moist patent, short decurrent, indistinctly auriculate, faintly plicate, from concave ovate and gradually acuminate base long cuspidate; borders broadly recurved at least at one side of nearly entire base to involute and distinctly denticulate acumen; cells pale, elongate, narrow, alar sub-quadrate and not much wider than other basal ones, all sparingly chlorophyllose; costa vanishing in acumen: perichætal leaves ecostate,

longer filiform-cuspidate, irregularly sinuolate, point patent or arcuate, basal cells larger rectangular: capsule small, at base distinctly gibbous, narrow, cylindric, curved; lid elongate-conic; pedicel very short, very faintly muriculate: monoicous. Mac. Cat. 199.—On old logs in woods: New Brunswick.

490. *Brachythecium Villardi* Ren. & Card.—Monoicous: similar in habit to *B. salebrosum*: stem depressed, creeping, radiculose, pinnate, branches ascending: leaves erect, imbricate or sub-secund, decurrent, ovate lanceolate, long and slenderly acuminate, plicate, margin more or less revolute, entire or in acumen faintly denticulate; costa reaching two-thirds length of leaf; cells narrow, elongated, linear, alar cells few, quadrate: pedicel rough: remaining characters not known. Bot. Centralbl. 44: 422. 1890.—Washington.

491. *Brachythecium cirrhosum* Sch.—Stems decumbent, 3-10 cm. long, stoloniferous, more or less branching; branches ascending or erect, inflated, julaceous: tufts yellowish or golden green: leaves imbricate, very concave, sub-cochleariform, oval oblong, very abruptly contracted into a very long filiform point, shining, plicate when dry, often inflexed on the borders, denticulate in the upper part, sometimes entire; costa simple or double, vanishing near middle; basal cells quadrate or rectangular, rounded, others linear: fructification unknown. Husnot, Musc. Gall. 338.—Greenland.

492. *Scleropodium caespitosum sublaeve* Ren. & Card.—Pedicel nearly smooth, slightly rough only below capsule. Bot. Gaz. 15: 61. 1890.—Sauvie's Island, Oregon.

493. *Scleropodium Krausei* (Müll.) Ren. & Card.—Monoicous: tufts low, rather robust and loose, pale green: stem sparingly branched; branches rather short, flexuous, round julaceous, turgescens, with obtuse apex: stem leaves closely or loosely crowded, narrowly oblong-lanceolate, cymbiform-concave, short pointed, point somewhat twisted, quite entire; base truncate, margin plane; costa very slender, yellowish, vanishing, often bifurcate; cells very narrow, vermicular, alar many small pellucid; perichætal leaves larger; seta rather short, red, smooth; capsule amblystegoid-cylindric, cernuous, coriaceous, ochraceous; lid conic, very short mamillate; annulus simple, rather broad; peristome teeth robust, broad, long, yellow, hyaline pointed, cristate; segments from a rather high yellow membrane, long, broad, sulcate, very smooth, not perforate nor gaping, cilia rudimentary, solitary. *Hypnum Krausei* C. Müll., Flora 70: 224. 1887.—Alaska.

494. *Isoetecium Cardoti* Kindb.—Rhizome creeping; secondary stems pinnate, ligneous, often curved, sometimes bearing rigid long and at the apex branching flagella: stem leaves ovate-lanceolate, subulate acuminate, faintly denticulate below, acumen serrate and twisted above; cells often yellowish, all long and narrow except basal, the alar and inner basal dark:

yellow or orange quadrate or rectangular; costa stout, reaching $\frac{3}{8}$ length of leaf; branch leaves shorter acuminate, sharply serrate above middle, borders often faintly reflexed to acumen; perigonal leaves sub-ovate, reddish at base; costa fine and short; perichætal leaves ecostate, from a short ovate base suddenly tapering to much longer, subulate acumen: capsule oval, horizontally patent or cernuous, teeth pale yellow; segments rimose in middle, prolonged into long and smooth cilia; seta smooth, arcuate above: monoicous. Bull. Torr. Bot. Club 17: 278.—On bases of trees and logs: Washington; Vancouver; British Columbia.

495. *Isoetium myurellum* Kindb.—Tufts very loose, dark green, not glossy; stems creeping; branches erect, tree-like and ramose; branchlets curved, attenuate; leaves of branches appressed when dry, small, ovate, blunt or short acuminate, twice serrate or denticulate all around, smooth, scarcely reflexed on borders; alar cells round quadrate, middle ones narrow, upper rhombic; costa three-fourths length of leaf: perichætal leaves ecostate, oblong, with a long denticulate horizontally patent acumen and narrow cells: capsule oval oblong, inclined; segments with a low basilar membrane and two short cilia; annulus double; lid conical acuminate; pedicel smooth: dioicous. Bull. Torr. Bot. Club 12: 278.—On decaying logs and on rocks: Vancouver Island; British Columbia.

496. *Eurhynchium strigosum* Barnes! Ren & Card.—Stems rather more robust, stem leaves larger, longer acuminate, branch leaves more elongated: capsule shorter broadly ovate. Bot. Gaz. 14: 97. 1889.—On logs: Lake Pend d'Oreille, Idaho.

497. *Eurhynchium strigosum fallax* Ren. & Card.—Form robust, resembling in habit *E. myosuroides*: stem leaves very large, triangular-lanceolate, obtuse; branch leaves rounded at apex: capsule like that of typical form. Bot. Gaz. 14: 98. 1889.—On logs: Lake Pend d'Oreille, Idaho.

498. *Eurhynchium sub-strigosum* Kindb.—Differs from *E. strigosum* in distant branches complanate: leaves long decurrent and twice greater, patent, sub-distichous: capsule very constricted below orifice; cilia apppendiculate: monoicous. Mac. Cat. 205.—On rocks: British Columbia.

499. *Eurhynchium crassinervium laxifretum* Kindb.—Leaves nearly entire or faintly denticulate above, shorter acuminate; cells larger: only male flowers found. Mac. Cat. 207.—On earth in woods: Queen's Co., N. B.

500. *Eurhynchium colpophyllum flagelliforme* Barnes.—Leaves lance ovate, small; branches long, almost flagelliform, attenuate. Bot. Gaz. 16: 207. 1891.—California.

501. *Eurhynchium Dawsoni* Kindb.—Stems pinnate, not or rarely radiculose; branchlets patent: leaves green or brownish, not glossy, not

or indistinctly papillose, not long-acuminate from broad ovate base, recurved on borders below, long decurrent, open erect, denticulate all around; areolation variable, often sub-rhomboidal; costa thick, reaching nearly to apex: probably dioicous. Bull. Torr. Bot. Club 17: 278.—On rocks: Vancouver Island; British Columbia.

502. *Eurhynchium semiasperum* C. M. & Kindb.—Plants loosely tufted, green: secondary stems sparingly radiculose, branching at one side; branches generally simple, acute: leaves sub-patent, from a broad-cordate base, fine acuminate, decurrent, nearly entire; alar cells numerous, reaching to costa, other cells narrower, the lower dilated; costa vanishing above middle: perichaetial leaves sub-oblong, short acuminate, erect, entire, ecostate: capsule small, sub-oval erect or inclined; teeth papillose above; segments shorter than high basilar membrane; lid narrow, short rostellate; pedicel rough at least to middle, smooth below: monoicous. Mac. Cat. 207.—On rocks in a brook, British Columbia.

503. *Eurhynchium Sullivantii Holzingeri* Ren. & Card.—Branches shorter, generally obtuse: leaves broader, shorter acuminate. Bot. Gaz. 19: 239. 1894.—District of Columbia.

504. *Raphidostegium sub-demissum* Kindb.—Differs from *R. demissum* in branches cuspidate, sub-julaceous: leaves smaller, long subulate, not or indistinctly recurved on borders: inflorescence dioicous. Mac. Cat. 208.—On rocks: Alaska.

505. *Raphidostegium micans submersum* Ren. & Card.—More robust: stems very much elongated, pinnate, intricate: leaves remote; perichaetial leaves longer. Revue Bryol. 20: 21. 1893.—Louisiana.

506. *Raphidostegium sub-adenatum* C. M. & Kindb.—Tufts green, intricate: branches short, sub-julaceous: leaves close, incurved at apex, ovate oblong, denticulate to middle, reflexed at margins below; inner cells sub-oblong, alar and marginal quadrate not vesiculose nor yellow; costa short, double; perichaetial leaves larger, longer, appressed, a little longer acuminate: capsule cylindric, curved, very much narrower than rostrate lid; pedicel arcuate or flexuous: probably monoicous. Mac. Cat. 209.—On trees in woods: Ontario; Quebec.

507. *Raphidostegium Kegelianum Floridanum* Ren. & Card.—Scarcely distinct from the S. American type by the shorter and broader capsule, rounded or less attenuate below. Bot. Gaz. 15: 61. 1890.—Trunks of palms: Florida.

508. *Raphidostegium Roellii* Ren. & Card.—Monoicous: delicate, densely caespitose, shining, yellowish green: leaves sub-homomallous or complanate, oblong lanceolate, decurrent, acuminate; margin plane or reflexed, denticulate above; costa double or very faint; cells elongated, narrowly linear, alar strongly dilated and hyaline or yellowish; perichaetial leaves long-

acuminate, above coarsely and irregularly dentate, costa divided or obsolete: capsule sub-erect, oblong, sub-symmetric; lid not known; teeth long subulate, densely articulate; cilia more or less elongated, nodulose. Bot. Centrbl. 44: 423. 1890. On trees: Washington.

509. *Thamnum alopecurum* Sch.—Plants robust: primary stems stoloniform, radicans, stout, with erect or inclined stems 8-12 cm. long; stems simple up to a certain height and distantly foliate, then branching; branches close, sub-distichous, having a dendroid aspect: tufts large and lax, dark green: branch leaves erect-spreading, loosely imbricate, oval oblong, acute, coarsely dentate above; costa strong, ridged dorsally, vanishing near apex; cells rounded, oval or oblong, a little longer at base: dioicous: pedicel smooth, arcuate at summit: capsule oblique or sub-horizontal, oval or oblong, contracted below orifice; lid conic, long rostrate. Husnot, Musc. Gall. 347.—In damp places along rocky cliffs: British Columbia.

510. *Thamnum Leibergii* Britton.—Dioicous: perichaetial leaves, ecostate, with recurved apices, entire or slightly serrulate; leaves costate to just below apex, entire or slightly serrulate below, coarsely serrate above: pedicel 1 cm. long, falling off with capsules when old; inner peristome, with three appendiculate regular cilia as long as the teeth or occasionally irregularly united into one or two and scarcely appendiculate. Bull. Torr. Bot. Club 16: 211. 1889.—On quartzite ledges, Idaho.

511. *Thamnum Holzingeri* Ren. & Card.—Green, slender, rather resembling in habit small forms of *Isoetium myosuroides*: primary stems creeping, stoloniform, secondary ascending or decumbent, more or less shrub-like, pinnate; branches complanate, generally attenuate: lower stem leaves small, erect spreading from a broadly deltoid base, ligulate-obtuse; costa vanishing about base of acumen; upper leaves larger, distichous, complanate, slightly asymmetric at base, oblong ligulate, obtuse or sub-obtuse; costa vanishing far from apex, sometimes forking above; branch leaves smaller, with the costa shorter and the lower margin inflexed: upper leaves acute; all the leaves plane and crenulate-serrulate on the margins, coarsely and irregularly dentate at the apex; cells parenchymatous, incrassate, short, chlorophyllose, roundish or sub-hexagonal above, ovate or oblong in the middle, sub-linear below, alar small rather obscure sub-quadrate or roundish; inner perichaetial leaves sub-vaginant, oblong lanceolate, long loriform-acuminate, serrulate; costa thin; cells narrower: pedicel smooth; capsule erect, oblong, sub-symmetric, constricted under orifice when dry; lid conic, obliquely beaked; teeth yellowish, lanceolate-acuminate, subulate; segments narrowly split along divisural line, cilia 2 long nodose. Bot. Gaz. 19: 239. 1894.—Oregon.

512. *Plagiothecium denticulatum microcarpum* Ren. & Card.—Capsule very short and turgid, scarcely 1.5 mm. long; pedicel thick, flexuous, often geniculate at base. Bot. Gaz. 14: 98. 1889.—Idaho; Washington.

513. *Plagiothecium denticulatum squarrosus* Kindb.—Distinctly squarrose when dry. Bull. Torr. Bot. Club 17: 279.—British Columbia; Behring Sea.

514. *Plagiothecium membranosum* Kindb.—Tufts dense, green, glossy: leaves distichous, crowded, patent, flat, ovate-oblong, acute or short acuminate, entire, or denticulate above middle, decurrent; cells very long and narrow, alar large, hyaline and sub-quadrate; costa none or obsolete: capsule cylindrical-obovate, horizontally curved; teeth yellow; pedicel smooth; lid unknown: probably dioicous. Mac. Cat. 215.—On dead wood: Ottawa and Belleville, Ont.

515. *Plagiothecium brevipungens* Kindb.—Tufts dense, dark green: stems prostrate, irregularly pinnate; branches attenuate: leaves crowded, scarcely decurrent; ovate-oblong, acute or short pointed, auricled, not plicate or reflexed on borders, entire or slightly denticulate at apex; upper cells long and narrow, alar very distinct quadrate inflated and hyaline; costa very short, thick and simple, or none: capsule curved, lid short, conical; pedicel smooth: monoicous. Mac. Cat. 215.—On stones: Ottawa, Ont.

516. *Plagiothecium aciculari-pungens* C. M. & Kindb.—Dioicous: tufts dense, radiculose, depressed, glossy green: stem irregularly divided; branches few and short, not attenuate: leaves concave, open erect or patent, decurrent, entire, ovate-oblong with a short needle-shaped recurved or patent point; margins narrowly recurved below middle at one side; areolation uniform and dilated; angular cells large and not numerous, sub-rectangular; costa none or indistinct: capsules not found. Mac. Cat. 216.—On earth: New Brunswick.

517. *Plagiothecium decursivifolium* Kindb.—Intermediate between *P. latebricola* and *P. pseudo-latebricola*: agrees with the last in branches complanate, leaves distichous, capsule oblique: differs in leaves broader, short-pointed, decurrent, alar cells not distinct but decurrent ones rectangular. Capsule arcuate when dry, and finally furrowed; lid longer, often curved. Mac. Cat. 277.—On cedar stumps in a swamp: Belleville, Ont.

518. *Plagiothecium Silesiacum* Sch.—Plants more robust than *P. striatellum*: stems inclined, radicans, often fasciculately branched; branches arcuate, procumbent: tufts lax, depressed, pale or yellowish green, shining: branch leaves erect-spreading, more or less secund, oval-lanceolate or lanceolate, gradually long acuminate, not decurrent, dentate in upper half; costa double, short, very faint; cells of the basal angles rectangular or oblong, middle linear; inner perichaetial leaves sheathing, terminated by a long, dentate, recurved acumen: monoicous: capsule oblique or horizontal, cylindrical, smooth, collum distinct, slightly contracted below mouth; lid conic; annulus simple; teeth pale, inner membrane half height of teeth. Husnot, Musc. Gall. 354.—Newfoundland; New Brunswick.

519. *Plagiothecium pseudo-latebricola* Kindb.—Tufts dense, radiclese, glossy green: stem irregularly branching; branches often curved, perfectly complanate, branchlets longer with smaller and narrower leaves: stem leaves small, distichous, not decurrent, shortly ovate lanceolate, suddenly tapering to a somewhat long filiform and straight point, entire; cells narrow, alar small few and sub-quadrate often wanting; costa obsolete; perichaetial leaves sinuate above: capsule obovate-oblong, oblique and faintly curved, often pendent; teeth pale; inner membrane clavate, cilia long, sub-appendiculate; lid conical: dioicous. Mac. Cat. 211.—On rotten wood: Columbia River, B. C.

520. *Plagiothecium bifariellum* Kindb.—Plants small, sparingly radiclese, loosely caespitose, dark or blackish green, not glossy: stems rigid, pinnate: leaves loose, small, spreading, smooth and not striate; stem leaves at base broadly ovate-cordate, decurrent, serrulate all around, abruptly attenuate to filiform hooked-deflexed and sub-entire acumen; cells narrow, linear, basal oblong; costa obsolete or reaching to middle: dioicous. Bull. Torr. Bot. Club 17: 279.—Wet places in woods: Vancouver Island.

521. *Plagiothecium attenuatiformum* Kindb.—Tufts green, faintly shining, loose, with few rhizoids: primary stem very short; branches elongate, long attenuate, finally flagelliform: leaves sub-distichous, lower broadly ovate, obtuse or obtusate, entire, long decurrent, concave, recurved at borders from base to above middle at least at one side; cells chlorophyllose, somewhat dilated, lowest very much wider and shorter and nearly uniform; costa generally short and double, rarely simple and reaching to middle; other leaves gradually smaller, narrower and more acute or acuminate: barren. Mac. Cat. 277.—On rocks: Quebec.

522. *Amblystegium fenestratum* Kindb.—Plants loosely coherent, green: stems capillary, irregularly ramulose: leaves small, spreading, very narrow, ovate lanceolate, acute, denticulate; cells dilated but elongate; costa more or less distinct: barren. Habit of *A. Sprucei*. Mac. Cat. 217.—On borders of a pond near London, Ont.

523. *Amblystegium spelrophyllum* Kindb.—Plants loosely caespitose, dark green: stem capillary, irregularly ramulose, not or sparingly radiclese: leaves small, long-distant, spreading, sub-cordate or oval oblong, blunt or sub-acute, entire or denticulate; cells short; costa sub-percurrent, broad, sometimes very distinct: barren: probably dioicous. Mac. Cat. 217.—On rocks: Canaan's Fork, N. B.

524. *Amblystegium serpens xanthodictyon* Kindb.—Tufts loose, yellow-brown: stem pinnate: leaves from an ovate-oblong base acuminate, denticulate or entire; alar and often also lower basal cells quadrate, yellowish, others oblong except inner, sublinear near costa and in acumen; perichaetial leaves very small, nerveless. Mac. Cat. 218.—On stones: St. Mary's river, Canada.

525. *Amblystegium Juratzkanum* Sch.—Closely related to *A. serpens*, but generally more robust: tufts dark olive green: leaves spreading; stem leaves distant, cordate-oval, short acuminate; branch leaves oval-lanceolate, longer acuminate, entire or denticulate; costa longer, reaching $\frac{3}{4}$ length of leaf, cells larger, sometimes smaller, very chlorophyllose, basal rectangular, middle sub-rhomboidal; perichaetial leaves less abruptly acuminate, costate and plicate: monoicous. Husnot, *Musc. Gall.* 358. According to Husnot this species differs from *A. hygrophilum* in dark green color, leaves longer acuminate, longer costate and cells larger.—On dead wood, stones, and bases of trees in damp woods: Ottawa and Owen Sound, Ont.; Revelstoke, B. C.

526. *Amblystegium hygrophilum* Sch.—Monoicous: stems 3-5 cm., depressed, delicate; branches erect or ascending: tufts pale or yellowish green: leaves spreading in all directions, or squarrose, small stem leaves cordate-oval, branch leaves oval, long acuminate, entire; costa weak, vanishing near middle: cells of basal angles rectangular, forming small auricles, middle cells linear-rhomboidal; internal perichaetial leaves oblong-lanceolate, long and finely acuminate, costate, entire: capsule sub-horizontal, oblong-cylindric, strongly arcuate when empty, contracted below mouth; lid conic; annulus very large. Husnot, *Musc. Gall.* 358.—Indiana; Illinois; Wisconsin.

527. *Amblystegium porphyrrhizon* Lindb.—Monoicous: slender, in habit between small forms of *A. riparium* and *A. serpens*; stem leaves rather remote, sub-squarrose-spreading, from deltoid-ovate to lanceolate ovate, slender, pale, excavate at the slightly decurrent angles; costa slender, yellowish, vanishing beyond middle; margin below obsoletely serrulate; cells narrowly oval-hexagonal, hyaline, quadrate loose and yellowish at angles: male fls. numerous, small, polyphyllous; leaves imbricate, short lanceolate, erect, from an ovate concave base; antheridia few; female fls. slender, sub-incurved: perichaetial branch strongly radicate; inner leaves erect, imbricate, long lanceolate, subulate-acuminate; costa slender, yellowish, vanishing toward base and apex; margin faintly serrulate; cells very narrow, vermicular: capsule oblique, oblong-cylindric, cylindric when dry and deoperculate, sub-arcuate, constricted below mouth, yellow-fuscescent, thin walled, loosely and irregularly areolate; annulus? Schimp. *Syn.* 715. 1876.—Kansas; Canada: N. W. Terr.; Rocky Mts.; British Columbia, Miquelon Is.

528. *Amblystegium Schlotthaueri* Ren. & Card.—Sub-species of *A. serpens*: similar in habit to compact form of *A. serpens* but differs in pedicel thicker, strongly twisted to right; capsule erect, slightly curved, ovoid, mouth dilated, collum elongate, plicate; cilia of endostome shorter, solitary. *Bot. Centralbl.* 44: 423. 1890.—Yellowstone Park, Wyoming.

ly related to *A. ser-*
 een: leaves spreading;
 ; branch leaves oval-
 costa longer, reaching
 y chlorophyllose, basal
 leaves less abruptly
 ot, Musc. Gall. 358.
pygrophilum in dark
 and cells larger.—On
 ls: Ottawa and Owen

ous: stems 3-5 cm., de-
 fts pale or yellowish
 ee, small stem leaves
 tire; costa weak, van-
 gular, forming small
 chaetial leaves oblong-
 : capsule sub-horizon-
 ty, contracted below
 asc. Gall. 358.—Indi-

onoicous: slender, in
serpens; stem leaves
 id-ovate to lanceolate
 ent angles; costa slen-
 below obsolete serru-
 e loose and yellowish
 eaves imbricate, short
 ridia few; female fls.
 radicant; inner leaves
 costa slender, yellow-
 rrrulate; cells very nar-
 ylindric when dry and
 th, yellow-fuscescent,
 ulus? Schimp. Syn.
 s.; British Columbia,

d.—Sub-species of *A.*
serpens but differs in
 rect, slightly curved,
 of endostome shorter,
 e Park, Wyoming.

529. *Amblystegium distantifolium* Kindb.—Stems irregularly branch-
 ing, creeping; leaves green, distant, spreading when dry, patent when
 moist, long oval lanceolate, long cuspidate, not or indistinctly decurrent,
 sharply serrulate all around, chlorophyllose; cells wide, sub-oblong; costa
 thick, percurrent or nearly excurrent: barren. Mac. Cat. 222.—On rocks,
 Newfoundland.

530. *Amblystegium dissitifolium* Kindb.—Tufts compact, sparingly
 radicant, green, not glossy: leaves very loosely disposed, long decurrent,
 ovate oblong, short-acuminate, denticulate all around; areolation loose,
 upper cells narrower, alar larger quadrate and well distinct, as chlorophyl-
 lose as the other cells; costa percurrent: probably dioicous; only male
 flowers found. Mac. Cat. 220.—On flat limestone rocks which receive
 dripping water: Canada.

531. *Amblystegium sub-compactum* C. M. & Kindb.—Differs from
A. compactum in stems thicker, leaves larger and longer; capsule asym-
 metric, curved in young state, at least doubly greater. The British Col-
 umbia specimens have a peculiar habit; tufts are decolorate below, bright
 green above, and stems erect. Mac. Cat. 221.—Growing in thick tufts at
 the bases of trees around springs and margins of bogs, also on wet
 rocks: British Columbia and Canada.

532. *Amblystegium riparium longifolium* Sch.—Leaves narrow, lanc-
 eolate, long and finely acuminate. Husnot, Musc. Gall. 363.—Vancouver
 Island and Washington.

533. *Amblystegium riparium serratum* Ren. & Card.—Plants slen-
 der, creeping; leaves narrow, serrulate at apex. Bot. Gaz. 14: 98. 1889.—
 Roots of trees: Kansas.

534. *Amblystegium Floridanum* Ren. & Card.—Very small, ap-
 pressed: leaves small, narrowly lanceolate, long acuminate, entire: cap-
 sules short, arcuate. Bot. Gaz. 14: 98. 1889, as *A. riparium*, var.—
 Florida; Louisiana.

535. *Amblystegium Kochii* Sch.—Closely related to *A. riparium*:
 stems depressed, with ascending or erect branches: leaves spreading in all
 directions; those of large branches cordate-oval or broad oval; those of
 small branches long acuminate, denticulate, costate for $\frac{3}{4}$ length, basal
 cells rectangular; middle sub-hexagonal, much larger than in *A. riparium*;
 pedicel long; capsule oval or oblong; annulus simple. Husnot, Musc.
 Gall. 362.—Kansas.

536. *Amblystegium homalostegium* Jgr. & Sauerb.—Monoicous: tufts
 low, broad, dirty green, rather dense and rigid: stem creeping, pseudo-pin-
 nate, with many short erect or curved densely aggregate slender simple
 branches: stem leaves densely imbricate, indistinctly secund, when wet
 making the stems appear julaceous and turgescant, broadly ovate from a
 cordate base, short acuminate, concave; margin plane, obsolete denticu-

late; costa double, short; cells minute, indistinct, narrowly elliptic, pale, slightly papillose, alar quadrate; perichaetial leaves lanceolate-acuminate, longer, secund, concave, longitudinally plicate, ecostate; yellowish; seta yellowish-red, ascending; capsule horizontal, minute, asymmetric-cylindric, strongly constricted in middle, gibbous or strumose at base, wide-mouthed, brown; lid minute, short apiculate; peristome teeth very prominent, connivent when wet, red, strongly cristate, segments yellow, smooth, broad, carinate, imperforate, cilia solitary, broad, shorter and paler. *Hypnum homalostegium* Müller, Flora 56: 484. 1873.—Trunks of trees, West Fowl River, Alabama.

537. *Hypnum Sommerfeltii* Myr.—Stems 15-30 mm. long, delicate, depressed, irregularly divided or sub-pinnate, branches ascending; tufts green or yellowish; leaves crowded, very spreading, sometimes sub-secund at extremity of branches; stem leaves broadly oval-lanceolate, long acuminate, denticulate below, ecostate or faintly bicostate; branch leaves oval lanceolate; basal cells quadrate or rectangular, forming yellowish auricles, others linear, broader than in *H. Halleri*; inner perichaetial leaves oblong, plicate; capsule sub-horizontal, oblong sub-cylindric, arcuate, contracted below mouth; lid convex conic; annulus large; monoicous. Husnot, Musc. Gall. 304.—On old logs, bases of trees, damp rocks and earth: Canada; British Columbia.

538. *Hypnum byssiframmeum* C. M. & Kindb.—Resembling a small form of *H. Sommerfeltii*, but leaves denticulate all around: barren. Mac. Cat. 323.—On the base of a dead tree, Ontario.

539. *Hypnum Macounii* Kindb.—Allied to *H. hespidulum*: habit of the European *H. Halleri*: monoicous: plants small, interlaced in dense brownish green tufts: stems pinnate, prostrate, sparingly radiculose: leaves densely crowded, squarrose-recurved, round deltoid, short acuminate, reflexed on borders of entire base to denticulate acumen; alar cells quadrate, pellucid, not numerous, others short oblong or short lanceolate; costa obsolete or none: capsule small, narrow cylindric, slightly curved; annulus simple; basilar membrane low; lid convex, obtuse, not apiculate; pedicel reddish brown, smooth. Mac. Cat. 224.—On earth and rocks: British Columbia; Rocky Mountains.

540. *Hypnum uncostatum* C. M. & Kindb.—Differs from *H. chrysophyllum* in dense tufts: stems more irregularly branching, creeping: leaves shorter-acuminate; alar cells smaller, not yellow; costa more distinct; perichaetial leaves gradually acuminate-subulate or filiform pointed with acumen arcuate: capsule smaller. Mac. Cat. 224.—Canada.

541. *Hypnum decursivalum* C. M. & Kindb.—Differs from *H. chrysophyllum* in leaves decurrent with broader base, borders recurved at angles; alar cells numerous, hyaline: barren. Mac. Cat. 224.—On old logs: British Columbia; Newfoundland.

542. *Hypnum Columbianae* Kindb.—Tufts dense: stems short, very tomentose, irregularly branching; branches short: leaves narrower than in *H. chrysophyllum*, from narrow ovate base gradually tapering into acumen; borders denticulate all around, recurved at angles; areolation often wide as in *Amblystegium*; costa reaching to acumen or sometimes longer; perichaetial leaves short-acuminate: capsule generally smaller and shorter than in *H. chrysophyllum*, curved; lid short-apiculate; peristome dark-yellow, not pale. Mac. Cat. 224.—On wet logs: British Columbia.

543. *Hypnum stellatum sub-decurvatum* Kindb.—Leaves smaller, decurrent, abruptly acuminate from a short ovate base; alar cells more numerous: not found fruiting. Mac. Cat. 225.—Growing with *Dicranum scoparium* on earth in a swamp, Ontario.

544. *Hypnum polygamum longiaerve* R. & C.—Leaves narrower, long acuminate; costa stronger, extending into acumen: capsule paler, narrower. Bot. Centralbl. 44: 423. 1890.—Victoria; Vancouver Island.

545. *Hypnum polygamum fallaciosum* Lindb.—Larger, often strikingly like *H. aduncum Kneiffi* in habit: costa very variable, forked, longer or shorter, or wanting. Milde, Bryol. Siles. 346. 1869.

546. *Hypnum aduncum pungens* H. Müll.—Leaves erect, sub-imbriate, apical ones inrolled at the point, short acuminate or subapiculate, straight or a little curved. Renault in Husn. Muscol. Gall. 387. 1894. Yellowstone Park.

547. *Hypnum aduncum attenuatum* Boul.—Slender, sometimes elongated: stem leaves short, deltoid, curved at point; costa with tendency to bifurcate; branch leaves small, without auricles. Ren. l. c.—Deer Lodge, Mont.

548. *Hypnum aduncum platyphyllum* Kindb.—Leaves very broad and short acuminate. Mac. Cat. 226.—On rocks in woods: Rockcliff, near Ottawa, Can.

549. *Hypnum aduncum RoeIII* Ren.—Leaves distant, spreading, flexuous, apical oblong, then gradually long subulate acuminate, acumen flexuous and twisted. Ren. *ibid.* 372.—Yellowstone Park.

550. *Hypnum aduncum flexile* Ren.—Emergent, more slender: stems flexuous: leaves usually narrower, flexuous or slightly homotropous prolonged into a twisted subula. Ren. *ibid.* 373.—Vancouver; Hobart, Ind.

551. *Hypnum capillifolium* Warnst.—Dioicous: stem erect (5-15 cm.), pinnate: leaves secund or falciform, oblong lanceolate, gradually narrowed from the base, long subulate by the excurrent costa which is strong, 90-140 μ wide at base, much thicker than the lamina; tissue delicate, basal cells distinctly and long excurrent, alar dilated, forming large convex auricles reaching almost to the costa. Ren. *ibid.* 379.—Idaho; Washington; Vancouver Is.; British Columbia.

552. *Hypnum symmetricum* R. & C.—Sub-species of *H. uncinatum*: leaves moderately striate, capsule narrow, cylindrical, always exactly erect, symmetric; seta sometimes in pairs. Ren. *ibid.* 379.—Rocky Mountain and Pacific Coast region.

553. *Hypnum fluitans Jeanbernati* Ren.—Monoicous: tufts pale green, sometimes rather dense, 6-10 cm. high: leaves feebly homotropous except at summit, oblong or oblong-lanceolate, narrowed into an often short and rather broad acumen, truncate at base, sinuate or slightly denticulate, sometimes dentate; costa slender (47μ), little surpassing the middle; median cells very long, alar little dilated, not vesicular, poorly delimited: seta 4-6 cm.—Ren. *ibid.* 381.

554. *Hypnum fluitans Delamarei* R. & C.—Monoicous: plants tall, slender (25 cm.): tufts floating, pale green at surface, dark brown below; some stems regularly pinnate: stem leaves falciform, strikingly recurved on some branches, on others simply secund, narrowly lanceolate, slenderly acuminate or short subulate, plainly dentate at summit; branch leaves linear, spreading, flexuous; costa colored, narrow ($40-60\mu$), little surpassing the middle: median cells very long and narrow, remainder elongated and compact to the base, basal with walls a little thickened, alar small, occupying almost all the base, not forming distinct auricles: capsule short, erect, blackish; seta 4-5 cm., surpassing the stem. Ren. *ibid.* 384. Miquelon Is.

555. *Hypnum fluitans pinatum* Boul.—Tufts yellowish green: stem rather short (8-10 cm.), erect, stout, usually pinnate: leaves falciform, oval or oval-oblong or narrower at base, slenderly acuminate or short subulate, usually denticulate at base and apex; costa reaching middle of point; auricles large, swollen, rounded composed of rather numerous cells with generally unthickened walls. Ren. *l. c.*—Miquelon Is.

556. *Hypnum fluitans falcifolium* Ren.—Usually purplish or mixed with green or brown: stem 5-10 cm or longer, pinnate: leaves falciform, rather distant, lanceolate narrowed into a long subula often spirally twisted and sparingly toothed costa broad ($80-90\mu$) at base, ending in the subula or reaching the point but not distinctly excurrent; median cells long and narrow; auricles composed of cells usually thickened and colored. Renauld, *ibid.* 387.—Yellowstone Park.

557. *Hypnum fluitans hemineuron* R. & C.—Tufts pale green, depressed, intricate, small: stem prostrate, ascending at tip, slender, very short (1-3 cm.), vaguely branched, radicles distinct: leaves slightly homotropous, oblong-lanceolate, short decurrent, acumen moderate, with some teeth or sub-entire; costa very slender ($33-48\mu$), sometimes simple to beyond the middle, more often bifurcate in stem leaves, more feeble and shorter and sometimes almost wanting in branch leaves; median cells loose, unequal as to length and breadth, basal almost equal, also a little broader,

es of *H. uncinatum*:
always exactly erect,
9.—Rocky Mountain

ous: tufts pale green,
homotropous except
to an often short and
slightly denticulate,
ng the middle; median
ly delimited: seta 4-6

onoicous: plants tall,
dark brown; thin;
strikingly recurved on
lanceolate, slenderly
summit; branch leaves
(60 μ), little surpassing
lander elongated and
ned, alar small, occu-
aricles: capsule short,
m. Ren. *ibid.* 384.

yellowish green: stem
leaves falciform, oval
ate or short subulate,
middle of point; auri-
erous cells with gen-
ly purplish or mixed
ate: leaves falciform,
subula often spirally
t base, ending in the
current; median cells
thickened and colored.

tufts pale green, de-
at tip, slender, very
leaves slightly homo-
moderate, with some
etimes simple to be-
ees, more feeble and
es; median cells loose,
also a little broader,

not forming distinctly delimited auricles. Ren. *ibid.* 388.—Packs Harbor, Labrador.

558. *Hypnum fluitans conflatum* C. M. & Kindb.—Stem slender, sub-filiform, distinctly pinnate, not radiculose: leaves small, concave, distant, denticulate all around; stem leaves decurrent, from a broad ovate base suddenly narrowed into a very short, subulate-filiform straight point; alar cells very large, hyaline or faintly yellowish, others nearly uniform, oblong-lanceolate, conflate; costa pale yellow, vanishing in the acumen; branch leaves narrower, oblong-lanceolate, more or less short acuminate, curved or straight: capsule very small, arcuate, contracted below mouth: dioicous. Mac. Cat. 230.—In pools and bogs: British Columbia; Ottawa, Ont.; Labrador.

559. *Hypnum Moseri* Kindb.—Differs from *H. uncinatum* in leaves not striate, but sometimes recurved at base; costa faint, often falling; differs from all other *Harpidia* in stem densely radiculose. Mac. Cat. 229.—On bases and trunks of poplar trees: New Brunswick; Newfoundland.

560. *Hypnum filicinum aculeatum* C. M. & Kindb.—Costa excurrent to a rigid point. Mac. Cat. 231.—On wet rocks: British Columbia.

561. *Hypnum decipiens* (DeNot.) Kindb.—Monoicous: habit of *Hypnum commutatum* or *flicinum*; loosely interwoven, with green innovations, scarcely shining; stems 4 cm., paraphyllia present, pinnate; branches simple, slender, spreading; stem leaves sub-squarrose, broadly cordate-deltoid, short cuspidate, decurrent, plicatulate; margin reflexed below, denticulate all around especially at apex; costa strong, vanishing below apex; branch leaves small, ovate-acuminate, secund falcate; costa reaching beyond middle; cells short, sub-rhombic to oblong, strikingly papillose especially on lower face, basal on decurrent, elongate, hyaline: perichaetial leaves membranaceous, pallescent, appressed, inner slenderly subulate, fimbriate: capsule on a long seta, thick-clavate from an erect base, cernuous, brown. *Thuidium decipiens* DeNot. Epil. 233. 1889.—On rocks: Rocky Mts.; Br. Columbia; Vancouver Is.

562. *Hypnum chloropterum* C. M. & Kindb.—Tufts laxly cohering, with few rhizoids, whitish or bright green, not shining: stem more or less densely pinnate, rigid: stem leaves small, patent also in dry state, loosely and long-decurrent, auricled, papillose at back, distinctly but faintly plicate, concave, broad-ovate, suddenly narrowed to a short (in the dry state twisted) acumen; borders broadly recurved below, sometimes to middle, denticulate all around from apex to auricles; areolation lax, more or less chlorophyllose, alar cells larger, sub-quadrata, numerous, others oval oblong, only the uppermost narrow; costa distinct, reaching above middle;

branch leaves more loosely disposed, sometimes ovate-oblong and narrow areolate; inner perichaetial leaves narrower, longer filiform pointed, nerveless; capsule small, sub-oblong, straight or curved; lid elongate-conic, oblique-apiculate, or rostellate; pedicel very rough, purple; monoicous. Mac. Cat. 231.—On rocks and on ground: Newfoundland, New Brunswick.

563. *Hypnum pseudo-fastigiatum* C. M. & Kindb.—Allied to *H. reptile*, but alar leaf cells more numerous, chlorophyllose and dusky, not decolorate; perichaetial leaves nerveless; capsule scarcely constricted below mouth. Mac. Cat. 235.—On bases of vines in woods, British Columbia. On rocks, Ontario.

564. *Hypnum fastigiatum* Brid.—Stems delicate, creeping, radiculose, much divided, erect in middle of tufts, spreading around outside, short arcuate at summit; paraphyllia quite numerous, lanceolate or digitate: tufts very large, depressed, yellowish green near surface, brownish within: leaves falciform-secund, stem leaves oval-lanceolate, branch leaves a little narrower, gradually narrowed into a long, sharp point, entire or superficially denticulate, plain on borders or slightly revolute, hyaline at base; costa bifurcate, narrow and very short; quadrate cells of basal angles quite numerous, middle ones linear, attenuate; inner perichaetial leaves half sheathing, acuminate, plicate, faintly bicostate: monoicous or dioicous: capsule erect or oblique, oblong-cylindric, arcuate, contracted below mouth; lid convex, apiculate, sometimes short rostrate; annulus narrow. Husnot, Musc. Gall. 400.—On dry rocks: Rocky Mountains; Greenland.

565. *Hypnum Waghornii* Kindb.—Differs from *H. fertile* in large and hyaline alar leaf cells, other basal ones not yellow: capsule tumid at base, slightly curved: paraphyllia very broad. Mac. Cat. 234.—Newfoundland.

566. *Hypnum revolutum* Mitt.—Stems 2-8 cm., ascending-erect, irregularly branched when tufts are compact, depressed and pinnate when loose; yellowish green, ferruginous within: leaves falciform-secund, oval or oblong-lanceolate, long acuminate, denticulate at summit, plicate when dry, strongly revolute from base to apex; costa none or double and short; cells of basal angles roundish quadrate or rectangular, rather numerous, forming small auricles, median cells linear, relatively short (1:6-8); perichaetial leaves strongly plicate: capsule sub-horizontal, rather large, oblong, arcuate, strongly contracted below mouth. Husnot, Muscol. Gall. 402. 1894.—Morley and Hector, Rocky Mts.; Greenland.

567. *Hypnum revolutum Villardi* Ren. & Card.—Leaves short acuminate, not or scarcely plicate, alar cells numerous, large. *H. Heufleri Villardi* R. & C., Bot. Centralbl. 44: 423. 1890.—Montana.

568. *Hypnum Canadense* Kindb.—Intermediate between *H. imponens* and *H. sub-imponens*: dioicous: densely caespitose, yellow or pale green:

stems creeping, densely pinnate ramulose; branches robust, thick and tumid; leaves close, falcate, with elongate-ovate or oblong base and short acumen, faintly or not striate, more or less denticulate all around, not reflexed on borders, larger than in *H. imponens*; cells very narrow, alar larger and pellucid, other basal cells yellow; paraphyllia few, subulate; perigonial leaves very broad-ovate, abruptly narrowed to a straight subulate point; capsule obovate, asymmetric or sub-cylindric and arcuate, thick and not striate; teeth yellow; segments orange, cilia short and not appendiculate. Mac. Cat. 236.—Newfoundland; Alaska; on stones: Nova Scotia and Quebec; on rotten logs: Rocky Mountains and Ontario.

569. *Hypnum cressiforme* Pyreniacum Ren.—Closely related to the variety *filiforme*, from which it is distinguished by the short acuminate leaves, quite strongly dentate. Fl. Miq. 55.—Miquelon Island.

570. *Hypnum Vaucherii* Lesq.—Plants resembling certain forms of *H. cressiforme*: stems erect-fastigiate: tufts compact, dark green or yellowish: leaves crowded and imbricate, more or less falciform-secund, sometimes erect so as to give branches a sub-julaceous appearance, oval or oval-lanceolate, entire or sinuate, plane on borders; costa very short, simple or bifurcated, one branch longer than the other; cells of basal angles more numerous and smaller than in *H. cressiforme*, walls thickened, middle cells broader and shorter, 6-8 times as long as broad: fruit unknown. Husnot, Musc. Gall. 406.—Montana.

571. *Hypnum Renaudii* Kindb.—Agrees with *H. curvifolium* in stem more or less pinnate, inner basal leaf cells finally yellow: with *H. Lindbergii* in leaves decurrent, alar cells very much dilated, capsule not plicate when dry: differs from both in entire leaves. *H. pratense* differs in leaves not striate nor decurrent, and alar cells not evolute. Mac. Cat. 238.—On earth and old logs and sometimes on rocks: British Columbia; Canada; Newfoundland.

572. *Hypnum Patientiæ* Lindb.—Closely related to *H. pratense*: differs from it by stems and branches not complanate, curved at summit: all leaves falciform-secund, broad oval-lanceolate, with a larger, entire acumen; costa none or double and very short; cells of basal angles large, forming hyaline auricles; middle ones linear, attenuate: capsu dric, arcuate, rarely fruiting. Husnot, Musc. Gall. 406.—On rocks: New Brunswick; Greenland; Miquelon Island; Newfoundland; Pennsylvania; Indiana; Wisconsin; Montana.

573. *Hypnum Patientiæ elatum* Sch.—Extensively cespitose, tufts yellowish or faintly rufescent: stems 2-3 inches long, erect, sub-fastigiate branched: leaves falcate and sub-hamate, narrower, long acuminate. Sch. Syn. 758. 1876.—Miquelon Island.

574. *Hypnum Patientiæ demissum* Sch.—Tufts deplanate, pale or

bright green, dark variegated: stems long, creeping, more or less regularly pinnate. l. c.—Miquelon Island.

575. *Hypnum Patientiæ Americanum* Ren. & Card.—Stems slender, prostrate, more or less distinctly pinnate: leaves smaller, with acumen shorter and broader. Bot. Gaz. 14: 99. 1889.—On decayed wood and sandy ground: Louisiana.

576. *Hypnum arcuatiforme* Kindb.—Tufts dense, green, not glossy: stem creeping, sub-pinnate: leaves arcuate, ovate lanceolate, generally short-acuminate or sub-obtuse, entire, decurrent, not striate; alar cells large, well defined, orange, others pale and narrow; costa none or short and double: capsule sub-cylindrical, curved, not striate nor furrowed, constricted below the wide mouth, teeth when dry incurved, pale yellow, hyaline margined; cilia long, appendiculate: probably dioicous. Mac. Cat. 238.—On earth near Ottawa, Ont.

577. *Hypnum Dieckii* Ren. & Card.—Tufts yellowish or rufescent; stem depressed, pinnate; branches ascending: leaves falcate-secund, strongly circinate, from broadly ovate base suddenly acuminate, acute or subulate; margin plane, entire, rarely obsolete denticulate above; costa double, short, sometimes very faint or prolonged to middle; cells linear, very narrow, obtuse, alar large, strongly inflated, hyaline or flavescent; perichastial leaves oblong-lanceolate, long acuminate, sub-entire, ecostate: dioicous: capsule horizontal or sub-pendulous, large, arcuate; lid unknown; teeth yellowish, strongly trabeculate, segments narrow, scarcely perforated along middle; cilia 2, nodulose. Bot. Centralbl. 44: 423. 1890. Oregon.

578. *Hypnum pseudo-pratense* Kind.—Nearly allied to *H. pratense*: tufts more compact; branches radiculose below: leaves more crowded, not decurrent, more distinctly denticulate near apex: inflorescence monoicous: capsules not found. Mac. Cat. 239.—On old logs in woods: Ontario.

579. *Hypnum Haldanianum* Roellii Ren. & Card.—Branches short, interwoven, leaves short and broad acuminate, areolation dense. Bot. Centralbl. 44: 424. 1890.—Tree trunks: Indiana.

580. *Hypnum flaccum* C. M. & Kindb.—Tufts large and loose, brown below, pale green above: stems pinnate, sparingly radiculose, red-brown; branches elongate, attenuate, distant and flaccid; paraphyllia few, multi-form: leaves oblong-lanceolate, shortly subulate-acuminate, not curved, concave-involute, patent, with excavate dark orange auricles at base; stem leaves decurrent, slightly recurved at basal angles; branch leaves loose, sub-distichous, not decurrent nor recurved; cells narrow, long linear, not chlorophyllose, alar large, sub-quadrate, inner basal narrow, pale orange; costa short, double and indistinct or none: barren. Mac. Cat. 240.—On old logs or rocks: New Brunswick; Ontario.

581. *Hypnum subflacum* C. M. & Kindb.—Tufts loose, glossy green: stem green, irregularly branching, not radiculose; branches few and long, flaccid, sub-compressed when dry; paraphyllia none: leaves striate, oblong-lanceolate, denticulate above to $\frac{1}{4}$, not curved, loosely appressed or sub-patent; basal angles hyaline, not excavate; stem leaves short-decurrent, acute or short-acuminate; branch leaves not decurrent, longer acuminate; cells narrow, long-linear, chlorophyllose, basal dilated, hyaline and irregular, sub-rectangular, inner rarely pale yellowish; costa none: dioicous: female plants not found. Mac. Cat. 240.—On earth: Ontario.

582. *Hypnum pseudo-drepanium* C. M. & Kindb.—Tufts loose, green, faintly glossy: secondary stems very long, flaccid, sub-pinnate, sparingly radiculose, faintly compressed; paraphyllia none; branchlets few and very short, curved at apex: leaves plicate, entire, from a short broad ovate base narrowed into a short incurved acute acumen, crowded, loosely appressed when dry, not decurrent, not distinctly chlorophyllose; basal cells hyaline, dilated, thick-walled, alar large, sub-rectangular, well-defined, others longer and narrower; auricles excavate; costa indistinct or short and double: dioicous; female plants not found. Mac. Cat. 240.—On old logs in woods: Ottawa, Ontario.

583. *Hypnum circulfolium* C. M. & Kindb.—Nearly allied to *H. dilatatum* Wils.: stem loosely foliate, denudate at the base; leaves patent when dry, sub-circular, faintly crenulate nearly all around, slightly decurrent, at the cordate base distinctly auriculate, in the middle carinate, narrowed above to an indistinct obtuse tip; alar cells large, inflated, sub-oval, apical also short, others narrow and flexuous; costa none or indistinct: barren. Mac. Cat. 242.—On rocks: New Brunswick.

584. *Hypnum pseudo-arcticum* Kindb.—Differs from *H. arcticum* in leaves crenulate, at least above middle; costa short and double, not reaching to middle; perichaetial leaves short acuminate, serrulate: peristomial segments rimose in middle: stem sparingly radiculose. Mac. Cat. 242.—On stones in brooks: British Columbia.

585. *Hypnum Gonlardi* Sch.—Tufts thick, very soft, variegated with red and green: branches flexuous-erect from a prostrate filiform eradiculose and partly denudate stem, very slender, soft, not radicant, sub-simple or oft divided, 2-4 cm. long; leaves small, rather distant, equally spreading when moist, coherent when dry; lower branch leaves minute, circular, others ovate-rotund, apex rounded or slightly obtuse pointed, strongly concave and sub-cochleariform, margin rather broadly recurved, quite entire, narrowly decurrent at angles, not excavate; costa bifurcate, long or almost obsolete; cells at apex rhombic, toward base flexuous fusiform, sub-vermicular or hexagonal-rhombic, at base loosely rhombic-hexagonal, at angles rectangular: flowers and fruit unknown. Synop. Musc.

Eur. 778. 1876.—On stones: Ste. Anne des Monts river, Gaspé Co., Quebec; Gunn river, Anticosti.

586. *Hypnum torrentis* C. M. & Kindb.—Differ from *H. Goulardi* in stems filiform, very rigid; leaves smaller, sub-circular, crenulate all around, reflexed at basal margins; costa thick and nearly percurrent; differs also from *H. arcticum* in loosely disposed decurrent leaves with large, angular cells. Mac. Cac. 243. —On sloping limestone rocks: British Columbia.

587. *Hypnum Norvegicum* Sch.—Appearance of *H. arcticum* but much smaller and more delicate; stems depressed, branches erect or ascending; leaves erect-spreading or sub-secund, small, oval or oval sub-orbicular, sinuate, obtuse; costa very short, bifurcate; cells of basal angles quadrate or rectangular, not forming distinct auricles; middle quite short, linear-sub-hexagonal; capsule oblique, oval or oblong; lid convex, very shortly apiculate; annulus large. Husnot, Musc. Gall. 413.—Greenland.

588. *Hypnum Columbieo-palustre* C. M. & Kindb.—Differs from *H. palustre* in leaves longer apiculate, faintly denticulate; margins recurved below at one side; alar cells dilated, well-defined; costa stout, vanishing above middle. Mac. Cat. 241.—On rocks: British Columbia.

589. *Hypnum polare* Kindb.—Loosely cespitose, tufts pale green, soft; secondary stem erect, fastigately branched, flaccid, eradiculose; leaves loosely disposed, erect-spreading and slightly secund, ovate-oblong, gradually short and acute acuminate, strongly concave, margin erect, thin, at basal angles very slightly decurrent, not excavate, obsolete crenulate only at very apex; costa simple, sub-terete, vanishing below apex; cells slightly hyaline, rhomboidal, 2 to 3 times as long as broad, middle cells longer and narrower, basal cells wider, alar cells few, hexagonal-rectangular, hyaline or slightly chlorophyllose. Schimp. Syn. 780. 1876.—Greenland.

590. *Hypnum engyrium Miquelonense* R. & C.—Leaves smaller, acumen short, obtuse or sub-obtuse; cells of basal angles hardly distinct. Revue Bryol. 20: 23. 1893.—Miquelon Is.

591. *Hypnum engyrium Mackayi* Sch.—Plants more robust; leaves erect-imbricate, distinctly denticulate at summit; costa simple or double, passing middle; auricles formed of smaller cells. Husnot, Musc. Gall. 412.—Tennessee.

592. *Hypnum ochraceum flaccidum* Milde. Stems long, loosely foliate; leaves throughout spreading every way, uniform brown, broad lanceolate, long pointed; costa long, double. Bryol. Siles. 376. 1869.—Montana; Oregon; Washington.

593. *Hypnum pseudo-montanum* Kindb.—Nearly allied to *H. montanum*; differing principally in stem longer and more robust; leaves larger and thinner, more loosely disposed, longer decurrent, not distinctly dentic-

ulate; alar cells large, reaching to costa which is simple and prolonged above middle; inflorescence monocious: capsules not found. Mac. Cat. 243. — On rocks in streams: British Columbia.

594. *Hypnum purum* L.—Stems 8-15 cm., depressed or ascending, simply pinnate or with pinnate branches, delicate, julaceous; tufts large, soft, often depressed, pale green; leaves imbricate, very concave, decurrent, denticulate all around, plicate; stem leaves broad ovate, strongly contracted at base, apex round with an erect or curved apiculus; branch leaves narrower, oblong; cells of basal angles quadrate or rectangular, pale green, forming small auricles; middle cells linear flexuous; inner perichaetial leaves lanceolate, long acuminate, costa: dioicous: capsule horizontal, oblong or sub-cylindric; lid conic, pointed; annulus double; teeth orange, segments wide open along keel. Husnot, Musc. Gall. 419.—Miquelon Is.

595. *Hypnum stramineum laxifolium* C. Müll. No description of this variety is at hand.—Greenland.

596. *Hypnum stramineum compactum* Milde.—Tufts thick, stems stronger and shorter, below (with leaves) yellow-brown; leaves shorter and broader. Bryol. Siles. 370. 1869.—Greenland.

597. *Hypnum stramineum exiguum* Ren.—Stems depressed, short, very delicate, filiform; leaves distant, very small. Fl. Mq. 57.—Miquelon Island.

598. *Hypnum occidentale* S. & L.—Plants densely cespitose; tufts intricate, depressed, bright green, sub-sericeous; stem much divided, filiform, sub-repent; branches prostrate, strongly branched, branchlets filiform or attenuate or thickish julaceous; leaves erect spreading when moist, imbricate when dry, ovate or lingulate, rather obtuse, concave, border denticulate above; costa thick, reaching middle; cells minute, oval-rhombic, alar quadrate, smaller; perichaetial leaves erect from a sheathing base, upper broadly ovate, uppermost lanceolate, obtusely short acuminate, reflexed; dioicous: capsule oblong-ovate to oblong-cylindric, inclined, sub-cernuous, exannulate, dilated below mouth when empty; seta smooth, scarcely 1 inch long; sub-cygnous; peristome teeth broadly lamellose within, segments entire, punctulate, cilia 2, shorter; lid long conic, acute, shortly oblique rotate. Sull. Icon. Musc. Suppl. 105. pl. 81. 1874.—Roots and base of trees, Oregon.

599. *Hypnum scorpioides Miquelonense* R. & C.—Robust, submersed; tufts black, apical leaves only golden yellow passing to red; stems 15-35 cm.: leaves of principal branches sub-imbricate, slightly secund, bluntly acuminate; leaves of secondary branches rather crowded, erect, falciform and flexuous at point, narrower, oblong, long and narrowly acuminate, needle-pointed, sometimes toothed at apex; costa sometimes short bifurcate, sometimes single, feeble, reaching or surpassing middle; median

cells usually sinuous, somewhat pitted. Ren. in Husn. Muscol. Gall. 394. 1894.—Miquelon Is.

600. *Hypnum incurvatum* Schrad.—Stem 2-4 cm., delicate, creeping, irregularly branching; tufts small, depressed, silky, green or slightly yellowish; leaves erect spreading, sub-secund, more or less arcuate, oblong-lanceolate, long-acuminate, entire or distantly toothed at apex; costa none or very short and faint, or bifurcate; cells of angles quadrate, middle cells short, 6-8 times as long as wide: monoleucous: two inner perichastial leaves sheathing, abruptly and narrowly acuminate, superficially denticulate at summit, not plicate, faintly costate: capsule horizontal, sometimes oblique, oblong or almost cylindric, arcuate, contracted below mouth; lid conic, short rostrate; annulus large. Husnot, Musc. Gall. 399.—Newfoundland.

601. *Hypnum dilatatum* Wils.—Stems 2-10 cm., prostrate, ascending, long denudate at base; tufts depressed, rather rigid: leaves sub-secund or secund, concave, sub-orbicular, obtuse or shortly and obtusely apiculate, narrowed at base, slightly denticulate at apex; costa bifurcate, very short or scarcely $\frac{1}{4}$ the leaf; cells of basilar angles large, hexagonal-rectangular, usually orange, forming rather distinct auricles, the rest longer than in *H. molle*: inner perichastial leaves sheathing, plicate: monoleucous: capsule oblique or horizontal, oblong; operculum convex-conic. Husn. Muscol. Gall. 413. *H. molle* Br. & Sch., not Dicks.—Greenland; Canada; Rocky Mts.; British Columbia; Idaho; Eastern States.

602. *Hylocomium squarrosum calvescens* (Wils.) Husnot.—Plants a little more robust, branches more numerous: stem leaves larger above, more distinctly denticulate, slightly plicate; leaf cells, pedicel and capsule similar to type. Husnot, Musc. Gall. 425. *H. calvescens* Wils.—On rocks: Nova Scotia; British Columbia.

603. *Hylocomium triquetrum Californicum* Ren. & Card.—Very robust: leaves strongly rugose undulate, strongly papillose above: capsule short. Bot. Gaz. 15: 61. 1890.—California.

- Micromitrium, 157, 176.
 Mielihoferia, 163, 211, 305.
 Mnium, 165, 219, 321.
 Myrinia, 167, 229, 333.
 Myurella, 167, 229.

 Neckera, 166, 167, 226, 331.

 Octoblepharum, 159.
 Edipodium, 159, 301.
 Oligotrichum, 164, 222, 324.
 Oreoweisia, 161.
 Orthothecium, 167, 168, 232, 336.
 Orthotrichum, 160, 164, 203, 293.

 Paludella, 165.
 Phacomitrium, 158.
 Phascum, 158, 177, 260.
 Philonotis, 165, 211, 305.
 Physcomitrella, 158.
 Physcomitrium, 159, 209, 301.
 Plagiothecium, 169, 241, 351.
 Platygyrium, 167, 168, 334.
 Pleuridium, 169, 177, 261.
 Pogonatum, 164, 222, 324.
 Polytrichum, 164, 223, 324.
 Pottia, 159, 162, 188, 274.
 Pseudoleskea, 168, 232, 337.
 Pterigynandrum, 166, 228, 332.
 Pterogonium, 166, 228.
 Pterigophyllum, 168.
 Ptychomitrium, 160, 201.
 Pylaisia, 166, 168, 230, 334.

 Raphidostegium, 169, 239, 350.
 Rhacomitrium, 160, 169, 289.

 Rhabdoweisia, 162, 179.
 Rhizogonium, 165.
 Rhynchostegium, 169, 240.

 Scleropodium, 169, 237, 348.
 Scouleria, 158, 197, 283.
 Seligeria, 161, 188, 273.
 Spharangium, 156, 176.
 Sphagnum, 157, 170, 251.
 Splachnum, 160, 206.
 Stereophyllum, 168.
 Syrrhopodon, 162, 207.

 Tayloria, 160, 208, 300.
 Tetraxis, 163, 207.
 Tetraplodon, 162, 208.
 Thamnum, 168, 240, 351.
 Thelia, 166, 229, 333.
 Thuidium, 168, 233, 339.
 Timmia, 165, 221, 323.
 Trematodon, 161, 180, 265.
 Trichodon, 161, 187, 272.
 Trichostomum, 161, 162, 163, 190, 275.
 Tripterocladium, 168, 234, 340.

 Ulota, 160, 164, 202, 292.

 Voitia, 158, 261.

 Webera, 165, 211, 306.
 Weisia, 162, 179, 262.

 Zieria, 165, 219.
 Zygodon, 158, 164, 201, 291.

a, 162, 179.
i, 165.
ium, 168, 240.

a, 169, 237, 348.
s, 197, 283.
l, 188, 273.
n, 158, 176.
57, 170, 251.
160, 208.
m, 168.
, 162, 207.

, 208, 300.
3, 207.
162, 208.
168, 240, 351.
229, 333.
88, 233, 339.
221, 323.
161, 180, 265.
61, 187, 272.
m, 161, 162, 163, 190, 275.
ium, 168, 234, 340.

64, 202, 292.

261.

211, 306.
179, 262.

219.
3, 164, 201, 291.



